


NORTHWEST ARKANSAS REGIONAL EMS PROTOCOLS



2023 Revisions

TABLE OF CONTENTS.....	b - e
INTRODUCTORY STATEMENT.....	f
PARTICIPATING AGENCIES.....	g
2019 DEPARTMENT MEDICAL DIRECTORS & SIGNATURE.....	h

SECTION ONE - PROTOCOLS

GENERAL PROTOCOLS

UNIVERSAL PATIENT CARE.....	1
CHEMICAL EXPOSURE (HAZMAT).....	2
CHEMICAL SEDATION FOR VIOLENT PATIENT.....	3
PAIN MANAGEMENT.....	4
SPINAL RESTRICTION.....	5
VASCULAR ACCESS.....	6

RESPIRATORY

OXYGEN ADMINISTRATION.....	7
GENERAL AIRWAY MANAGEMENT.....	8
ADVANCED AIRWAY MANAGEMENT.....	9
PHARMACOLOGICAL ASSISTED INTUBATION (PAI).....	10
ALLERGIC REACTION—ANAPHYLAXIS.....	11
ASTHMA.....	12
COPD.....	13
PULMONARY EDEMA.....	14
STRIDOR (UPPER AIRWAY OBSTRUCTION).....	15

CARDIAC / RESUSCITATION

ACUTE CORONARY SYNDROMES.....	16
AUTOMATED DEFIBRILLATION/ CPR.....	17
DEATH/WITHHOLDING RESUSCITATION.....	18
DO NOT RESUSCITATE.....	19
TERMINATION OF RESUSCITATION PROTOCOL.....	20
ASYSTOLE.....	21
BRADYCARDIA.....	22
CARDIAC ARREST.....	23
NARROW COMPLEX TACHYCARDIA.....	24
POST RESUSCITATION.....	25
PULSELESS ELECTRICAL ACTIVITY.....	26
VENTRICULAR FIBRILLATION / PULSLESS V-TACH.....	27
WIDE COMPLEX TACHYCARDIA.....	28

MEDICAL/ENVIRONMENTAL EMERGENCIES

ABDOMINAL PAIN.....	29
ACUTE STROKE—CVA.....	30
ALTERED LEVEL OF CONSCIOUSNESS/MENTATION.....	31
BITES, STINGS, and ENVENOMATIONS.....	32
DIABETIC EMERGENCIES.....	33
HEAT EMERGENCIES.....	34
HYPERTENSIVE CRISIS.....	35
HYPOTENSION—SYMPTOMATIC.....	36
HYPOTHERMIA.....	37

NAUSEA—VOMITING—DIARRHEA.....	38
NEAR DROWNING/DROWNING	39
OVERDOSE/TOXIC EXPOSURE.....	40
SEIZURE.....	41
SEPSIS.....	42
SYNCOPE.....	43
TRAUMA	
BURNS.....	44
CHEST TRAUMA	45
DESTINATION PROTOCOL.....	46
GLASGOW COMA SCORE.....	48
DIVING EMERGENCIES (SCUBA)	49
EXTREMITY TRAUMA.....	50
SHOCK- TRAUMA	51
TASER REMOVAL	52
TRAUMA.....	53
TRAUMATIC BRAIN INJURY	54
OBSTETRICS	
CHILDBIRTH.....	55
COMPLICATIONS IN CHILDBIRTH (Multiples/Prolapsed Limb/Nuchal & Prolapsed Cord/Pre-Term).....	56
COMPLICATIONS IN CHILDBIRTH (Shoulder Dystocia/Breech)	57
ECLAMPSIA-PREECLAMPSIA.....	58
POST PARTEM CARE.....	59
NEWBORN RESUSCITATION	60
PEDIATRIC	
UNIVERSAL PATIENT CARE - <i>Pediatric</i>	61
AIRWAY MANAGEMENT - <i>Pediatric</i>	62
ALLERGIC REACTION – ANAPHYLAXIS - <i>Pediatric</i>	63
ASTHMA - PEDIATRIC	64
ASYSTOLE/PULSELESS ELECTRICAL ACTIVITY - <i>Pediatric</i>	65
BRADYCARDIA - <i>Pediatric</i>	66
DIABETIC EMERGENCIES - <i>Pediatric</i>	67
MULTIPLE TRAUMA - <i>Pediatric</i>	68
OVERDOSE/TOXIC EXPOSURE - <i>Pediatric</i>	69
POST RESUSCITATION - <i>Pediatric</i>	70
SEIZURE - <i>Pediatric</i>	71
STRIDOR(UPPER AIRWAY OBSTRUCTION) - <i>Pediatric</i>	72
TACHYCARDIA - <i>Pediatric</i>	73
VENTRICULAR FIBRILLATION/PULSELESS V-TACH - <i>Pediatric</i>	74

SECTION 2 - MEDICATIONS - Referenced in Protocols

ADENOSINE (ADENOCARD).....	76
ALBUTEROL (PROVENTIL, VENTOLIN)	77
AMIODARONE (CORDARONE).....	78
ASPIRIN (ASA, BAYER, ECOTRIN, ST. JOSEPH, OTHERS).....	79
ATROPINE SULFATE	80
CEFAZOLIN (ANCEF).....	81
CETACAINE.....	82
DEXAMETHASONE (DEXAMETHAZONE)	83
DEXTROSE 10%.....	84
DIAZEPAM (VALIUM).....	85
DILTIAZEM (CARDIZEM)	86
DIPHENHYDRAMINE (BENADRYL)	87
DOPAMINE (INTROPIN)	88
EPINEPHRINE (ADRENALIN) ADULT SHEET	89
ETOMIDATE (AMIDATE).....	90
FENTANYL CITRATE (SUBLIMAZE)	91
FUROSEMIDE (LASIX)	92
GEODON (ZIPRASIDONE).....	93
GLUCAGON	94
HALOPERIDOL LACTATE (HALDOL)	95
HEPARIN SODIUM	96
HYDROXYZINE (VISTARIL, ATARAX, APRESOLINE)	97
IPRATROPIUM (ATROVENT)	98
KETAMINE (KETALAR).....	99
KETOROLAC TROMETHAMINE (TORADOL).....	100
LABETALOL (NORMODYNE, TRANDATE)	101
LEVALBUTEROL (XOPENEX)	102
LIDOCAINE (XYLOCAINE).....	103
LORAZEPAM (ATIVAN)	104
MAGNESIUM SULFATE.....	105
METHYLPREDNISOLONE (SOLU-MEDROL).....	106
MIDAZOLAM HYDROCHLORIDE (VERSED)	107
MORPHINE SULFATE	108
NALOXONE (NARCAN)	109
NITROGLYCERIN (NITROSTAT AND OTHERS).....	110
ONDANSETRON HYDROCHLORIDE - ZOFRAN	111
OXYMETAZOLINE HYDROCHLORIDE (AFRIN)	112
OXYTOCIN (PITOCIN)	113
PROCAINAMIDE (PRONESTYL)	114
PROMETHAZINE (PHENERGAN).....	115
RACEMIC EPINEPHRINE	116
ROCURONIUM BROMIDE	117
SODIUM BICARBONATE	118
SUCCINYLCHOLINE (ANECTINE).....	119
TETRACAINE (PONTACAINE).....	120
THIAMINE (BETAXIN).....	121
TRANEXAMIC ACID (TXA).....	122
VECURONIUM (NORCURON).....	123
VERAPAMIL (ISOPTIN).....	124

SECTION 3 GUIDELINES (HOW TO PERFORM SKILL LISTED IN PROTOCOLS)

12 LEAD EKG	126
RIGHT SIDED AND POSTERIOR LEAD EKG.....	127
APGAR.....	128
BLOOD DRAW.....	129
BLOOD GLUCOSE ANALYSIS	130
BLOOD PRODUCTS.....	131
CAPNOGRAPHY -ELECTRONIC & COLORIMETRIC / ETCO2.....	133
CARBON MONOXIDE OXIMETRY	134
CHEST DECOMPRESSION.....	135
CONTINUOUS POSITIVE AIRWAY RESSURE (CPAP).....	136
CRICOTHYROTOMY--NEEDLE.....	137
CRICOTHYROTOMY--SURGICAL.....	138
CRIME SCENE.....	139
DECONTAMINATION - EMERGENCY.....	140
ELECTRICAL THERAPY.....	141
GLASCOW COMA SCALE.....	142
INTRAOSSEOUS ACCESS HUMERAL HEAD.....	143
INTRAOSSEOUS ACCESS TIBIAL.....	144
INTUBATION--NASOTRACHEAL (BLIND).....	145
INTUBATION--OROTRACHEAL	146
IV ACCESS-EXISTING CENTRAL VENOUS LINES	147
IV ACCESS—EXTERNAL JUGULAR.....	149
KING AIRWAY.....	150
LEFT VENTRICULAR ASSIST DEVICE (LVAD).....	151
MEDICATION ADMINISTRATION.....	152
MIST REPORT.....	157
ORTHOSTATIC VITAL SIGNS.....	158
PCR SHORT FORM.....	159
PHYSICIAN ON SCENE.....	160
PULSE OXIMETRY.....	161
RESTRAINTS.....	162
RULE OF NINES.....	163
SAFETY EQUIPMENT RREMOVAL.....	164
STROKE SCREEN BEFAST.....	165
TOXICOLOGICAL SYNDROMES.....	166
TRAUMA ALERT.....	167
TRAUMA SCORE ADULT	168
TRIAGE.....	169

INTRODUCTORY STATEMENT

The following protocols, guidelines, and instructional material were developed as a collective effort by a group of dedicated, charitable, and knowledgeable paramedics, EMS educators, and Medical Directors of the EMS agencies of Northwest Arkansas. These individuals, who recognized a need for a "Standard of Excellence" voluntarily formed a regional Task Force which initiated, researched and reviewed the following algorithms and have chosen to implement them as guidelines and as a basis for the standard of care in our area.

This collaborative effort provides a dynamic document that is based on national and state standards of care to include but not limited to:

- Arkansas Department of Health
- American Heart Association—Advanced Cardiac Life Support
- American Academy of Pediatrics—Pediatric Advanced Life Support American College of Surgeons—
- Advanced Trauma Life Support American College of Surgeons
- National Association of Emergency Medical Technicians
- National EMS Scope of Practice and Instructional Guidelines
- American College of Emergency Physicians

This protocol booklet is meant to be reviewed continuously and updated as national, state and regional standards change and scientific research and literature support.

The purpose of the following algorithms is multifaceted. The primary purpose is to establish a foundation and a minimum standard of care for the pre-hospital care delivered in our area. The intent is to provide current, well-researched, and accepted standards with the ultimate goal of minimizing the morbidity and mortality of our patients and to provide guidelines for the treatment of specific emergency conditions in the pre-hospital setting. This is best served by active EMS Medical Directors and dedicated Paramedics/EMT s supported by continued education, review and quality improvement, and continuous pursuit of excellence.

Although no document can specifically address every possible variation of injury or disease, this manual provides a foundation for the acute care of the patients we serve. The education, experience, and judgment of the pre-hospital provider should be recognized as the paramount part of sound clinical decision-making processes regarding pre-hospital care. The complexity of emergency medicine and the pre-hospital setting require a team-approach use of every appropriate, accepted, and available resource to provide optimal patient care. In many cases, that resource is On-line Medical Control for consultation, advice, guidance, and authorization or modification for treatment not specifically addressed in this manual. The specifics of this requirement are to be determined by the Medical Director responsible for that particular EMS service and the paramedics they oversee, and is intentionally not addressed in this manual, for that reason.

The departmental policies are the responsibility of that individual agency and Medical Director due to the specifics of that EMS agency, but are encouraged to support the premise of regional care and collective effort these guidelines were founded on.

The provision of emergency care does not, and should not, occur in isolation. It requires many individuals and organizations working together towards a common goal—optimizing our patient's clinical outcome. We hope that the efforts provided by this founding Task Force and the resultant work provides a basis for the future development of a regional EMS approach to the "Standard of Excellence" we strive for in the care of our patients and the people of Northwest Arkansas.

Northwest Arkansas EMS Regional Protocol Participating Agencies

<i>Bella Vista Fire</i>	<i>Bentonville Fire</i>	<i>Eureka Springs Fire</i>
<i>Franklin County EMS</i>	<i>Gravette EMS and Fire</i>	<i>Jacksonville Fire Department</i>
<i>Johnson Regional EMS</i>	<i>Lowell Fire</i>	<i>Madison County EMS</i>
<i>North East Benton County (NEBCO) Fire and EMS</i>	<i>Northwest EMS</i>	<i>Ozark EMS</i>
<i>Pea Ridge EMS</i>	<i>PULSE EMS</i>	<i>Rogers Fire</i>
<i>Sebastian County EMS</i>	<i>Siloam Springs Fire</i>	<i>Southern Ambulance</i>
<i>Springdale Fire</i>	<i>Total Life Care (TLC)</i>	<i>Washington County EMS Authority (CEMS)</i>

Northwest Arkansas Regional EMS Protocols

All rights reserved. No part of the contents of this book may be reproduced or transmitted in any form or by any means without the written permission of the publisher.

Use of this document or its contents as guidelines or protocols for non-participating EMS providers or services is prohibited without the written permission of the publisher.


Use of these protocols – Protocols in brackets with grey on the sides are Advanced Procedures Utilized by paramedics only.



Non Grey sides are skill sets for all levels of licensure:



All Levels of Licensure



Northwest Arkansas EMS Regional Protocol

Department Medical Directors

Department	Name of Medical Director
Bella Vista Fire	Brad Johnson
Bentonville Fire	Brad Johnson
Centerton Fire Department	Brad Johnson
Eureka Springs Fire	Greg Kresse
Franklin County EMS	Jason Richey
Gravette EMS	Donald Tribbey
Jacksonville Fire Department	Darren Flamik
Johnson Regional EMS	James B Cotner
Lowell Fire	Mark Rucker
Madison County EMS	Burdge Green
NEBCO	Brad Johnson
Northwest EMS	Joe Olivi
Ozark EMS	William C Pittman
Pea Ridge EMS/Fire	Brad Johnson
Rogers Fire	Brad Johnson
Sebastian County EMS	Michael Hillis
Siloam Springs Fire	Gary Brooks
Southern	Chadwick Simpson & Don Ball
Springdale Fire	Mark Rucker
Total Life Care	James Holden
Washington County EMS	Dalton Lee Gray

MEDICAL DIRECTOR 2019- Revisions

As Medical Director for: _____

I have reviewed the Revised Northwest Regional Protocol Manual for 2019 and approve it for use. Portions that are not used or changed have been amended and my signature attached to that amendment page.

Signature:

_____ Date

UNIVERSAL PATIENT CARE

Scene Survey

- Assess the environmental conditions
- Mechanism of illness or injury
 - consider spinal precautions
- Considerations as indicated for airborne or droplet isolation
- If hazardous conditions are present (such as swift water, hazardous materials, electrical hazard, or confined space) contact an appropriate agency before approaching the patient. Wait for the designated specialist to secure the scene and patient as necessary.

Form a **General Impression** of the patient's condition &
Conduct an appropriate Adult or Pediatric Assessment

Primary Survey

General Airway Protocol

Vital signs:

- Stable patients should have a documented assessment, and trending vital signs at a minimum of 15-minute intervals.
- Unstable patients should have a documented assessment, and trending vital signs at a minimum of 5-minute intervals.
- Minimum vitals signs are:
 - Respirations
 - Pulse
 - Blood Pressure
 - Temperature
 - Pulse Oximetry
 - Blood Glucose
 - End Tidal Capnography (As indicated)
 - Cardiac Monitor (As indicated)

Vascular Access Protocol

Does the patient fit a specific protocol?	YES	Proceed To An Appropriate Protocol
	NO	Contact Medical Control As Indicated

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• A patient care report (PCR) must be completed for every patient contact, and a report must be left in ER	

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while **Blue font identifies an ALS skill/treatment or consideration.**

UNIVERSAL PATIENT CARE

prior to leaving ER- either short form or completed encounter form. • Pediatric patients (for these protocols) are patients who are from birth to adolescence (puberty). • The initial assessment must be appropriate to the patient's condition, mechanism of injury, and severity of illness.	
--	--

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

CHEMICAL EXPOSURE (HAZ-MAT)

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
- Identify substance - Type of exposure (ingestion, inhalation, contact) - Length of exposure - Quantity of substance/size of incident - Number of patients - Signs and symptoms of responders or others exposed - Past medical history - Other Trauma - Fire Department HAZMAT response - Decontamination if required	- Burns, pain, swelling - Eye irritation (burning, tearing, redness) - Dyspnea - CNS effects (headache, dizziness, altered mental status, seizure, coma) - Cardiac dysrhythmia - Nausea/Vomiting	- Trauma - Thermal burns - Cardiac - Respiratory (asthma) - Other medical (history) - Anxiety (hyperventilation) - Psychological - Overdose/Poisoning

Universal Patient Care Protocol

Remove patient from exposure area

Emergency Decontamination Procedure

- Notify Receiving Hospital

Type of Exposure

- Consider all, treat simultaneously

<u>Skin Contact</u>	<u>Inhalation</u>	<u>Ingestion</u>	<u>Burns</u>
If EYE involvement, remove contact lenses flush continuously with Normal Saline in each affected eye. Consider using Tetracaine Remove clothing and/or expose the area Flush the area with water or Normal Saline for 10 - 20 minutes	Fresh Air/ Oxygen Airway Management Protocol Monitor EKG, 12-Lead, if available, Proceed to Appropriate Protocol (Respiratory, Cardiac, Altered Mental Status, etc.)	Proceed to Overdose/Toxic Exposure Protocol	Proceed to Burn Protocol

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

CHEMICAL EXPOSURE (HAZ-MAT)

Contact Medical Control As Needed

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• 1st arriving units must perform a risk assessment. Situations involving gasses and vapors noticeably affecting victims, bystanders, and responders require SCBA at a minimum to remove patients from the area. • The act of removing patients from the exposure area reduces exposure to the patient and caregiver substantially. Do not treat patients in the hazard area. • Emergency decontamination will reduce the risk still further and eliminate almost all risks to the caregiver. Emergency Decontamination is the most important treatment when the chemicals are causing the symptoms. • Exam: ABCs, vital signs, mental status, skin, HEENT, neck, heart, lungs, abdomen, back, extremities, neurological. • Receiving hospitals must be notified early of chemical name(s), type of exposure, decontamination performed, and the number of patients. Notify with the information you have and update as newer information is received. • Medical control may order a high dose of Atropine (2-5 mg) for Organophosphate poisoning. • Reference the Emergency Response Guide (ERG)	• Burn Protocol • Airway Management • Overdose Toxic Exposure • Atropine

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Chemical Sedation For Violent Patient

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• Drug use • Hypoxia • Severe Epilepsy • Anxiety • Uremia • Psychiatric • Trauma • Poisoning • Acidosis • Stroke • Overdose • Shock • Diabetes	• Elevated HR • Uncontrolled Rage • Disorientation	• Head Injury • Hypoxia • Tumor • Hypo/hyperglycemia • Electrolyte imbalance • Hypertensive Crisis

Universal Patient Care Protocol

Is the Patient using physical force with the intent to harm themselves or another person?

<u>Can Police Restrain The Patient For A Reasonable Period Of Time?</u>		
	No	Retreat and stage
	<u>Yes</u> Select one of the medications and use until sedated or maximum dosage is administered. If no success, contact medical control for possible additional drug(s).	Haldol Or Droperidol Or Geodon/Ziprasidone Or Midazolam Or Ketamine Or Lorazepam Or Diazepam
	Consider reducing the amount of medication initially for patients who are elderly, pediatric, or have co-morbidity	

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Chemical Sedation For Violent Patient

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• Definition of <u>VIOLENT PATIENT</u>: Any patient using physical force with the intent to harm themselves or another person. • Definition of <u>EXCITED DELIRIUM</u>: Combination of delirium, psychomotor agitation, anxiety, hallucinations, speech disturbances, disorientation, violent/ bizarre behavior, insensitivity to pain, hyperthermia, and increased strength. Can be a result of cocaine, crack cocaine, methamphetamine, amphetamines, and other stimulant agents. • IV access should be accomplished prior to chemical sedation whenever possible. If not possible due to safety concerns, obtain as soon as possible after chemical sedation. • Consider physically restraining the patient after chemical sedation. • Constantly reassess any chemically or physically restrained patient for asphyxiation. DO NOT transport prone. • ETCO2 monitoring for any sedated patient if available. • Avoid the use of Ketamine with patients who have a psychiatric history. ○ Consider Contacting Medical Control prior to the administration of Ketamine for these patients.	Haldol Or Droperidol Or Geodon/Ziprasidone Or Midazolam Or Ketamine Or Lorazepam Or Diazepam

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while **Blue font identifies an ALS skill/treatment or consideration.**

Pain Management

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• Age • Location • Duration • Severity on 1-10 Scale • Past Medical History • Medications • Drug Allergies	• Severity • Quality • Radiation • Relation to Respiration • Increased with Palpation	

<u>Universal Patient Care Protocol</u>
--

Acute Coronary Syndrome with Chest Pain	Pain management as need for persistent ACS chest pain following administration of NTG, monitoring BP >90mmHg
Pain Related to IO Use	See also IO Guideline Lidocaine- administer slowly prior to IO bolus or flush on alert/ conscious patient
General/Other Pain	Refer to specific protocol Consider treatment with appropriate medication • Fentanyl • Morphine • Ketamine • Ketorolac • Acetaminophen • Ibuprofen • Dilaudid Consider adding Benzodiazepine; B/P based (max dose is pressure based) • Diazepam • Midazolam • Lorazepam

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Pain Management

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• Use of Pulse Oximetry and/or capnography should be considered when using pain medication • Use of a Benzodiazepine may be indicated for cocaine and or meth overdose with Chest pain. • Ensure an exam and, at minimum, the topics below are assessed before administering any pain medication. ○ Mental status ○ Area of pain ○ Neuro ○ Vital signs • Toradol (Ketorolac Tromethamine) has been shown to be effective when used for kidney stone pain and migraine pain.	Nitroglycerin Lidocaine Fentanyl Morphine Ketamine Diazepam Midazolam Lorazepam Acetaminophen Ibuprofen Dilaudid Ketorolac

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Spinal Restriction

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• Mechanism of Injury • Unknown history with unresponsiveness	• Neck pain/back pain • Tenderness, crepitus, or deformity on palpation of spine • Numbness, tingling-paresthesia • Limited range of motion	• Muscular-lateral neck or back pain

Universal Patient Care Protocol

Alert, Oriented and Neurologically intact, and able to communicate?	NO	Spinal Restriction
---	----	--------------------

Following the assessment chart below with the identification of no complaint,
NO SPINAL RESTRICTION REQUIRED

Evidence of major injury which may distract patient's awareness of pain?	YES	Spinal Restriction
Evidence of intoxication or Mental impairment?	YES	Spinal Restriction
Complaining of Neck Pain?	YES	Spinal Restriction
Pain on palpation of spinous processes of Cervical, Thoracic or Lumbo-Sacral Spine?	YES	Spinal Restriction
Neck pain to patient's range of motion?	YES	Spinal Restriction

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Spinal Restriction

Spinal Restriction:

If the patient is ambulatory, place an appropriately sized C Collar & Position the patient directly on the Cot in the position of comfort, limiting movement of the spine during the process. Utilize the cot securing straps/system to secure the patient. Supine/Prone Patients should be moved to the cot utilizing an appropriate device. Patients who are stable, alert, and without neurological deficits who are sitting or standing may be allowed to self-extricate to the ambulance cot after placement of a cervical collar - limiting movement of the spine during the process. If a Long Spine Board or Scoop Stretcher, etc. is used for extrication or patient movement, the patient should be taken off the device (if time allows) & Placed directly on the EMS Stretcher

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• Refer to Safety Equipment Removal Guidelines if the patient is wearing Safety gear • If Patient's condition may be worsened by spinal restriction, then spinal restriction may not be prudent for that patient. • The decision to NOT implement spinal restriction in a patient is the responsibility of the EMS provider; if there is a concern, perform spinal restriction. • The decision not to apply spinal restriction should be thoroughly documented on the patient care report. • Patient's range of motion should not be assisted. The patient should touch their chin to chest, extend their neck (look up) and turn their head from side to side (shoulder to shoulder) without pain. • Major injuries which may distract a patient's awareness of pain include pelvic fracture, femur fracture, extensive burns or soft tissue injury, acute abdomen, or significant chest injury. • Proper restraints on the EMS cot should be adhered to for all spinal injured patients. Use padding when necessary to limit movement. • Consider leaving children in their car seats unless they need immediate treatment or the car seat is damaged. • Remove seriously injured children from the child seat, or if it is potentially damaged in the crash. • Seriously injured children may require spinal restriction.	

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while [Blue font identifies an ALS skill/treatment or consideration.](#)

Vascular Access

Universal Patient Care Protocol

Assess Need for IV Access:

Patient may become unstable en route or at ED

Patient may require blood lab values at ED

Patient requires IV fluids and /or medications

Attempt intravenous access of peripheral extremities with an appropriately sized IV catheter.

Unsuccessful X2

Or

Appropriate landmarks for peripheral access are unavailable?

Consider

<u>Stable</u>	<u>Unstable</u>
Transport without IV	IO Placement
	External Jugular (puberty or >)
	Access of existing central venous lines

<u>Notes</u>	<u>Associated Protocols/Resources</u>
- When performing I.O.-Proximal Humeral or Tibial Placement may be used. - When performing I.O.-Large Adult needle is necessary for Proximal Humeral placement - In cases of severe illness or injury requiring immediate fluid or drug administration, an IO may be considered prior to peripheral IV attempts if IV assessment is unlikely or impossible - Consider pain management for conscious patients receiving IO placement. - In post-mastectomy patients, avoid IV, blood draw, or injection in the arm on the affected side.	

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Vascular Access

- | | |
|---|--|
| <ul style="list-style-type: none">• In the setting of cardiac arrest, any preexisting dialysis shunt may be used but should otherwise be avoided.• Lower extremity IV sites are discouraged in patients with vascular disease or diabetes.• Any venous catheter which has already been accessed prior to EMS arrival may be considered.• Proximal humeral IO placement results in better circulation of medications, especially during cardiac arrest. | |
|---|--|

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Oxygen Administration

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• COPD • Asthma • Emphysema • Trauma • Obstruction	• Poor perfusion • Low O2 Saturation • Inadequate Ventilation • Skin color changes • LOC changes	• Chronic Disease • Physical obstruction • Shock • Drug Overdose • CO Poisoning

Indications:

- The overall goal of oxygen therapy is to avoid tissue hypoxia
- The most common indications for oxygen saturation in the acute setting are the presence of arterial hypoxemia or a failure of the oxygen-hemoglobin transport system
- Arterial hypoxemia is defined as an oxygen saturation of less than 88% and may result from impaired gas exchange in the lung, inadequate alveolar ventilation or a shunt that allows venous blood into the arterial circulation
- A failure of the oxygen-hemoglobin transport system can result from a reduced oxygen carrying capacity in the blood (e.g. Anemia, carbon monoxide poisoning) or reduced tissue perfusion (e.g. Shock)
- If accuracy of Pulse ox is suspected and hypoxia is suspected, or in doubt – administer oxygen

Universal Patient Care Protocol

Is there a need for an Advanced Airway?	NO	Proceed to: "General Airway Protocol" AND/OR "Oxygen Admin and Titration Protocol"
Airway Obstructed	YES	AHA BLS Guidelines for "Foreign Body Airway Obstruction"
Apnea, Agonal Respirations and/ or no Gag Reflex?	YES	Direct Laryngoscopy
Can the patient swallow, is there a need to secure the airway / intubate the patient	YES / NO	Consider/Proceed to PAI Protocol if applicable

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while **Blue font identifies an ALS skill/treatment or consideration.**

Oxygen Administration

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• Continue with 100% O₂ in any patient with suspected carrying capacity: • Known or expected anemia (ie. Trauma, GI Bleed, etc) • Presence of Toxins such as Carbon monoxide, Methemoglobin, or cyanide. (Note Paraquat poisoning is a contraindication to oxygen unless below 88%) • Titrate O₂ target range of 88 – 92% in patients with exacerbation of COPD, Asthma, and Chronic neuromuscular disease with difficulty breathing. • No supplemental O₂ is recommended for cardiac and CVA patients with SpO₂ ≥ 94%. If O₂ is required to achieve SpO₂ ≥ 94% titrate to ≤ 96% • Hyperoxia resulting from high FiO₂ administration producing saturations higher than 94-96% (or PaO₂ > 100) in acute conditions can cause structural damage to the lungs, as well as potentially cause increases in free radicals and post-reperfusion tissue damage. In addition, patients who are chronically hypoxic (COPD, ALS, MS, etc.) have shifted their oxygen dissociation curve and require lower oxygen saturations. • With prolonged oxygen therapy there is an increase in blood oxygen level, which may suppress peripheral chemoreceptors, depress ventilation/perfusion balance (V/Q) and cause an increase in dead space to tidal volume ratio and increase in PCO₂.	

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

General Airway Management

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• Trauma • Head Injury • CVA • Asthma • COPD • Known difficult airway • Facial fractures • Pulmonary edema • Respiratory Distress	• Hoarseness • Limited neck movement • Limited mouth opening • Short thyromental distance • Short heavy neck • Receding mandible/overbite • Large swollen tongue • Obesity • Long incisors	• LOC • Airway injury • Airway swelling • Burns • Foreign body • Epiglottitis

Universal Patient Care Protocol

Airway, Ventilation, and Level of Consciousness Adequate	YES	Proceed to: "Oxygen Admin and Titration Protocol"
--	------------	--

Airway, Ventilation, and Level of Consciousness Adequate	NO	Manual Airway Maneuver
		Airway Adjunct (i.e. OPA, NPA)
		Bag-Valve-Mask Ventilation
		Suctioning as needed
		EtCo2 Monitoring (if available)

Airway Obstructed	YES	AHA BLS Guidelines for "Foreign Body Airway Obstruction"
-------------------	------------	--

Need for Advanced Airway Procedures	YES	Proceed to "Advanced Airway Procedures" Protocol
-------------------------------------	------------	--

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

General Airway Management

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• Keep it simple – use progressively invasive maneuvers only as necessary • Pulse Oximetry & Capnography should be used for all airway/ventilation problems • Maintain spinal precautions , neutral alignment when trauma suspected	

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Advanced Airway Management

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• Trauma • Head Injury • CVA • Asthma • COPD • Known difficult airway • Facial fractures • Pulmonary edema • Respiratory Distress	• Hoarseness • Limited neck movement • Limited mouth opening • Short thyromental distance • Short, heavy neck • Receding mandible/overbite • Large, swollen tongue • Obesity • Long incisors	• LOC • Airway injury • Airway swelling • Burns • Foreign body • Epiglottitis

Universal Patient Care Protocol

Is there a need for an Advanced Airway?	NO	Proceed to: "General Airway Protocol" AND/OR "Oxygen Admin and Titration Protocol"
---	-----------	--

Airway Obstructed	YES	AHA BLS Guidelines for "Foreign Body Airway Obstruction"
-------------------	------------	--

Apnea, Agonal Respirations and / or no Gag Reflex?	YES	Direct Laryngoscopy
--	------------	---------------------

Can the patient swallow, is there a need to secure the airway / intubate the patient	YES NO	Consider/Proceed to PAI Protocol Secure the airway as necessary
--	-------------------------	--

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• Document ET or Supraglottic Tube Placement at the transfer of the patient. • Keep it simple – Use progressively invasive maneuvers only when necessary.	

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Advanced Airway Management

- | | |
|---|--|
| <ul style="list-style-type: none">• Clinical End Tidal CO2 monitoring should be used with all advanced airways.• Pulse Oximetry is used for all airway/ventilation problems when circulation allows and is maintained above 90% at all times.• Maintain spinal precautions and neutral alignment when trauma is suspected.• Once ET or Supraglottic tube is placed and confirmed, apply cervical collar and maintain C spine immobilization to prevent displacement. | |
|---|--|

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Pharmacological Assisted Intubation (PAI)

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• Trauma • Head Injury • CVA • Asthma • COPD • Known difficult airway • Facial fractures • Pulmonary edema • Respiratory Distress	• Hoarseness • Limited neck movement • Limited mouth opening • Short thyromental distance • Short heavy neck • Receding mandible/overbite • Large swollen tongue • Obesity • Long incisors	• LOC • Airway injury • Airway swelling • Burns • Foreign body • Epiglottitis

<u>Indication</u>	<u>Contra-Indications</u>
• Altered mental status where the loss of airway is inevitable • Airway compromise is a real possibility before or during transport • Breathing is not adequate and likely require intubation during transport • Patients with trauma injuries not able to protect their airway • Multi System trauma affecting adequacy of ventilations • Inability to maintain O2 Sat. of >90% with BVM and NPA/OPA • Anticipated clinical course	• (PAI) is not approved in patients < 8 YO in Arkansas • Do NOT attempt (PAI) if Cricothyrotomy would NOT be possible in the case of a failed (PAI) Attempt • Do not attempt (PAI) if ventilations and intubation would be difficult or impossible such as Epiglottis or partial airway obstruction

<u>Universal Patient Care Protocol</u>
--

<u>Pre-Oxygenation</u>
• Capnography monitored throughout event/procedure • Pre-Oxygenation Phase (100% Oxygen) 3-5 minutes prior to Laryngoscopy • Prepare and apply Apneic Oxygenation

<u>Induction</u>
• Administer Sedative / Anesthetic Agent

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Pharmacological Assisted Intubation (PAI)

<u>Crash Airway</u> (When appropriate)
Indications
Unconscious / Unresponsive need for paralytic administration to perform intubation
- Administer Paralytic - Increase apneic oxygenation to 10-15 lpm until confirmed intubation

<u>Laryngoscopy</u>
Perform intubation as necessary

<u>Post Intubation Management</u>
- Continued sedation with post-intubation sedative - Monitor, anticipate and treat pain for management/discomfort of intubation - Monitor for / anticipate hypotension - Consider long-acting paralytic only if sedation methods are not effective in keeping the patient sedated

<u>Notes</u>	<u>Associated Protocols/Resources</u>
- Document ET or Supraglottic Tube Placement at the transfer of the patient. - Keep it simple – Use progressively invasive maneuvers only when necessary. - Clinical End Tidal CO2 monitoring should be used with all advanced airways. - Pulse Oximetry is used for all airway/ventilation problems...when circulation allows and maintained above 90% at all times. - Maintain spinal precautions and neutral alignment when trauma is suspected. - In an attempt to define intubation in the emergent setting and to collect/record accurate data, this group shall define an intubation attempt as “the introduction of the laryngoscope with the intent of passing a ET tube through the cords.” It is strongly recommended that providers, prior to intubation, attempt to obtain a view of the oropharynx to identify concerns prior to an intubation attempt. If landmarks are identified and intubation can be accomplished, it should be done with a first pass success recorded. - After making adjustments following a single view, if the tube is placed with the second introduction of the laryngoscope, it should be recorded as a first pass success. All	

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Pharmacological Assisted Intubation (PAI)

additional attempts shall be recorded in successive order. • Once ET or Supraglottic tube is placed and confirmed, apply a cervical collar and maintain C spine immobilization to prevent displacement of the tube • Unsuccessful intubation attempts ○ Use a secondary device ASAP ○ Ventilation performed with BVM and adjunct placed/used as needed ○ In the event a patient cannot be ventilated by any other means perform a surgical airway as a means of last resort.	
--	--

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Allergic Reaction

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
- Known allergic reaction to bites, stings, food, medications, etc. - Possible ingestion of or contact with an allergen.	- Dyspnea, often with sneezing, wheezing, or coughing - Facial swelling - Urticaria - Abdominal cramps - Nausea, vomiting, diarrhea - Tachycardia - Hypotension	- Asthma - Pulmonary embolism - History of ACE Inhibitor use – angioedema - Capnography – determine if bronchospasm is present

Universal Patient Care Protocol

<u>Determine Reaction</u>	
Allergic Reaction - Mild symptoms (Urticaria, erythema, mild wheezing, etc.)	Anaphylaxis - Moderate & Severe (Hypoxemia, hypotension, history of allergic reactions, etc.)
Diphenhydramine	Epinephrine - Give early on- IM if needed. If still symptomatic administer Epinephrine IV - Alive patients should be administered a 1:100k ratio of Epinephrine by IV.
Bronchodilator Consider Ipratropium (Atrovent) or levalbuterol if Albuterol is Ineffective	Fluid Bolus 250-500cc, repeat bolus titrate to BP
Consider Epinephrine	Diphenhydramine
	Bronchodilator Consider Ipratropium (Atrovent) or levalbuterol if Albuterol is Ineffective
	Consider Decadron or Solu-Medrol

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Allergic Reaction

<u>Monitor and Observe for rebound effects</u>	
- Continued Hypotension? - Decreased peripheral perfusion? - Shock?	Consider - Glucagon - Vasopressors

<u>Notes</u>	<u>Associated Protocols/Resources</u>
- Consider Epinephrine IM, & diphenhydramine early in the allergic process, administration prior to histamine release will provide more rapid results. - When signs of histamine release are noted, the process is well underway and will require aggressive treatment. - Epinephrine has a short half-life and may require repeated doses. - Closely monitor patients for rebound signs and symptoms. - Any patient suffering from an allergic reaction should be evaluated by a physician.	Hypotension Protocol

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Asthma

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• Asthma • COPD: Emphysema, Bronchitis • CHF: Congestive Heart Failure • Home Oxygen use • Home Nebulizer Use • Medications: Steroids • Inhalation of Possible Chemical or biological agents • Exposure to chemical or biological agents	• Shortness of breath • Pursed-Lip breathing • Accessory muscle use, retractions, nasal flaring, fatigue • Inability to speak in sentences • Audible Wheezing or rhonchi • Fever, cough • Cyanosis • Lung sounds: ◦ Wet? ◦ Diminished? ◦ Bilaterally? ◦ Expiratory ◦ Wheezing?	• COPD • CHF, Pulmonary Edema • Anaphylaxis • Pneumonia • Pulmonary Embolus • Cardiac • Hyperventilation • Inhaled toxin • DKA • Pneumothorax

Universal Patient Care Protocol

Oxygen Administration Protocol

General Airway Management Protocol

Apply waveform Capnography

12 Lead ECG

Fluid Bolus

Bronchodilators

Consider Epinephrine

Consider Steroid

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Asthma

Consider Advanced Airway Protocol

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• If in severe Asthma/COPD distress, treatment may occur simultaneously with IV, ECG, and 12-Lead. Consideration of Mag Sulfate in the updraft, contraindications are ◦ Heart block or myocardial damage ◦ Hypermagnesia • IV administered Mag Sulfate should be diluted in 100ml fluid. Be aware of contraindications of Mag Sulfate • Pulse Oximetry & Capnography should be monitored continuously for all patients with respiratory distress and/or respiratory failure. • Remember: almost all cardiac problems produce some degree of respiratory distress. • Patients with a history of asthma, who have had prior hospitalization for asthma, and/or present with initial O2 saturations of <90% are at increased risk for the rapid decline in spite of initial improvement with your treatments. • A silent chest in the setting of severe respiratory distress is a pre-respiratory arrest sign. • Use extreme caution when administering epinephrine in patients who are >50 years of age, have a history of cardiac disease, or if the patient's heart rate is > 150. Epinephrine may precipitate cardiac ischemia. • Respiratory distress can be the result of metabolic acidosis from overdose and/or DKA, head injury, or trauma.	Magnesium Sulfate Drug Sheet

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

COPD

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• Asthma • COPD: ◦ Emphysema, Bronchitis • Congestive Heart Failure (CHF) • Home Oxygen Use • Medications: ◦ Inhaled Steroids • Inhalation of Possible Chemical or biological agents • Exposure to chemical or biological agents	• Shortness of Breath • Pursed-Lip breathing • Accessory muscle use, retractions, nasal flaring, fatigue • Inability to speak in sentences • Audible Wheezing or rhonchi • Fever, Cough • Cyanosis • Lung Sounds: ◦ Wet? Diminished? ◦ Bilaterally equal? ◦ Expiratory Wheezing?	• Asthma • CHF, Pulmonary Edema • Anaphylaxis • Pneumonia • Pulmonary Embolism • Cardiac • Hyperventilation • Inhaled Toxin • DKA • Pneumothorax

Universal Patient Care Protocol

Oxygen Administration Protocol

General Airway Management Protocol

Apply waveform Capnography

12 Lead ECG

Consider CPAP / Bi-PAP if available

Bronchodilators

Steroids

If ventilation efforts are not sufficient to deliver bronchodilators Effectively, consider a brief period of placing the bronchodilator in line with positive pressure, BVM, CPAP/BIPAP.
Careful attention to the patient's level of consciousness and ventilation must be maintained.

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

COPD

Consider Advanced Airway

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• COPD patients in severe respiratory distress should have oxygen delivery titrated to a Pulse OX greater than or equal to 92%. • If Asthma/COPD is in severe distress, treatment may occur simultaneously with IV, EKG, and 12-lead. • Consideration of Mag Sulfate in the updraft or IV, contraindications are ○ Heart block or myocardial damage ○ Hypermagnesemia • Remember: almost all cardiac problems produce some degree of respiratory distress. • A silent chest in severe respiratory distress is a pre-respiratory arrest sign. • Respiratory distress can be the result of metabolic acidosis from overdose and/or DKA, head injury, or trauma. • Preferred: Pulse Oximetry & Capnography should be monitored continuously for all patients with respiratory distress and/or respiratory failure. • NIVP- noninvasive Ventilation pressures – accomplished without intubation, preferably with BIPAP	

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Pulmonary Edema

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• Congestive heart failure • Past medical history • Medications ◦ Digoxin ◦ Lanoxin ◦ Lasix ◦ Viagra (or equivalent) • Cardiac history • Prior MI	• Respiratory distress • Bilateral Rales with decreased lung sounds • Jugular venous distension • Dependent edema (pedal, ascites) • Pink frothy sputum • Apprehension, orthopnea • Hepato-Jugular reflux • 3rd & 4th heart sound murmurs	• MI • Asthma • Pulmonary Hypertension • COPD, Cor Pulmonale • Anaphylaxis • Pleural Effusion, Pneumonia • Pulmonary Embolus • Drug Overdose, Toxic Exposure • Cardiac Tamponade

Universal Patient Care Protocol

Oxygen Administration Protocol

General Airway Management Protocol

Apply waveform Capnography

12 Lead ECG

Consider positive pressure ventilation or CPAP / Bi-PAP if available

- Consider sedation due to anxiety

Nitroglycerine

- Continue use as long as blood pressure allows

Hypotension

- Consider Cardiogenic Shock
 - Vasopressures

If Wheezing is present consider Bronchodilator

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Pulmonary Edema

If ventilation efforts are not sufficient to deliver bronchodilators effectively, consider a brief period of placing the bronchodilator in line with positive pressure, BVM, CPAP/BIPAP.

Careful attention to the patient's level of consciousness and ventilation must be maintained.

Consider Advanced Airway or PAI If the patient becomes exhausted with increased work of breathing secondary to hypoxia

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• If CPAP treatment has begun, notify ER so they can obtain a CPAP/BiPAP to continue the CPAP/BiPAP treatment upon arrival. • Caution with NTG if systolic BP < 120. • Consider tachycardia as the cause of pulmonary edema (especially V-tach.) Treat tachycardia. • Use Nitroglycerin with caution if acute inferior myocardial infarction is in progress. • Consider Nitro Drip if available • Furosemide may be considered for use in Pulmonary Edema. Follow AHA guidelines regarding the use of Furosemide.	

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Stridor (Upper Airway Obstruction)

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• Recent Intubation • Respiratory syncytial virus (RSV) • Respiratory Failure • Inhaled Toxins • Bronchiolitis • Medications: ◦ Inhaled Steroids • Inhalation of: ◦ Possible Chemical or biological agents • Exposure to chemical or biological agents	• Shortness of breath • Tripod positioning • Neck extended • Drooling • Barking cough • Pursed-lip breathing • Accessory muscle use, retractions, • nasal flaring, fatigue • Inability to speak in sentences • Audible stridor • Cyanosis	• Epiglottitis, Croup • CHF, Pulmonary Edema • Anaphylaxis • Pneumonia • Pulmonary embolus • Cardiac • Hyperventilation • DKA • Pneumothorax • Asthma, COPD

Universal Patient Care Protocol

Oxygen Administration Protocol

General Airway Management Protocol

Apply waveform Capnography

12 Lead ECG

Consider / Identify Etiology

- If stridor is secondary to trauma or burn injuries Proceed to Advanced Airway Management Protocol / PAI Protocol

Administer Epinephrine

- Nebulized Racemic Epinephrine
- Nebulized Epinephrine
 - 1:1k epi 0.3-0.5mg in 3cc NS

Steroid

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Stridor (Upper Airway Obstruction)

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• If heart rate increases greater than 20 beats per minute (bpm) while administering nebulized Racemic Epinephrine, or EPI 1:1000, then further dilute the treatment or stop administration. • Use Humidified Oxygen • Refrain from oropharynx manipulation	

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

ACUTE CORONARY SYNDROMES

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• Onset and Location of Pain • Provocation • Quality of Pain on 1-10 Scale • Radiation • Severity • Relationship to Exertion and Breathing • Previous Cardiac History • Risk Factors	• Radiating Pain (Jaw Pain) • Diaphoresis • Dyspnea • Palpitations • Weakness • Nausea or Vomiting • Feeling of Impending Doom	• Pulmonary Problems • Ulcers/GI Disorders • Medications • Anaphylaxis • Recent Trauma • Hyperventilation/Anxiety • Dissecting Thoracic Aneurysm

Universal Patient Care Protocol

General Airway Management Protocol

12 Lead ECG

- Obtain 12 Lead ECG within 10 minutes of patient contact
- Monitor a 12 Lead while proceeding with other treatment
- Repeat 12 Lead q 5 – 10 minutes or at rhythm change(s)
- Transmit or fax 12 Lead ECG to receiving hospital or notify receiving facility of possible STEMI or AMI within 5 minutes from ECG time
- Expedited transport to most appropriate facility ASAP

Consider / Identify Etiology

- Rate problem
- Inferior AMI suspected? (II, III, aVF)
 - Perform R-sided ECG V4-R
 - Titrate fluids to B/P & Lung Sounds
- If AMI is suspected in absence of ST elevation – consider 12 Lead with V7, V8, and V9 Posterior

Aspirin

Nitro

- May repeat Q 3-5 minutes until the pain is controlled or B/P drops below 90mmHg Systolic
 - And/Or initiate IV Nitro Drip
 - Ensure IV dose would equal or exceed SL dose, the goal is to alleviate pain, titrate dose to blood pressure.

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

ACUTE CORONARY SYNDROMES

Pain Management

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• Consider a Nitroglycerin Drip if chest pain does not resolve with oral administration of NTG. • Use caution when administering NTG if BP < 120 systolic, without venous access. • Chest pain that has moved or migrated should raise suspicion of an aortic dissection. • Use of a Benzodiazepine may be indicated for cocaine and or meth overdose with Chest pain.	

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

AMERICAN HEART ASSOCIATION GUIDELINES

Utilization of current:
American Heart Association guidelines
when performing
BLS, ACLS and PALS

It is recommended to utilize current AHA guidelines and standards at all times. If they differ from the NW Regional Protocols, the AHA guidelines should supersede these protocols.

These change frequently and are not listed specifically here.

AMERICAN HEART ASSOCIATION GUIDELINES

DEATH/WITHHOLDING RESUSCITATION

Purpose:

- The purpose of this protocol is to honor those who have obviously expired prior to EMS arrival and to honor the advanced directives of the patient as required by law.

Procedure:

- CPR and ALS treatment are to be withheld only if the patient is obviously dead or a valid written "Do Not Resuscitate" order is present. If a patient is in complete cardiopulmonary arrest (clinically dead) and meets one or more of the criteria below, CPR and ALS therapy need not be initiated:
 - Body decomposition
 - Injuries incompatible with life or entrapment with prolonged extrication time, decapitation, burning beyond recognition, massive open or penetrating trauma to the head or chest with obvious organ destruction
 - *Rigor mortis
 - *Dependent lividity
 - *Extended downtime with asystole on the EKG
- * **= must perform 4 lead ECG to verify asystole on the monitor looking in three leads minimum**
- If a bystander or EMR has initiated CPR or automatic defibrillation prior to paramedic arrival and any of the above criteria (signs of obvious death) are present, CPR and ALS therapy may be discontinued by the Paramedic. Once resuscitation is initiated, continue resuscitation efforts until either:
 - Resuscitation efforts meet the criteria for implementing the Termination of Resuscitation Protocol.
 - Patient care responsibilities are transferred to another appropriate caregiver.
- If doubt exists, or there is any question about the **validity** of a DNR order start resuscitation immediately. If there is a misunderstanding with family members or others present at the scene or if there are other concerns about following the DNR orders, contact the attending physician or medical control for guidance. When a DNR order is present unless otherwise **specifically** restricted, care shall be administered to provide comfort or alleviate pain except in those practices described as cardiopulmonary resuscitation. Depending on the needs of the patient this may include:
 - Basic airway management (BLS) including suctioning
 - Oxygen administration (including CPAP)
 - Pain Management
 - Trauma care
 - Transport
 - Family support

DEATH/WITHHOLDING RESUSCITATION

Do Not Resuscitate Form

- A DNR Order executed properly requires EMS personnel to withhold or withdraw cardiopulmonary resuscitation including intubation and advanced airway management, artificial ventilation, defibrillation, administration of cardiac resuscitation medications, and related procedures, from the patient in the event of a cardiac or respiratory arrest. The DNR Order form may be any document that includes the words "DNR", "No Code" or similar language, the physician's signature, and the date. Copies of the original are acceptable. The form may be found (but is not limited to) at the back door of the patient's bedroom, the nightstand by the patient's bed, the door of the refrigerator, or the patient's wallet. The patient, attending physician, or healthcare proxy may revoke the EMS/DNR order at any time. Document the presence of the DNR Order on the Patient Care Report. Include a copy of the DNR order with the PCR unless impracticable (ie single copy left with the coroner)

POLST (Physician Order for Life-Sustaining Treatment) Portable Medical Treatment Order

- **POLST is a process and a form**
- **POLST has [different names in different states](#).**
 - At the national level, it is simply called **POLST: Portable Medical Orders**, or POLST for short. Portable means that the order is valid outside the clinic or doctor's office, similar to a drug prescription.
- **POLST is many things, including:**
 - A process. Part of advanced care planning, which helps you live the best life possible.
 - Conversation. A good talk with your provider about your medical condition, treatment options, and what you want.
 - A medical order form that travels with you (called a POLST form).
- **POLST communicates your wishes as medical orders**
 - A POLST form tells all healthcare providers during a medical emergency what you want:
 - "Take me to the hospital" or "I want to stay here"
 - "Yes, attempt CPR" or "No, don't attempt CPR"
 - "These are the medical treatments I want"
 - "This is the care plan I want to be followed"
- **POLST is for the seriously ill or frail**
 - POLST gives seriously ill or frail people more specific direction over their health care treatments compared to advance directives and more options than Do Not Resuscitate (DNR) orders.
 - Who Needs a POLST, a Portable Medical Order?
 - Why is POLST not for healthy people?

Use the link to go to the Arkansas Department of health for more information and to access resources, obtain forms, etc...[POLST Resources](#)

DO NOT RESUSCITATE

INDICATION

An EMS/DNR Order form approved by the Department of Health executed properly requires EMS personnel to withhold or withdraw cardiopulmonary resuscitation to include intubation and advanced airway management, artificial ventilation, defibrillation, administration of cardiac resuscitation medications, and related procedures, from the patient in the event of a cardiac or respiratory arrest

PROCEDURE

The EMS/DNR Order form must be a document as approved by the Arkansas Board of Health, **or** one created or used by a physician that include the words “DNR”, “No Code” or similar language, the physician’s signature and the date. Copies of the original are acceptable.

The form may be found (but is not limited to):

- The back of the door of the patient’s bedroom.
- The nightstand by the patient’s bed.
- The door of the refrigerator
- The patient’s wallet

Care shall be administered to provide comfort or alleviate pain except those practices described above as cardiopulmonary resuscitation. Depending on the needs of the patient this may include:

- Basic airway management (BLS) including suctioning
- Oxygen administration
- Pain Management
- Trauma care
- Transport
- Family support

If there is a misunderstanding with family members or others present at the scene or if there are other concerns about following the EMS/DNR orders, attempt to contact the attending physician or medical control for guidance. If there is any question about the validity of an EMS/DNR order, resuscitate while contacting medical control.

The patient, attending physician, or healthcare proxy may revoke the EMS/DNR order at any time.

Document the presence of the EMS/DNR Order on the Patient Care Report. Include a copy of the EMS/DNR order with the PCR unless impracticable (ie single copy left with coroner)

DO NOT RESUSCITATE

TERMINATION OF RESUSCITATION PROTOCOL

INDICATIONS

- Unsuccessful Cardiopulmonary resuscitation (CPR) and other advanced life support (ALS) interventions may be discontinued prior to transport or arrival at the hospital when this procedure is followed.
- The purpose of this protocol is to allow for the discontinuation of prehospital resuscitation after delivery of adequate and appropriate ALS therapy.
- $ET\ CO_2 < 10$ for 10 minutes of high quality CPR


PROCEDURE

1. The following criteria must be met before consulting Medical Control for discontinuation of prehospital resuscitation attempts:
 - Patient must be 18 years of age or older
 - Adequate CPR has been administered
 - Endotracheal intubation or other advanced airway has been successfully accomplished with adequate ventilation
 - Vascular access has been achieved
 - No evidence or suspicion of any of the following:
 - Drug/toxin overdose
 - Active internal bleeding
 - Hypothermia
 - Rhythm-appropriate medications and if indicated defibrillation have been administered according to protocol for a total of 3 cycles of drug therapy (epinephrine) without return of spontaneous circulation (palpable pulse)
 - All Paramedic personnel involved in patient care agree that discontinuation of the resuscitation is appropriate
2. If all of the above criteria are met and discontinuation of prehospital resuscitation is desired, **Consult Medical Control**

TERMINATION OF RESUSCITATION

ASYSTOLE

History • Past medical history • Medications • Events leading to arrest • End stage renal disease • Suspected hypothermia • Suspected drug overdose • DNR or Living Will • Estimated down time	Signs and Symptoms • Pulseless • Apneic • No electrical activity on ECG	Differential • Medical or Trauma • Hypoxia • Potassium (Hypo / Hyper) • Drug Overdose • Acidosis • Hypothermia • Device (ECG Lead) error • Death
--	--	--




NOTES:

- Exam: ABCs, Vital Signs, Mental Status.
- Always confirm Asystole in 3 or more leads.
- Efforts should be toward determining the cause of the arrest.
- Higher dose epinephrine may be indicated in presence of Beta Blocker or Calcium Channel Blocker Overdose or anaphylactic shock with cardiac arrest

ASYTOLE

BRADYCARDIA

History - Past History of Bradycardia - Medications: Beta Blockers, Clonidine, Calcium Channel Blockers, Digitalis - Pacemaker - Nausea and Vomiting	Signs and Symptoms - HR < 60/min - Chest Pain - Respiratory Distress - Acute Coronary Syndrome - Hypotension - Decreased LOC - Weakness	Differential - AMI - Hypoxia - Hypothermia - Stroke - Head Injury - Vasovagal - Athletes
--	---	---




NOTES:

- Atropine is often ineffective for third-degree heart block or Mobitz type II second-degree heart block.
- Consider and treat causes of Bradycardia
- Atropine should be used with caution in patients with suspected AMI.
- Attempting to increase the rate of an asymptomatic patient is contraindicated.
- Right ventricular infarction may present with bradycardia, consider fluid challenge in the absence of pulmonary edema.
- PVCs may occur if the rate falls below 30-40 beats/min. Do not treat PVCs in bradycardic arrhythmias.
- Versed 2.5 mg slow IV, max of 5 mg, may use as a sedative agent in conjunction with pain management for Pacing
- Do not delay pacing in high degree AV block or critical patients presenting in bradycardia

BRADYCARDIA

CARDIAC ARREST

History • Events leading to cardiac arrest • Estimated downtime • Past medical history • Medications • Existence of terminal illness • Signs of lividity, or rigor mortis • State DNR or Living Will	Signs and Symptoms • Unresponsive • Apneic, agonal • Pulseless	Differential • Medical vs. Trauma • Ventricular Fibrillation, Pulseless Ventricular Tachycardia • Asystole • PEA
---	---	--




NOTES:

- CPR should not be interrupted, except under certain circumstances—endotracheal intubation, moving patient up or down stairs.
- If prolonged BLS prior to arrival consider NG/OG tube placement.
- Cardiopulmonary resuscitation may be discontinued: See Termination of Resuscitation Protocol.
- Be aware of any appropriate DNR, call medical control if you are uncertain on how to proceed.
- Use of Mechanical CPR device is appropriate and should be used if available if patient meets criteria for it's use.
- CPR is defined as continuous compressions with ventilation administered without interruptions in compressions regardless if patient is intubated or not.

CARDIAC ARREST

NARROW COMPLEX TACHYCARDIA

History • Medications (Aminophylline, decongestants, thyroid supplements, diet pills, Digoxin) • Diet • Illicit drugs (methamphetamine, cocaine, stimulants) • Past Medical History • History of Palpitations/heart racing • Syncope • Near Drowning	Signs and Symptoms • HR >150/min • QRS <0.12 sec • Dizziness, chest pain, shortness of breath • Potential presenting rhythm • Sinus Tachycardia • Atrial Fibrillation/ Flutter • Multifocal Atrial Tachycardia	Differential • Heart Disease (WPW, Valvular) • Sick Sinus Syndrome • Electrolyte Imbalance • Exertion, Pain, Emotional Stress, Fever • Drug Effect/ Overdose • Hyperthyroidism • Sinus Tach
---	---	--




NOTES:

- Establish rapid heart rate as cause of signs and symptoms.
- If Wolfe Parkinsons White (WPW) is suspected – do not administer Adenosine or Calcium Channel Blockers
- Adenosine 3 mg if patient is taking dipyridamole or Cobalazine

NARROW COMPLEX TACHYCARDIA

POST RESUSCITATION

History • Cardiac Arrest • Respiratory Arrest	Signs and Symptoms • Return of Pulse	Differential • Continue to address specific differentials associated with original dysrhythmia
---	---	---


**NOTES:**

- A 12 lead EKG should be obtained as soon as possible to determine the presence of an acute coronary syndrome.
- ETT and/or alternate airway device should not be removed unless Medical Control is contacted.
- Versed and/or Ketamine may be used for sedation in order to maintain a controlled airway
- Narrow Complex Tachycardia in the post resuscitation phase may be due to Epinephrine and/or Atropine therapy and usually does not require treatment-monitor BP.
- Consider NG/OG tube placement for gastric decompression.
- Place second IV if possible.
- Consider temperature regulation; allow mild hypothermia and treat hyperthermia

POST RESUSCITATION

PULSELESS ELECTRICAL ACTIVITY

History • Events leading up to arrest • Estimated down time • Past medical history/ medications • Renal failure/dialysis • DNR • Hypothermia • Suspected Overdose (digitalis, tricyclics, Beta-blockers, Calcium channel blockers)	Signs and Symptoms • Unresponsive, Apneic, pulseless with organized electrical activity	Differential • Medical vs. Trauma etiology • Massive Myocardial Infarction • Pulmonary Embolism • See Frequent Causes
---	--	---




NOTES:

- Hypoxia is the most common cause of reversible PEA
- Use Atropine for Vagal stimulation in Cardiac Arrest to increase heart rate.

PULSELESS ELECTRICAL ACTIVITY

VENTRICULAR FIBRILLATION/PULSELESS V-TACH

History - Estimated down time - Past medical history/ medications - Events leading to arrest - Renal failure/dialysis - DNR - Hypothermia - Electrocution	Signs and Symptoms Unresponsive, Apneic, pulseless	Differential - Medical vs. Trauma etiology - Artifact - Asystole - Device failure
--	---	---


**NOTES:**

- Left Ventricular Assist Device (LVAD) patients may not have a palpable pulse – this does not indicate that perfusion is compromised- check color and LOC. See guideline section for further details on LVAD
- Defibrillation takes precedence over all treatment once the defibrillator is available.
- For hypothermic patients defibrillation may not be effective, see Hypothermia Protocol.
- If patient successfully converted with Automatic Implantable Cardiac Defibrillator (AICD), consider antiarrhythmic therapy, contact Medical Control.
- Magnesium Sulfate for V-fib refractory to above treatments, for digitalis toxicity, and for Torsades.
- If Amiodarone is contraindicated or unavailable– Lidocaine may be used.
- If V Fib is not responding to Defibrillations, consider changing pad placement and continue shocking

VENTRICULAR FIBRILLATION/ PULSELESS V-TACH

WIDE COMPLEX TACHYCARDIA

History	Signs and Symptoms	Differential
Prior cardiac history	- Heart rate typically > 150/min - Wide Complex	- Aberrantly conducted SVT - Aberrantly conducted A Fib




NOTES:

- You may consider precordial thump or cough if witnessed V Tach occurs
- For unstable V Tach with regular rhythm – you may consider use of antiarrhythmic – consider if vitals are stable.
- Polymorphic: more than one origin (shape); Monomorphic: one origin (shape.)
- 90% of wide complex tachycardias are V-Tach.
- Irregular wide complex tachycardia may be Atrial Fibrillation with WPW or aberrant conduction – administer Amiodarone – do not cardiovert.
- Look for dissociated P waves on EKG.
- Medications may be given simultaneously with cardioversion. **Promptly cardiovert for hemodynamic instability!**
- Measure baseline QT intervals on all patients: Meds that prolong QT—Procainamide, Amiodarone, Quinidine.
- Check medications already on board—do not mix medications that prolong the QT interval.
- Do not mix the use of antiarrhythmic medications.
- May go directly to synchronized cardioversion as in unstable. Check for V-Tach clues and/or 12 lead
- Do not** administer Calcium Channel Blockers for V Tach or suspected V Tach or unknown wide complex tachycardia
- If Amiodarone is contraindicated or unavailable – Lidocaine may be used**

WIDE COMPLEX TACHYCARDIA

ABDOMINAL PAIN

History • Past medical, surgical history • Medications • Sepsis • Fever • Time of last meal • Last bowel movement/emesis • Menstrual history (pregnancy)	Signs and Symptoms • Pain (location/migration) • Tenderness (palpation) • Nausea/Vomiting • Dysuria • Constipation • Vaginal bleeding/discharge • Pregnancy • Associated symptoms: Fever, headache, weakness, malaise, myalgias, cough, mental status changes, rash • Syncope	Differential • Pneumonia • Liver (hepatitis, CHF) • Peptic Ulcer Disease/Gastritis • Gallbladder • Myocardial Infarction • Pancreatitis • Kidney Stone • Abdominal Aneurysm • Appendicitis • Bladder/Prostate Disorder • Pelvic (PID, Ectopic Pregnancy, Ovarian Cyst) • Spleen Enlargement • Diverticulitis • Bowel Obstruction • Gastroenteritis (infection) • Dehydration
---	--	--




NOTES:

- Abdominal pain in women of child-bearing age should be treated as ectopic pregnancy until proven otherwise.

ABDOMINAL PAIN

ACUTE STROKE—CVA

History • Previous CVA, TIAs • Previous cardiac, vascular surgery • Associated disease: diabetes, hypertension, arteriosclerosis, Atherosclerotic Coronary Artery Disease • Atrial fibrillation • Medications, blood thinners • History of trauma	Signs and Symptoms • Altered mental status • Weakness/paralysis • Blindness or other sensory loss • Aphasia/dysarthria • Syncope • Vertigo/dizziness • Vomiting • Headache • Seizures • Respiratory pattern change • Hypertension/hypotension	Differential • Diabetic Emergency • Stroke: Thrombotic/Embolic/Hemorrhagic • Tumor • Head trauma • Central nervous system injury • Seizure, Sepsis • Toxic ingestion/Overdose • Alcohol intoxication • Environmental exposure • Psychiatric disorder
---	--	--




NOTES:

- Thrombolytic therapy may be possible with any acute stroke defined by duration of symptoms of less than 3- 4 1/2 hours. Scene times and transport times should be minimized in this setting.
- Onset of symptoms is defined as the last witnessed time the patient was symptom free (i.e. awakening with stroke symptoms would be defined as an onset time of the previous night when the patient was symptom free).
- Elevated blood pressure is commonly present with CVA. Contact Medical Control and consider treatment if diastolic is > 120 mmHg or systolic > 230 mmHg. Be alert for airway problems (swallowing difficulty, vomiting).

ACUTE STROKE—CVA

ALTERED LEVEL OF CONSCIOUSNESS/MENTATION

History	Signs and Symptoms	Differential
- Known diabetic, medic alert tag - Seizures - Drugs, drug paraphernalia - Report of illicit drug use or possible overdose/ingestion - Fever/febrile illness	- Weakness/paralysis - Changes in baseline mental status - Bizarre behavior - Hypoglycemia/hyperglycemia - Syncopal - Vertigo/dizziness - Headache - Seizures - Respiratory pattern change - Hypertension/hypotension	- Hypoxia - Cardiac Dysrhythmias - Diabetic Emergency - Central nervous system Disorder/Injury/Disease - Toxic ingestion/Overdose - Drug/Alcohol intoxication - Environmental exposure - Psychiatric disorder


**NOTES:**

- All possible causes of altered mental status should be considered. (AEIOUTIPS)
- Observe and assess environment to gather information for medical control.
- Proceed to more specific protocol when cause of Altered LOC determined.
- Be alert for airway problems (swallowing difficulty, vomiting) AND MANAGE AGGRESSIVELY.
- Hypoglycemia can present as a localized neurological deficit, especially in the elderly.
- Consider restraints if necessary for patient's and/or personnel safety.
- Consider stimuli to wake patients PRN. If patient does respond to stimuli, there may still be an underlying medical condition that requires attention.
- Consider antipsychotic or sedative for acute psychosis or severe agitation.

ALTERED LEVEL OF CONSCIOUSNESS/MENTATION

BITES, STINGS, and ENVENOMATIONS

History - Type and time of bite - Location of bite - Prior first aid given - Past medical history - Progression of signs and symptoms	Signs and Symptoms - Parasthesias, pain - Chills/weakness - Nausea/vomiting - Headache - Hypotension - Discoloration, edema - Difficulty breathing or swallowing - Cardio – Respiratory Arrest - Dysrhythmias	Differential - AMI - Medical illness - Other toxic exposure - Anaphylaxis
--	--	---


**NOTES:**

- Do not apply ice or constrictive bandage (tourniquet) to site.
- All dog bites/attacks **must** be reported to law enforcement.
- Progression of swelling should be marked every five minutes in order to monitor.
- Identification of the animal/substance should be made only if rescuer safety is not compromised.
- Patients who appear asymptomatic should be transported for observation. Some signs and symptoms may take up to 24 hours to appear.
- Human bites should always be transported to ED due to high possibility of infection.
- Maintain affected extremity below level of heart – dependent.

BITES, STINGS, AND ENVENOMATIONS

DIABETIC EMERGENCIES

History - Known diabetic, medic alert tag - Possible illicit drug use - Chronic alcohol abuse - Medications - Excessive thirst, hunger or urination	Signs and Symptoms - Decreased mental status - Bizarre behavior - Cool diaphoretic skin - Fruity, ketotic breath - Kussmaul respirations - Signs of dehydration	Differential - Trauma - CVA, seizure, sepsis - Cardiac - Shock - Toxic ingestion/ alcohol intoxication - Environmental exposure - Psychiatric disorder
--	---	---



NOTES:

- Underlying coronary disease including AMI or CVA should also be considered with middle aged – elderly patients presenting as diabetic emergencies.
- Ascertain the patient's insulin regimen (dosage) for ED reference.

DIABETIC EMERGENCIES

Fever

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• Age • Report of fever • Severity • Past medical history • Medications • Previous doses of Ibuprofen or Tylenol	• Warm or hot skin • Flushed • Sweaty • Chills • Headache • Mental status changes	• Sepsis • Drug reaction • Overdose • Heatstroke • Metabolic Abnormalities

Universal Patient Care Protocol

Temperature greater than 100.4 F (38 C)

Consider antipyretic


- Has the patient received antipyretic medication prior to EMS contact?
 - Identify time and name of the medication
 - Has it been greater than 4-6 hours
 - Readminister antipyretic as needed
 - Antipyretics can be alternated by brand/trade name at the 2 or 3-hour mark to maintain a 4-6 hour recommendation between dose and still not exceed the maximum daily doses per the manufacturer
- Monitor and reassess as needed

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• Febrile seizures are more likely in children with a history of febrile seizures and/or with a rapid change in temperature • Patients with a history of liver failure should not receive acetaminophen • Rehydration with fluids increases the patient's ability to sweat and improves heat loss • NSAIDS should not be used in the setting of environmental heat emergencies	

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

HEAT EMERGENCIES

History • Age • Exposure to increased temperatures and/or humidity • Past medical history/medications • Extreme exertion • Time and length of exposure • Poor PO intake • Fatigue and/or muscle cramping • Not conditioned for hot/humid environment	Signs and Symptoms • Altered mental status or unconsciousness • Hot, dry or sweaty skin • Hypotension or shock • Seizures • Nausea • Heat Cramps- benign muscle cramping secondary to dehydration and not associated with an elevated temperature. • Heat Exhaustion -: dehydration, salt depletion, dizziness, fever, mental status changes, headache, cramping, nausea and vomiting. Vital signs: tachycardia, hypotension and elevated temperature • Heat Stroke- Heat Stroke: dry skin, dehydration, tachycardia, hypotension, temperature > 104°F (40°C) and an altered mental status. True emergency, must be RAPIDLY cooled.	Differential • Fever • Dehydration • Medications • Hyperthyroidism • Delirium tremens • CNS lesions or tumors
--	--	---




NOTES:

- Extremes of age are more prone to heat emergencies (i.e. young, old).
- Predisposed by use of: tricyclic antidepressants, phenothiazines, anticholinergic medications, and alcohol.
- Cocaine, amphetamines, and salicylates may elevate body temperature.
- Sweating generally disappears as body temperature rises above 104°F (40° C).
- Intense shivering may occur as patient is cooled.

HEAT EMERGENCIES

HYPERTENSIVE CRISIS

History • Documented Hypertension • Medications • Pregnancy • Viagra • Diabetic/Renal impairment • Recent trauma	Signs and Symptoms • Headache • Chest pain • Dyspnea • Blurred vision • Signs & symptoms Acute Ischemic Attack/CVA • Weakness • Vertigo • Epistaxis	Differential • Central nervous system injury • AMI • Aneurysm • Preeclampsia • Hypertensive Encephalopathy • Emotional Crisis
--	---	---




NOTES:

- See Preeclampsia/Eclampsia Protocol if patient is pregnant and has no previous history of hypertension.
- Do not attempt to rapidly decrease the BP if the patient is exhibiting signs and symptoms of Acute Ischemic Attack/CVA.
- Patient should be transported with the head elevated if possible.
- Nitroglycerine may be indicated if Labetalol is not available.
- Avoid Nitroglycerine if the patient has taken Erectile Dysfunction Medication within the last 24 hours.
- Reassess BP after each medication administration.

HYPERTENSIVE CRISES

HYPOTENSION—SYMPTOMATIC

History - Blood loss: GI, vaginal, acute abdominal aneurysm - Fluid loss: vomiting, diarrhea, fever, infection - Cardiac: AMI, CHF - Medications: narcotics, antihypertensives, anticoagulants - Allergic reaction - Pregnancy - Recent surgery or long bone fracture	Signs and Symptoms - Restlessness, thirst - Confusion, change in level of consciousness/mentation - Weak/rapid pulse - Pale, cool, diaphoretic, clammy skin - Hemodynamic instability - Delayed capillary refill - Hemorrhage - Tarry Stool - Chocolate colored emesis	Differential - Shock: hypovolemic, cardiogenic, septic, neurogenic, anaphylactic - Ectopic pregnancy - Dysrhythmias - Pulmonary embolus - Tension pneumothorax - Medication effect/Overdose - Vasovagal/Syncope
--	---	--




NOTES:

- Maintain a Mean Arterial Pressure (MAP) above 60.
- Oxygen is still the most important drug to administer to patients in shock.
- It is always a good idea to ask patients what their normal BP is, if known.
- Consider all possible causes of shock and treat per appropriate protocol.
- A systolic BP between 90-100 mm Hg may be normal for a healthy, physically fit individual.
- 3rd trimester pregnant patients will become hypotensive when placed supine—be sure to place them left-laterally recumbent or elevate right side.
- Pregnant patients will shunt blood away from the fetus. Aggressive fluid resuscitation may be necessary. When in doubt, contact medical control.
- A Dopamine infusion should not be abruptly stopped, but should be titrated.
- In children -Maintain perfusion with fluid resuscitation, systolic BP of $70 + 2 \times \text{age}$ if over 1 year old.
- Dialysis patient – use of excessive fluid in dialysis patients maybe harmful.

HYPOTENSION—SYMPTOMATIC

HYPOTHERMIA

History - Extremes of age - Past medical history/medications - Exposure to environment even in normal temperatures - Drug use, alcohol, barbiturates - Wet - Infection, Sepsis - Length of exposure	Signs and Symptoms - Cold, clammy skin - Shivering - Altered mental status or unconsciousness - Extremity pain or sensory abnormality - Bradycardia - Hypotension or shock	Differential - Sepsis - Hypoglycemia - CNS Dysfunction: stroke, head injury, spinal cord injury
--	--	--


**NOTES:**

- NO PATIENT IS PRONOUNCED DEAD UNTIL WARM AND DEAD.
- Mild Hypothermia – normal body temp of 95° – 89.8° F
- Moderate Hypothermia-89.7° – 82.5° F
- Severe Hypothermia 82.4° – 75.2° F
- Profound Less than 75.2°
- Hypothermia may cause severe bradycardia.
- The patient must be rewarmed before treatments will be effective. In cardiac arrests, provide first round defibrillations and vasopressors as rewarming occurs while withholding antiarrhythmic's until patient is rewarmed to 86F or 30C, (Withhold repeat efforts until rewarmed) then antiarrhythmic's may be given if indicated at prolonged intervals until the patients temp is 93F or 34C at which time refer to appropriate ACLS protocol.
- Shivering stops below 90°F (32°C).
- Hot packs should be placed in the armpits and groin. Care should be taken not to place the packs in direct contact with the skin. Use a towel or 4X4 as a barrier.
- Hypothermia may be predisposed by use of: tricyclic antidepressants, phenothiazines, anticholinergic medications, and alcohol.

HYPOTHERMIA

NAUSEA—VOMITING—DIARRHEA

History • Time of last meal • Onset/duration of complaint • Other sick contacts • Coffee ground emesis, dark tarry stools • Past medical history • Recent surgery or trauma • Medications: new meds • Menstrual history/pregnancy • Radiation/chemotherapy • Toxic exposure • Motion sickness	Signs and Symptoms • Pain – OPQRST • Abdominal Distension • Fever • Headache • Blurred vision • Nausea increases with movement • Weakness • Diaphoresis • Dysuria • Bradycardia/tachycardia	Differential • Myocardial Infarction • CNS (Headache, stroke, recent head trauma) • GI bleed • Drugs (antibiotics, narcotics, chemotherapy) • Diabetic Ketoacidosis • Pneumonia • Influenza • Food or Toxin induced • Pregnancy • Vertigo • Vagal Response
--	---	---




NOTES:

- Silent AMI may present with Nausea/Vomiting

NAUSEA—VOMITING—DIARRHEA

NEAR DROWNING/DROWNING

History • Diving/SCUBA Diving • Events leading to submersion • Drug use, alcohol, barbiturates • Length of time submerged • Type and temperature of water • Extremes of age • Past medical history/medications	Signs and Symptoms • Apnea • Hypothermia • Paralysis • Shortness of Breath • Arrhythmias	Differential • Trauma • Pre-existing medical condition • Drug/alcohol ingestion
---	---	--


**NOTES:**

- Near drowning patients who have any resuscitation efforts should always be transported to the hospital due to secondary pulmonary edema.
- Asymptomatic patients should be transported for observation. Symptoms may be delayed for 24 hours.
- Blood glucose should be assessed in patients with extended submersions.
- If the safety of rescuers is not compromised, patients found in the water should have spinal restriction protocol before removal from water.
- Consider the use of PEEP.

DROWNING/NEAR DROWNING

OVERDOSE/TOXIC EXPOSURE

History	Signs and Symptoms	Differential
- Ingestion or suspected ingestion of a potentially toxic substance - Substance ingested, route, quantity, time - Reason (suicidal, accidental, criminal), prior history - Available medications in home - Past medical history, medications	- Mental status changes - Hypotension/Hypertension - Decreased respiratory rate - Tachycardia, dysrhythmias - Seizures - Pupils status - Signs of illicit drug use	- Reasons for Coma (AEIOUTIPS) - Tricyclic antidepressants - Acetaminophen (Tylenol) - Depressants - Stimulants - Anticholinergic - Cardiac medications - Solvents, Alcohols, Cleaning Agents,




NOTES:

- Do not rely on patient history of ingestion, especially in suicide attempts.
- Bring bottles, contents, emesis to ED.
- Consider polysubstance (multiple drugs).
- Cardiac Meds:** dysrhythmias and mental status changes
- Tricyclic Antidepressants:** 4 major areas of toxicity-seizures, dysrhythmias, hypotension, decreased mental status or coma; Rapid progression from alert mental status to death.
- Acetaminophen:** Initially normal or N/V. If not detected and treated, causes irreversible liver failure.
- Depressants:** ↓HR, ↓BP, ↓ respirations, ↓ temperature, nonspecific pupils.
- Stimulants:** ↑HR, ↑BP, ↑respirations, ↑ temperature, dilated pupils, seizure.
- For Salicylate overdose** – intubated patients should be ventilated at a faster rate – CO₂ at 35 is target reading.

OVERDOSE/TOXIC EXPOSURE

SEIZURE

History • Documented seizure disorder • Medications • Pregnancy • Trauma	Signs and Symptoms • Decreased mental status • Sleepiness • Incontinence • Observed seizure activity • Evidence of Trauma	Differential • CNS Injury of Hx of • Tumor • Hypoxia • Fever • Eclampsia • Renal failure • Drug use • Infection • Alcohol withdrawal • Metabolic disorder • Electrolyte imbalance • Pseudoseizures
--	--	--



NOTES:

- Anticonvulsants should only be used when patient exhibits **ACTIVE, CONTINUOUS** seizure.
- See Preeclampsia/Eclampsia protocol if patient is pregnant and has period of recovery or consciousness.
- Assess possibility of recent traumatic event and drug abuse (i.e. stimulants).
- Remember, febrile seizures in infants and children are relatively benign; most common cause of seizure in pediatric patient – should be transported to the ED for physician evaluation.
- If pseudoseizures are a consideration, consider stimulus.
- If PAI is used to control airway, note that seizure activity may still be present but not visible

SEIZURE

Sepsis

<u>History</u>	<u>Signs and Symptoms</u>	<u>Differential</u>
• Fever • Recent history of Diagnosed infection • Recent surgery • Compromised immune system • Age (more common in very young and very old) • Suspected Infection • Indwelling Catheter	• Altered level of consciousness • Heart rate > 90 beats per minute • Respiratory rate > 20 breaths per minute • Decreased urine output • Abdominal pain • Delayed capillary refill • Temperature > 101 F or < 96.8 F	• Overdose • Shock-Hypovolemic, • Cardiogenic, Anaphylactic • Dehydration • Drug Interaction

Universal Patient Care Protocol

End Tidal Capnography

<u>Diagnosed or suspected infection and any two of the following:</u> • Altered Mental Status • Heart Rate > 90 BPM • Respiratory Rate > 20 BPM • Systolic B/P < 90 mmHg systolic • Temperature > 101F or < 96.8F • ETCO2 < 26 mmHG without hyperventilation present	No	Proceed to appropriate protocol
	Yes	Transport to appropriate facility

Administer Crystalloid Fluids as needed

- Confirm and reassess lung sounds
- **Treat Hypotension to maintain a MAP of 60mmHg**
 - Do not exceed 30 ml/kg of fluid administration
 - Consider Vasopressors early

Considerations

- Continued assessment
- Consider secondary vascular access
- Contact Medical Control

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.


Sepsis

<u>Notes</u>	<u>Associated Protocols/Resources</u>
• Early detection of sepsis is critical to patient survival. Notify receiving facility if sepsis is suspected • A history of dialysis or CHF does not contraindicate aggressive fluid therapy. • Increased serum lactate is commonly used to diagnose sepsis. As lactate increases, ETCO₂ decreases, making capnography an excellent indicator of sepsis in the field.	Hypotension Epinephrine Levophed Push Dose Pressors

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

SYNCOPE

History • Cardiac, stroke, seizure • Occult blood loss, GI, Ectopic rupture • Last Menstrual Period, vaginal bleeding, pregnancy • Fluid loss: nausea, vomiting, diarrhea • Past medical history • Medications • Emotion, stressful event	Signs and Symptoms • Loss of Consciousness with recovery • Lightheadedness, dizziness • Palpitations, slow or rapid pulse • Pulse irregularity • Decreased blood pressure • Numbness and/or tingling in extremities • Carpal/pedal spasms	Differential • Vasovagal • Cardiac syncope • Urination/defecation • Psychiatric • Stroke • Shock • Hypoglycemia • Seizure • Toxicological (overdose/alcohol) • Medication • Hyperventilation
--	--	---



NOTES:

- Assess for signs and symptoms of trauma if associated or possible fall with syncope.
- Patients suffering syncopal episodes should be transported.
- Over 25% of geriatric syncope is dysrhythmia based.
- Be alert for airway problems (swallowing difficulty, vomiting) AND MANAGE AGGRESSIVELY.

SYNCOPE

CHEST TRAUMA

History • Time and Mechanism (blunt, penetrating, crushing, etc.) of injury • Damage to structure or vehicle • Location in structure or vehicle • Others injured or dead • Speed and details of MVC • Restraints/protective equipment • Past medical history/medications	Signs and Symptoms • Pain—specific or general • Cyanosis • Dyspnea/increased work of breathing • Anxiety • Tachycardia • Hypotension • Bruising/hematoma • Bleeding • Fractured ribs • Abnormal breath sounds • Open/sucking wounds • Crepitus, paradoxical movement • Decreased ventilatory compliance • Sub Cutaneous Emphysema	Differential • Trauma vs. Medical • Angina/AMI/CHF • Pericarditis • Pulmonary embolism • Asthma/COPD • Pneumothorax (tension, spontaneous) • Hemothorax • Aortic dissection or rupture • Blunt vs. penetrating
---	---	---



NOTES:

- Consider air transport when indicated.
- Significant chest injuries must be addressed – treated immediately when found – definitive care is surgery – rapid transport.
- For Large flail sections without pneumothorax – consider positive pressure ventilation

CHEST TRAUMA

DESTINATION PROTOCOL

(Always use the current Destination Protocol approved by NW TRAC)

Regional Destination Protocol for the Northwest Arkansas Trauma

Regional Advisory Council Area (NW TRAC) *Approved per NWA TRAC- 3/16/2012*

Trauma Transport Destination Guideline

Northwest Arkansas Regional Trauma Advisory Council

The following reflects the Pre-hospital Triage and Decision Scheme of the ADOH Rules and Regulations for Trauma Systems, and American College of Surgeons Field Triage Decision Scheme.

All trauma patients shall have a trauma band and be evaluated against the criteria to determine the need for rapid transport to the appropriate **level trauma center**.

If the trauma patient meets any one of the MAJOR or MODERATE criteria listed below consider the patient a **trauma alert** and notify dispatch as soon as possible.

On scene times for patients meeting the trauma alert criteria shall be 10 minutes or less.

Destination shall be determined by the severity of injury and the distance to the closest appropriate Trauma Center.

Transport of the **trauma alert** patient to the receiving facility shall be in the emergency mode, unless otherwise determined by Medical Control.

Always utilize the most current NW TRAC Destination Protocol, as of publication of this document, the following pages were currently being utilized.

Helicopter transport should be considered when time is critical and transport to a higher level appropriate Trauma Center is warranted from the scene. Early activation and concurrent dispatch of helicopter transport should be considered when the dispatcher identifies the potential for MAJOR or MODERATE trauma injuries.

Inability to establish or maintain an adequate airway or control excessive hemorrhage for trauma patients requires transport to the closest appropriate facility.


Multiple trauma patient situations may require interaction with Arkansas Trauma Communications Center (ATCC) or area trauma centers distribute trauma patients to avoid overtaxing the trauma centers.

ATCC should be notified for all Moderate and Major Trauma Injuries

(This Protocol is continued on next two pages)

DESTINATION PROTOCOL

Field Triage Decision Scheme: The Arkansas Trauma Triage Protocol



TS-Field Triage Decision Scheme rev. October 28, 2013

DESTINATION PROTOCOL

GLASGOW COMA SCORE

ADULT

MOTOR RESPONSE		EYE OPENING		VERBAL RESPONSE	
Obeys commands	6	Spontaneous	4	Oriented	5
Localizes	5	To Voice	3	Confused	4
Withdrawal	4	To Pain	2	Inappropriate	3
Flexion	3	None	1	Incomprehensible	2
Extension	2			None	1
None	1				

PEDIATRIC - recommended from 4years of age to adult

MOTOR RESPONSE		EYE OPENING		VERBAL RESPONSE	
Obeys commands	6	Spontaneous	4	Oriented & converses	5
Localizes	5	Verbal command	3	Disoriented & converses	4
Withdrawal	4	To pain	2	Inappropriate	3
Flexion-withdrawal	3	No response	1	Incomprehensible	2
Flexion-abnormal	2			None	1
None	1				


INFANT - recommended for birth to 4 years of age

MOTOR RESPONSE		EYE OPENING		VERBAL RESPONSE	
Spontaneous	6	Spontaneous	4	Smiles, oriented to sound, interacts appropriate	5
Localizes pain	5	Reacts to speech	3	Crying - consolable Interacts - inappropriate	4
Withdraws in response to pain	4	Reacts to pain	2	Crying - inconsistently consolable; interacts – restless	3
Abnormal flexion in response to pain	3	No response	1	Crying - inconsolable Interacts - restless	2
Abnormal extension in response to pain	2			No response	1
No response	1				

GLASCOW COMA SCORE

DIVING EMERGENCIES (SCUBA)

History • Diving/SCUBA Diving • Events leading to dive/ascent • Dive within 36 hours of event • Depth of dive • Length of dive • Description of ascent	Signs and Symptoms • Headache, disorientation, vertigo • Nausea, abdominal pain • Chest Pain, Dyspnea, visual disturbances • Joint pain, paralysis • Seizure, decreased LOC • Pulmonary Edema • Cardiac Arrest	Differential • AMI, CVA, Seizure • Diabetic condition • Trauma • Carbon monoxide/toxins
--	---	---


**NOTES:**

- Decompression symptoms usually manifest within 20 minutes of surfacing.
- PT's who receive oxygen may become symptom free but still require hyperbaric treatment
- Air embolism is the most serious complication of pulmonary barotraumas.
- If diver loses consciousness immediately after surfacing, an air embolism should be suspected.
- Evaluate patient for presence of pneumothorax.
- If possible bring the divers dive records (Dive Tables) to the hospital with the patient.
- DAN (Diver Alert Network): (919)-684-8111 or (919)-684-4326

DIVING EMERGENCIES (SCUBA)

EXTREMITY TRAUMA

History - Type of injury - Mechanism: crush, penetrating, amputation - Time of injury - Open vs. closed wound or fracture - Wound contamination - Age - Past medical history - Medications	Signs and Symptoms - Pain, swelling, - Deformity - Altered sensation or motor function - Diminished pulse or capillary refill - Decreased extremity temperature	Differential - Abrasion - Contusion - Laceration - Sprain/Strain - Dislocation - Fracture - Amputation
--	--	---



NOTES:

- If evidence of open fracture exists it is permissible to administer antibiotic (Cefazolin) to patient
- In amputations and pulseless extremities, time is critical. Transport and notify medical control immediately.
- Notify hospital of **Trauma Alert** as soon as practical.
- Document and mark distal pulses.
- Hip, knee, and elbow fractures and/or dislocations have a high instance of vascular compromise.
- Splint in position found unless: no distal pulse, unable to transport patient in position found, there is severe pain with angulation.
- Do not attempt to realign open fractures unless necessary for transport, document exposed bone ends.
- Blood loss may be concealed or not apparent with extremity injuries.
- Lacerations must be evaluated for repair within 6 hours of the injury.
- Extremity injuries must be managed and splinted with appropriate splinting device following any immediate interventions required to manage the patient's ABC's.
- Splint, elevate, and cool injured extremity as indicated.

EXTREMITY TRAUMA

SHOCK - TRAUMA

History • Amputation • Blunt or Penetrating Trauma • Multiple Long Bone Fractures • Flail Chest • Major Abdominal Injury • Pelvic Fracture	Signs and Symptoms • Restlessness, thirst • Confusion, change in level of consciousness/mentation • Weak/rapid pulse • Pale, cool, diaphoretic, clammy skin • Hemodynamic instability • Delayed capillary refill • Signs of poor perfusion	Differential • Shock: hypovolemic, cardiogenic, septic, neurogenic, anaphylactic • Ectopic pregnancy • Dysrhythmias • Pulmonary embolus • Tension pneumothorax • Medication effect/Overdose • Vasovagal/Syncope
--	---	--


**NOTES:**

- Once Blood pressure is at 90 systolic for hemorrhagic shock, fluid should be titrated to maintain at 90 and not higher due to risk of increased bleeding
- TXA is contraindicated in: patients under 16, Renal Failure, Allergy to TXA, History of Thromboembolism, Known aneurismal SAH (Sub arachnoid hemorrhage, and in injuries greater than 3 hours old.
- Do not delay fluid bolus for administration of TXA

SHOCK - TRAUMA

TASER REMOVAL


History • Traumatic Injury • Drug Abuse • Cardiac History • History of Asthma • Psychiatric History	Signs and Symptoms • Shortness of Breath • Chest Pain • Numbness/Weakness • Altered LOC • Intoxication/Substance Abuse • Signs of Trauma • Palpitations	Differential (Life Threatening) • Psychiatric Illness • Substance Abuse • Traumatic Injury • Traumatic Brain Injury • Cardiac Dysrhythmia • Spinal injury
--	--	---



TASER REMOVAL

TRAUMA

History - Type of injury - Mechanism of injury, damage to structure or vehicle - Location in vehicle or structure - Others injured or dead - Speed or other details of MVC - Restraints, protective equipment - Past medical history - Medications	Signs and Symptoms - Pain, swelling, - Deformity, lesions, bleeding - Altered mental status or unconscious - Hypotension or shock - Cardio-Respiratory Arrest	Differential (Life Threatening) Chest - Tension Pneumothorax, Flail/ Sucking/Open Chest Wound, Pericardial Tamponade, Hemothorax - Intra-abdominal Bleeding - Pelvis/Femur Fracture - Spine Fracture/Cord Injury - Head Injury (see Head Trauma) - Extremity Fracture/Dislocation - HEENT (Airway Obstruction)
--	--	--




NOTES:

- Mechanism is an indicator of serious injury.
- If transport delayed begin IV fluids on scene, otherwise establish IVs enroute.
- Consider Blood-Y tubing for second IV with Normal Saline.
- Attempt to maintain perfusion with fluid resuscitation, systolic BP of 80-90 mmHg. Systolic BP of > 100 mmHg can lead to increased bleeding at injury site.
- Consider Sheet or commercial pelvic binder device for pelvic fractures if available
- Pregnant trauma patients can lose up to 2 liters of blood before showing signs and symptoms of shock.

TRAUMA

TRAUMATIC BRAIN INJURY

History - Type and time of injury - Mechanism of injury, damage to structure or vehicle - Location in vehicle or structure - Loss of consciousness - Restraints, protective equipment (helmet use, or damage to helmet?) - Evidence of multi-system trauma - Seizures - Preceding events - Past medical history - Medications	Signs and Symptoms - Hematoma, depressions, lacerations - Altered mental status - Unresponsiveness - Nausea/Vomiting - Pupil status - Abnormal respiratory pattern - Apnea - Antegrade or retrograde amnesia - Blood from nose or ears - Exposed brain tissue - Cushing's response	Differential - Traumatic brain injury - Skull fracture - Epidural or subdural hematoma - Spinal injury - Physical abuse/Assault - CVA - Diabetic emergency - Seizure - Syncope - Substance ingestion (drugs, alcohol, other)
---	---	--



NOTES:

- If GCS < 14, consider Air and/or rapid transport to most appropriate facility. Also refer to the Destination Protocol.
- Consider **Trauma Alert**.
- If head injured patient is combative with an unprotected airway – consider pharmacological intervention and intubation.
- Hyperventilate (20/min and/or ETCO₂ 28-32 Torr) the patient **ONLY** if evidence of herniation (blown pupil & flaccidity or decerebrate posturing).
- Hyperventilate (25/min children 35/min for infants and/or PCO₂ 28-32) **ONLY** if evidence of herniation (blown pupil, flaccidity and/or decerebrate posturing). Normal ventilation for others: 20/min for children 30/min for infants. Avoid fluid bolus if isolated head injury.
- 75% of patients with significant head trauma have serious injuries to other organ systems: Do complete assessments.
- Hypotension in **head injury** patients increases mortality by 50%. Titrate fluids to maintain a **systolic BP of at least 100 mmHg** in adults.
- Increased intracranial pressure (ICP) may cause bradycardia and hypertension (Cushing's Response).
- Patients with suspected head trauma should be closely monitored and reassessed for any change in their mental status.
- Anticipate vomiting. Have suction and airway equipment ready and close at hand.
- Scalp lacerations can result in significant blood loss. Apply bulky dressings with moderate pressure PRN.

TRAUMATIC BRAIN INJURY

CHILDBIRTH

History - Length of gestation - Parity and gravidity/Twins? - Previous cesarean delivery - Prenatal care/physician - Alcohol or drug use - Infectious disease status - Previous OB/GYN emergencies (eclampsia, diabetes, premature labor, ectopic pregnancy, etc.)	Signs and Symptoms - Membranes ruptured - Contraction frequency and intensity - Urge to push/bear down - Crowning - Bloody show – mucous plug - Vaginal bleeding - Cramps - Meconium	Differential - Braxton Hicks - Contractions following trauma - Multiple fetuses - Premature - Abdominal pain
---	--	---




NOTES:

- Oxygen should be administered to all mothers during delivery.
- There may still be time to transport to the hospital before delivery, when contractions < 2minutes apart, and patient not crowning.
- Do not rupture membranes unless the baby's head has been delivered and the membranes must be cleared from the mouth and nose.
- The mother may need coaching, support and guidance (breathing, when to push etc.) through the birthing process.
- Abruptio placenta, placenta previa, and ruptured uterus are maternal complications that may be encountered in the pre-hospital setting. These situations may present with severe abdominal pain, hypotension, and/or significant vaginal bleeding. Rapid transport.
- Enlist the help of Midwife, or staff if at a Birthing Facility.

COMPLICATIONS IN CHILDBIRTH

(Multiples/Prolapsed Limb/Nuchal & Prolapsed Cord/Pre-Term)

History - Length of gestation - Parity and gravidity - Previous cesarean delivery - Alcohol or drug use - Infectious disease status - Previous OB/Gyn emergencies (eclampsia, diabetes, premature labor, ectopic pregnancy etc.)	Signs and Symptoms - Membranes ruptured - Contraction frequency and intensity - Urge to push/bear down - Crowning - Bloody show—mucous plug - Vaginal bleeding - Cramps - Meconium	Differential - Braxton Hicks - Contractions following trauma - Multiple fetuses - Premature - Abdominal pain
--	--	---

**NOTES:**


- The above Protocols were developed to serve as a guide for the pre-hospital setting in the event that birth is imminent and complications occur.
- Contact should be made with Medical Control as soon as possible for assistance.
- Rapid Transport for all complications in childbirth.
- A key to neonatal resuscitation is keeping the baby warm.

COMPLICATIONS IN CHILDBIRTH

(Multiple Fetus/Prolapsed Limb/Nuchal Cord/Prolapsed Cord/Pre-Term)

COMPLICATIONS IN CHILDBIRTH (Shoulder Dystocia/Breech)

History	Signs and Symptoms	Differential
- Length of gestation - Parity and gravidity - Previous cesarean delivery - Alcohol or drug use - Infectious disease status - Previous OB/GYN emergencies (eclampsia, diabetes, premature labor, ectopic pregnancy, etc.)	- Membranes ruptured - Contraction frequency and intensity - Urge to push/bear down - Crowning - Bloody show—mucous plug - Vaginal bleeding - Cramps - Meconium	- Braxton Hicks - Contractions following trauma - Multiple fetuses - Premature - Abdominal pain




NOTES:

- Dystocia—difficult birth, may be produced when the size of the fetus is larger than the size of the pelvic outlet.
- During complicated deliveries the fetus may become bradycardic and hypoxic. Complications during birth can be life threatening to the mother and the fetus.
- Rapid transport.
- Prepare to manage a depressed baby. Hypoxia, hypothermia, and hypoglycemia should be addressed. Proceed to the Appropriate Newborn Resuscitation Protocol.

COMPLICATIONS IN CHILDBIRTH (Shoulder Dystocia/Breech)

ECLAMPSIA-PREECLAMPSIA

History • Pregnancy • Advanced maternal age • Chronic hypertension • Chronic renal disease • Diabetes • Lupus • Multiple gestation	Signs and Symptoms • Hypertension - 140/90 mm/Hg or a rise of 20 mm/Hg systolic and 10 mm/Hg diastolic over pregnant BP • Proteinuria • Excessive weight gain with Edema • Headache, dizziness, confusion • Seizure, coma • Blurred vision • Nausea/vomiting • Fetal Distress	Differential • Seizure disorder
---	---	--


**NOTES:**

- Handle the patient GENTLY and minimize sensory stimulation (e.g. darken ambulance lights) to avoid precipitating seizures.
- Eclampsia can occur from 20 weeks gestation and up to 1 month postpartum.
- Preeclampsia may affect previously healthy, normotensive mothers.
- Significant increase in risk to the mother and fetus – TRUE EMERGENCY!
- Place the mother in the left lateral recumbent position to maintain or improve uteroplacental blood flow and to minimize risk of insult to the fetus.
- Anticipate seizures at any moment, and be prepared to provide airway, ventilatory, and circulatory support.
- Eclampsia may be associated with apnea during the seizures.
- Labor can begin spontaneously and progress rapidly.

ECLAMPSIA-PREECLAMPSIA

POST-PARTEM CARE

History • Pregnancy • Length of gestation • Parity and gravidity/Twins? • Previous cesarean delivery • Prenatal care/physician • Alcohol or drug use • Infectious disease status • Previous OB/Gyn emergencies (eclampsia, diabetes, premature labor, ectopic pregnancy, etc.)	Signs and Symptoms • Delivery of baby within 48 hours • Shock • Significant vaginal bleeding • Estimated blood loss (EBL) - # of pads soaked?	Differential • Trauma • Gynecological procedure/Surgery
--	---	---




NOTES:

- This protocol addresses significant hemorrhage after the placenta has delivered. Do not confuse with treatment for patients experiencing significant hemorrhage during delivery.
- Consider Pitocin. Call Medical Control for advice about administering Pitocin
- Rapid transport.

POST-PARTEM CARE

NEWBORN RESUSCITATION - Pediatric

History • Difficult delivery • Fetal distress • Term	Signs and Symptoms • Apnea, Cyanosis • Bradycardia, pulseless • Meconium	Differential • Hypothermia • Hypoglycemia
---	---	---




NOTES:

- **All newborns:** once the body is delivered, dry the baby, replace wet towels with dry ones, and wrap the baby in a thermal blanket or dry towel.
- Cover the head to preserve warmth.
- If infant is already breathing or crying, tracheal suctioning may be omitted if meconium is present.
- Tracheal doses of epinephrine should always be 1:10,000 for newly born.
- **Do not use concentrated doses of medications—cerebral hemorrhage may result.**
- **APGAR** at 1 and 5 minutes.

NEWBORN RESUSCITATION PEDIATRIC

UNIVERSAL PATIENT CARE - Pediatric




NOTES:

- This protocol provides general guidelines for patient management. Refer to additional protocols for treatment of specific conditions.
- A length-based resuscitation tape or other care guide shall be available to assist EMS personnel to quickly determine appropriate equipment size, normal vital signs, and correct medication doses.
- If hazardous conditions are present (such as swift water, hazardous materials, electrical hazard, or confined space) contact an appropriate agency before approaching the patient. Wait for the designated specialist to secure the scene and patient as necessary.
- Reassess the patient frequently.
- Expose the child only as necessary to perform further assessments. Maintain the child's body temperature throughout the examination.
- If the child's condition is critical or unstable, initiate transport. Perform focused history and detailed physical examination enroute to the hospital if patient status and management of resources permit.
- If the child's condition is stable, perform focused history and detailed physical examination on the scene, then initiate transport.
- Contact Medical Control for additional instructions.
- If spinal trauma is suspected, continue manual stabilization, place in rigid cervical collar, and apply an immobilization device.
- PAT – Pediatric Assessment Triangle: Appearance/Work of Breathing/Circulation.

UNIVERSAL PATIENT CARE PEDIATRIC

AIRWAY MANAGEMENT - Pediatric

History	Signs and Symptoms of Difficult Airway	Differential
• Trauma • Head Injury • Asthma • COPD • Known difficult airway • Facial fractures • Pulmonary edema	• Hoarseness • Limited neck movement • Limited mouth opening • Short thyro-mental distance • Short heavy neck, obesity • Receding mandible/overbite • Large swollen tongue • Long incisors	• LOC • Airway injury • Airway swelling • Burns • Foreign body • Epiglottitis


**NOTES:**

- Intubation is not necessarily required for adequate ventilation, which is the goal. BVM may be adequate.
- Keep it simple...when possible – use progressively invasive maneuvers **ONLY** when necessary.
- Clinical End-Tidal CO₂ (capnography) monitoring should be used with all advanced airways.
- Only use hyperventilation for head injury when signs of herniation are present – ventilate to torr CO₂ of 28 - 32.
- Assess for signs of respiratory distress, failure, or arrest. If present, refer to the appropriate protocol for treatment options.
- If the child is not breathing or breathing is inadequate, initiate assisted ventilation using a bag-valve-mask device with high flow, 100% oxygen. Begin with 2 slow, deep breaths of about 1-1/2 seconds duration until chest rises, then ventilate at 20 breaths/minute for all ages (except neonates at 40). If abdominal distention arises, consider placing an orogastric tube to decompress the stomach.
- If breathing is adequate administer oxygen to an oxygen saturation of 93 – 99% in infants and children (see newborn for administration to newborns)

AIRWAY MANAGEMENT PEDIATRIC

ALLERGIC REACTION – ANAPHYLAXIS - Pediatric

History - Known allergic reaction to bites, stings, food, medications etc. - Possible ingestion of or contact with allergen.	Signs and Symptoms - Dyspnea, often with sneezing, wheezing, or coughing - Facial swelling - Urticaria - Abdominal cramps - Nausea, vomiting, diarrhea - Tachycardia - Falling blood pressure	Differential - Asthma - Pulmonary embolism
--	--	--


**NOTES:**

- Consider Epinephrine IM, diphenhydramine early in the allergic process, administration prior to histamine release will provide more rapid results. When signs of histamine release are noted, the process is well under way and will require aggressive treatment.
- Epinephrine has a short half-life and may require repeat doses.
- Closely monitor patients for rebound signs and symptoms. Any patients suffering from an allergic reaction should be evaluated by a physician.
- For patients with signs of anaphylaxis – hypotensive, despite treatment, **consult medical control** for a glucagon order. Can be repeated every 5 minutes until hypotension resolves.

ALLERGIC REACTION—ANAPHYLAXIS PEDIATRIC

ASTHMA - Pediatric

History - Asthma - COPD: Emphysema, Bronchitis - CHF: Congestive Heart Failure - Home Oxygen use - Home Nebulizer Use - Medications: Steroids, Inhalation, Possible Chemical or biological exposure	Signs and symptoms - Shortness of breath - Pursed-Lips breathing - Accessory muscle use, retractions, nasal flaring, fatigue - Inability to speak in sentences - Audible Wheezing or rhonchi - Fever, cough - Cyanosis - Lung sounds: Wet? Diminished? Bilaterally? Expiratory Wheezing?	Differential Asthma, COPD CHF, Pulmonary Edema - Anaphylaxis Pneumonia Pulmonary Embolus Cardiac - Hyperventilation Inhaled toxin - DKA Pneumothorax Epiglottitis, Croup
---	--	---




Notes:

- In severe distress, treatment may occur simultaneous with IV, EKG. Consideration of Mag Sulfate in the updraft or IV as directed by Medical Control.
- Remember: almost all cardiac problems produce some degree of respiratory distress.
- Pulse Oximetry should be monitored continuously for all patients with respiratory distress and/or respiratory failure.
- Patients with a history of asthma, who have had prior hospitalization for asthma, and/or present with initial O₂ saturations of <90% are at increased risk for rapid decline in spite of initial improvement with your treatments.
- A silent chest in the setting of severe respiratory distress is a pre-respiratory arrest sign.
- Versed may be administered prior to intubation of a conscious patient who is *in extremis* and has not responded to treatment.
- Use all available personal protective equipment and clothing if toxic inhalation or exposure is a possible etiology.
- Provide high flow O₂ and transport for patients who are hyperventilating when the cause is unknown.

ASTHMA - PEDIATRIC

ASYSTOLE/PULSELESS ELECTRICAL ACTIVITY - Pediatric

History - Events leading up to arrest - Estimated down time - Past medical history/ medications - Renal failure/dialysis - DNR - Hypothermia - Suspected Overdose (Digitalis, Tricyclics, Beta-blockers, Calcium channel blockers) - Respiratory failure	Signs and Symptoms Unresponsive, Apneic, Pulseless with organized electrical activity	Differential - Medical vs. Trauma etiology - Hypovolemia (Trauma, AAA, GI) - Hypothermia - Drug Overdose - Massive Myocardial Infarction - Hypoxia - Tension Pneumothorax - Pulmonary Embolism - Acidosis - Hyperkalemia - Device error (lead off) - Death
--	--	--



NOTES:

- For trauma patients determine the underlying cause of arrest and provide definitive treatment i.e. fluid resuscitation, pleural decompression.
- Reassess Advanced Airway placement frequently, i.e. after every patient move, change in patient condition.
- For hypothermic patients pharmacologic treatment may not be effective until patient is warmed; see Hypothermia Protocol.
- Considerations for Sodium Bicarb-known preexisting hyperkalemia, bicarbonate responsive acidosis (e.g. Diabetic ketoacidosis), or overdose (e.g. Tricyclics, cocaine, diphenhydramine) to alkalinize the urine in aspirin or other overdose.
- Atropine administration is indicated for poisoning from organo phosphates

ASYSTOLE/ PULSELESS ELECTRICAL ACTIVITY- PEDIATRIC

BRADYCARDIA - Pediatric

History • Respiratory • Medications: Beta Blockers, Clonidine, Calcium Channel Blockers, Digitalis • Pacemaker • Nausea and Vomiting • Past cardiac history	Signs and Symptoms • HR < 60/min • Respiratory distress/Failure • Nausea/Vomiting • Cardiopulmonary Failure • Hypotension • Decreased LOC • Weakness	Differential • Respiratory Failure/Hypoxia • Hypothermia • Toxic Exposure/Overdose • Head Injury • Vasovagal • Cardiac • CVA
--	--	---


**NOTES:**

- Respiratory failure is the usual cause of bradycardia in pediatric patients—assist ventilations.
- Epinephrine is more effective than Atropine for hypoxic bradycardia.
- CPR should be started if ventilation fails to improve heart rate.
- Atropine is not indicated unless history of heart disease or vagal cause of bradycardia is suspected.
- Attempting to increase the rate of an asymptomatic patient is contraindicated.
- Adult pads should be used in patients down to the age of 1 year so long as they do not touch; If adult pads are too large, use pediatric pads.
- Versed 0.1 mg/kg IV, max of 5 mg or appropriate Benzodiazepine, may be used as a sedative agent when needed for pacing discomfort.

BRADYCARDIA PEDIATRIC

DIABETIC EMERGENCIES - Pediatric

History - Known diabetic, medic alert tag - Change in routine, meds, or diet - Recent illness or infection - Possible illicit drug use - Alcohol abuse - Medications - History of trauma	Signs and Symptoms - Decreased mental status/Seizures - Bizarre behavior - Cool diaphoretic skin - Fruity, ketotic breath - Kussmaul respirations - Signs of dehydration - Excessive thirst, hunger, or urination	Differential - Head trauma - CVA, seizure, sepsis - Cardiac - Shock - Toxic ingestion/alcohol intoxication - Environmental exposure - Psychiatric disorder
---	--	---




NOTES:

- If Glucagon is given and patient responds, remember that glucagon depletes glucose stores so dextrose must be administered soon.
- Perform blood glucose checks on ALL patients with altered mental status.
- Glucometer reading from 60–80 in a patient with serious symptoms may indicate hypoglycemia—Administer Dextrose.
- If in doubt about glucometer reading—administer Dextrose.
- Consider oral glucose in the alert diabetic patient who is expected to maintain his/her own airway.
- Consult Medical Control for Thiamine administration for patients suspected of malnutrition i.e. history of chemotherapy, etc.
- Perform blood glucose checks on all seizure patients. Undiagnosed DKA and hypoglycemia from other causes can precipitate seizure activity.
- Consider endotracheal intubation in patients with altered blood glucose levels who do not respond to Dextrose and Narcan.
- Ascertain the patient's insulin regimen (dose, type, & schedule) for ED reference.

DIABETIC EMERGENCIES PEDIATRIC

MULTIPLE TRAUMA - Pediatric

History	Signs and Symptoms	Differential (Life Threatening)
- Type of injury - Mechanism of injury, damage to structure or vehicle - Location in vehicle or structure - Others injured or dead - Speed or other details of MVC - Restraints, child seats, & helmets - Past medical history - Medications	- Pain, swelling, - Deformity, lesions, bleeding - Altered mental status or unconscious - Hypotension or shock - Cardio-Respiratory Arrest - Altered PAT - Bruising or hematomas	- Chest - Tension Pneumothorax, Flail Chest, Sucking/Open Chest Wound, Pericardial Tamponade, Hemothorax - Intra-abdominal Bleeding - Pelvis/Femur Fracture - Spine Fracture/Cord Injury - Head Injury (see Head Trauma) - Extremity Fracture/Dislocation - HEENT (Airway Obstruction)




NOTES:

- Approximately 60% of multiple trauma patients have a concomitant head injury.
- Unrecognized hemorrhage the leading cause of preventable death in trauma care. Increasing heart rates often reflect untreated hemorrhage.
- Maintain perfusion with fluid resuscitation, systolic BP of $70 + 2 \times \text{age}$ if over 1 year old. Increased BP can cause increased bleeding at injury site.
- Mechanism of injury is the earliest predictor of serious injury.
- If transport delayed begin IV fluids on-scene, otherwise establish enroute. Consider Blood-Y tubing for second IV.
- Remove seriously injured children from the child seat if potentially damaged in the crash. Seriously injured children require supine immobilization.
- Attempt to keep siblings, parents, and or friends together.

MULTIPLE TRAUMA PEDIATRIC

OVERDOSE/TOXIC EXPOSURE - Pediatric

History	Signs and Symptoms	Differential
- Suspected toxic exposure - Age of patient - Substance, route, quantity, time - Reason (suicidal, accidental, criminal, terrorism), prior history - Available medications in home - Past medical history, medications	- Mental status changes - Hypotension/Hypertension - Decreased respiratory rate - Tachycardia, dysrhythmias - Seizures - Pupils status - Signs of illicit drug use	- Reasons for Coma (AEIOUTIPS) - Tricyclic antidepressants - Acetaminophen (Tylenol) - Depressants - Stimulants - Anticholinergic - Cardiac medications - Solvents, Alcohols, Cleaning Agents,

**NOTES:**

- Many ingestions in under 5 year age group are minor—usually single agent ingestions. Assess thoroughly!
- Do **not** force administration of oral antidotes or NG/OG tube in the alert/semi-alert child.
- Teenagers often have multiple agent ingestions. Assess for trauma and suicide attempts.
- Perform ET tube placement prior to NG/OG tube in unresponsive patients.
- Do not rely on patient history of ingestion, especially in suicide attempts.
- Attempt to bring bottles, contents, emesis to ED.
- Consider restraints if necessary for patient's and/or personnel protection.
- Consider Calling Poison Control – American Association of Poison Control – 800-222-1222


- Cardiac Meds:** dysrhythmias and mental status changes
- Tricyclic Antidepressants:** 4 major areas of toxicity—seizures, dysrhythmias, hypotension, decreased mental status or coma; Rapid progression from alert mental status to death.
- Acetaminophen:** Initially normal or N/V. If not detected and treated, causes irreversible liver failure.

- Depressants:** ↓HR, ↓BP, ↓ respirations, ↓ temperature, nonspecific pupils.
- Stimulants:** ↑HR, ↑BP, ↑respirations, ↑ temperature, dilated pupils, seizure.
- Beta Blockers/Ca⁺ Channel Blocker** ↓ HR ↓ BP give glucagon

OVERDOSE/TOXIC EXPOSURE PEDIATRIC

POST RESUSCITATION - Pediatric

History • Cardiac Arrest • Respiratory Arrest	Signs and Symptoms • Return of Pulse • Increasing heart rate • Skin color change	Differential • Continue to address specific differentials associated with original dysrhythmia
---	---	---




NOTES:

- Find and treat cause of cardiac arrest. Respiratory failure is the most frequent cause in pediatric patients.
- A 12 lead EKG should be obtained as soon as possible to determine the presence of an acute coronary syndrome.
- Continuously recheck tube placement. Secure tube then immobilize patient with CID to prevent tube dislodgement.
- Assess ventilation and respiratory status— treat as indicated.
- Benzodiazepines may be used for sedation in order to maintain a controlled airway.
- Narrow Complex Tachycardia in the post resuscitation phase may be due to epinephrine therapy and usually does not require treatment— monitor BP.
- Consider OG tube placement for gastric decompression.
- Consider temperature regulation. Correct hyperthermia, allow **mild** hypothermia.

SEIZURE - Pediatric

History • Documented seizure disorder • Medications • Pregnancy • Trauma – Recent or Remote • Recent illness • Diabetes • Fever	Signs and Symptoms • Decreased mental status • Sleepiness • Incontinence • Observed seizure activity • Evidence of Trauma • Photophobia • Increased sensitivity to touch and sound	Differential • Fever • Hypoxia • Hypoglycemia • CNS Injury or Tumor • Eclampsia • Renal failure • Drug use • Infection • Alcohol/illicit drug withdrawal • Metabolic disorder • Electrolyte imbalance
--	---	--




NOTES:

- Short-term **Febrile seizures** in infants and children are relatively benign; most common cause of seizure in pediatric patient—should be transported to the ED for physician evaluation. Rapid change in temperature is typical cause. Evaporative cooling to bring temp down.
- Anticonvulsants should only be used when patient has ACTIVE, CONTINUOUS seizures or no ALERT period between seizures.
- Hypoxia & hypoglycemia during status can cause permanent brain damage—ensure good airway breathing circulation and blood sugar.
- Status may exist if patient continues to have any focal seizure activity after generalized seizure (the brain may still be seizing).
- Be prepared to control airway and assist ventilation; consider nasal trumpet airway and nasal intubation for patients with clenched jaw.
- Assess possibility of recent traumatic event and drug abuse or toxic exposure (i.e. stimulants)
- Consider positioning the patient in lateral recumbent, recovery position.

SEIZURE PEDIATRIC

STRIDOR (UPPER AIRWAY OBSTRUCTION) - Pediatric

History - Recent Intubation - Respiratory syncytial virus (RSV) - Respiratory Failure - Inhaled Toxins - Bronchiolitis - Medications: Steroids, Inhalation, Possible Chemical or biological exposure	Signs and symptoms - Shortness of breath - Tripod positioning - Neck extended - Drizzling - Barking cough - Pursed-Lips breathing - Accessory muscle use, retractions, nasal flaring, fatigue - Inability to speak in sentences - Audible stridor - Cyanosis	Differential - Epiglottitis, Croup - CHF, Pulmonary Edema - Anaphylaxis - Pneumonia - Pulmonary Embolus - Cardiac - Hyperventilation - DKA - Pneumothorax - Asthma, COPD
--	--	--


**Notes:**

- If heart rate increases greater than 20 beats per minute while administering nebulized Racemic Epinephrine or Epi 1:1,000 the further dilute the treatment or stop administration.
- Stridor is caused by narrowing of the upper airway structures above the carina. Prompt identification and proper treatment is imperative.
- Stridor is commonly noted during Anaphylaxis, Croup, trauma to the trachea, or burns to the upper airway.
- Beta 2 agonist/Bronchodilators have little to no effect when used to treat Stridor.
- Pulse Oximetry should be monitored continuously for all patients with respiratory distress and/or respiratory failure.
- A silent chest in the setting of severe respiratory distress is a pre-respiratory arrest sign.
- Use all available personal protective equipment and clothing if toxic inhalation or exposure is a possible etiology.
- Provide humidified oxygen for pediatric patients in stridor.

STRIDOR (UPPER AIRWAY OBSTRUCTION) - PEDIATRIC

TACHYCARDIA - Pediatric

History • Medications (Aminophylline, decongestants, thyroid supplements, diet pills, Digoxin) • Diet, N/V, dehydration • Congenital Heart Disease • Illicit drugs (methamphetamine, cocaine, stimulants) • Past Medical History • History of palpitations/heart racing • Syncope • Near Drowning	Signs and Symptoms • HR > 180/min • QRS < 0.08 sec vs. > 0.08 sec • Dizziness, chest pain, shortness of breath • Sudden onset • Poor perfusion • Potential presenting rhythm: • Sinus Tachycardia • Atrial Fibrillation/ Flutter • PSVT • Ventricular Tachycardia	Differential • Congenital Heart Disease (WPW, Long QT syndrome, Valvular) • Myocardial Infarction • Electrolyte imbalance • Exertion, pain, emotional stress, fever • Hypoxia • Hypovolemia or anemia • Drug effect/overdose • Hyperthyroidism • Pulmonary Edema
---	--	---




NOTES:

- Abnormal tachycardias in children are rare. Rate changes with activity, respirations in Sinus Tach.
- Establish rapid heart rate as cause of signs and symptoms.
- Note/record EKG changes during **Vagal** maneuvers and Adenosine administration.
- Signs of poor perfusion: LOC change, weak or absent radial pulses, poor capillary refill, pale, mottled or cyanotic skin, or low blood pressure.
- **A child with narrow QRS Tach: Dehydration or volume depletion usually indicate Sinus Tachycardia. Do not use this protocol. Use Hypotension Protocol.**
- Promptly cardiovert hemodynamically unstable, the more unstable the patient, the more urgent the need for cardioversion.
- Continuous pulse oximetry for all Tachycardia patients.
- Document all rhythm changes and therapeutic interventions with EKG strips.
- Calcium Channel Blockers (Cardizem, Verapamil etc.) are contraindicated in patients with WPW.
- QRS > 0.08 sec (2 little squares) means Ventricular Tachycardia.

VENTRICULAR FIBRILLATION/PULSELESS V-TACH - Pediatric

History - Estimated down time - Past medical history/ medications - Events leading to arrest - Hypothermia - Electrocution - Toxins - Heart Hx/ WPW / Long QT syndrome	Signs and Symptoms Unresponsive, Apneic, pulseless	Differential - Medical vs. Trauma etiology - Artifact - Asystole - Device failure
---	---	---



NOTES:

- Pattern should be drug- CPR- shock, drug- CPR- shock, etc.
- Reassess ETT placement frequently, i.e. after every patient move, change in patient condition.
- Search for cause of Cardiac Arrest.
- If defibrillation is successful and patient rearrests, return to previously successful Joule setting and administer antiarrhythmic medicine.
- Defibrillation takes precedence over all treatment once the defibrillator is available.
- For hypothermic patients defibrillation may not be effective, see Hypothermia Protocol.
- Spinal immobilize electrocution patients.
- For trauma patients determine the underlying cause of arrest and provide definitive treatment i.e. fluid resuscitation, pleural decompression.
- If patient successfully converted with Defibrillation: Start antiarrhythmic therapy as above.
- Considerations for Sodium Bicarb: known preexisting hyperkalemia, bicarbonate responsive acidosis (e.g. Diabetic ketoacidosis), or overdose (e.g. Tricyclics, cocaine, diphenhydramine) to alkalinize the urine in aspirin or other overdose.
- Adult pads should be used in patients down to the age of 1 year so long as they do not touch; If adult pads are too large, use pediatric pads.

VENTRICULAR FIBRILLATION/PULSELESS V-TACH PEDIATRIC

MEDICATION SECTION

This section is protocol. Not all medications listed here will be carried or used by your service. Drugs included here are listed because they are referenced in the previous protocol section either in the diagram or in the notes section.

Please see your service policy on which drugs you carry and use for the individual protocols in the previous section

It is necessary to have multiple drugs for individual protocols due to drug shortages and availability.

Acetaminophen

NAME	Acetaminophen (Tylenol, APAP)				
CLASS	Analgesic, antipyretic				
ACTION	Acetaminophen has analgesic and antipyretic properties with effects equivalent to those of aspirin. Its analgesic and antipyretic effects are likely the result of the inhibition of prostaglandin E2 (PGE2). It does not fall into the non-steroidal anti-inflammatory class of medications as it has no anti-inflammatory effects. Acetaminophen acts on a variant of cyclooxygenase (COX3) that is only expressed in the central nervous system. Unlike aspirin, it has no effect on COX1 or COX2, and therefore, it has no effect on platelets. Acetaminophen elevates the pain threshold and readjusts the hypothalamic temperature regulatory center.				
ONSET/DURATION	<table><tr><td>Oral</td></tr><tr><td>Onset: 30-60 minutes Duration: 4-6 hours </td></tr></table> <table><tr><td>IV</td></tr><tr><td>Onset 5-10 minutes Duration - The peak analgesic effect is reached at approximately 1 hour. - The duration of action for analgesia is 4 to 6 hours </td></tr></table>	Oral	Onset: 30-60 minutes Duration: 4-6 hours	IV	Onset 5-10 minutes Duration - The peak analgesic effect is reached at approximately 1 hour. - The duration of action for analgesia is 4 to 6 hours
Oral					
Onset: 30-60 minutes Duration: 4-6 hours					
IV					
Onset 5-10 minutes Duration - The peak analgesic effect is reached at approximately 1 hour. - The duration of action for analgesia is 4 to 6 hours					
INDICATIONS	- Mild to moderate pain - Fever - over 38.0 Celsius taken orally or rectally for those under 2 years of age - Febrile seizure				
CONTRAINDICATIONS	- Known hypersensitivity - Environmental hyperthermia - Severe hepatic impairment - Liver disease - Less than 2 months old				

Acetaminophen

ADVERSE REACTIONS	• Nausea/vomiting • Skin irritation • Hearing impairment • Possible GI irritation • Anaphylaxis • Liver problems • Eye irritation • Sores in mouth and nose • Stevens-Johnson Syndrome (rare) • Toxic epidermal necrolysis (rare)												
DOSE AND ROUTE	<table border="1"> <tr> <th colspan="2"><u>Oral</u></th></tr> <tr> <td data-bbox="760 737 906 1150"> Adult </td><td data-bbox="906 737 1442 1150"> Tablet dosing by weight (For patients able to swallow pills) • 21 – 33kg of body weight give 325mg by mouth • 33 – 43kg of body weight give 500mg by mouth • 43 – 66kg of body weight give 650mg by mouth • > 66kg of body weight give 1,000mg by mouth </td></tr> <tr> <td data-bbox="760 1150 906 1276"> Pedi </td><td data-bbox="906 1150 1442 1276"> • 15 ml/kg/dose every 4-6 hours • Max liquid dose of 1,000 mg Round to the nearest ML </td></tr> </table> <table border="1"> <tr> <th colspan="2"><u>IV</u></th></tr> <tr> <td data-bbox="760 1472 906 1535"> Adult </td><td data-bbox="906 1472 1430 1535"> 1g IV </td></tr> <tr> <td data-bbox="760 1535 906 1598"> Pedi </td><td data-bbox="906 1535 1430 1598"> 15mg/kg IV (max 1g) </td></tr> </table>	<u>Oral</u>		Adult	Tablet dosing by weight (For patients able to swallow pills) • 21 – 33kg of body weight give 325mg by mouth • 33 – 43kg of body weight give 500mg by mouth • 43 – 66kg of body weight give 650mg by mouth • > 66kg of body weight give 1,000mg by mouth	Pedi	• 15 ml/kg/dose every 4-6 hours • Max liquid dose of 1,000 mg Round to the nearest ML	<u>IV</u>		Adult	1g IV	Pedi	15mg/kg IV (max 1g)
<u>Oral</u>													
Adult	Tablet dosing by weight (For patients able to swallow pills) • 21 – 33kg of body weight give 325mg by mouth • 33 – 43kg of body weight give 500mg by mouth • 43 – 66kg of body weight give 650mg by mouth • > 66kg of body weight give 1,000mg by mouth												
Pedi	• 15 ml/kg/dose every 4-6 hours • Max liquid dose of 1,000 mg Round to the nearest ML												
<u>IV</u>													
Adult	1g IV												
Pedi	15mg/kg IV (max 1g)												
NOTES													

NAME	ADENOSINE (Adenocard)
CLASS	Endogenous nucleoside, miscellaneous antidysrhythmic agent
ACTION	Adenosine slows supraventricular tachycardias (SVTs) by decreasing electrical conduction through the AV node without causing negative inotropic effects.
ONSET/DURATION	Onset: Immediate Duration: 10 seconds
INDICATIONS	Diagnosis and treatment of SVT, in adults and pediatric patients. Monomorphic Wide Complex Tachycardia
CONTRAINDICATIONS	• Second or third degree AV block, or sick sinus syndrome • Hypersensitivity to adenosine • Atrial flutter, atrial fibrillation, ventricular tachycardia (Adenosine is usually not effective in converting these rhythms to sinus rhythm) • Wolff-Parkinson-White syndrome
ADVERSE REACTIONS	• Hypotension • Shortness of Breath • Transient periods of sinus bradycardia, sinus pause or bradyasystole • Nausea
DOSE AND ROUTE	Adult: Rx: SVT & Monomorphic Wide Complex Tachycardia: 6 mg IV rapidly over 1-3 seconds. Flush with 20 ml bolus of NS, elevate IV arm. If no effect in 1-2 minutes, give 12 mg over 1-3 seconds. May repeat 12 mg bolus one more time. Pediatric: Rx: SVT & Monomorphic Wide Complex Tachycardia: 0.1-0.2 mg/kg IV, IO rapidly, up to 6 mg. If no effect, may double dose. Maximum: 12 mg total dose.
NOTES	Run monitor strip while administering Adenosine. This will often allow you to see the underlying rhythm when the rate slows. If underlying rhythm is Atrial Fib or Atrial Flutter, discontinue Adenosine and consult medical control. Stimulants such as caffeine may cause Adenosine to be ineffective.

Albuterol

NAME	Albuterol (Proventil, Ventolin)												
CLASS	Sympathomimetic, Bronchodilator, Beta 2 agonist												
ACTION	- Albuterol is a sympathomimetic that is selective for beta 2 adrenergic receptors. It relaxes the smooth muscle of the bronchial tree and peripheral vasculature by stimulating adrenergic receptors of the sympathetic nervous system - Albuterol has an additive effect with insulin and glucose, which helps shift potassium into the intracellular space in cases of hyperkalemia.												
ONSET/DURATION	Onset: 5-15 minute after inhalation Duration: 3-4 hours after inhalation												
INDICATIONS	- Relief of bronchospasm in patients with reversible obstructive airway disease In: COPD, Asthma, Pulmonary Edema, Allergic Reaction. - Compartment Syndrome, Hyperkalemia												
CONTRAINDICATIONS	- Prior hypersensitivity reaction to albuterol. - Cardiac dysrhythmias associated with tachycardia.												
ADVERSE REACTIONS	- Usually dose related - Restlessness, apprehension - Dizziness - Palpitations, tachycardia - Dysrhythmias												
DOSE AND ROUTE	<table> <tr> <td>Adult</td><td> <table> <tr> <td>Bronchospasm - Asthma - COPD - Allergic reaction </td><td>2.5mg</td></tr> <tr> <td>Hyperkalemia</td><td>10mg</td></tr> </table> </td></tr> <tr> <td>Pedi</td><td> - Use adult dosage for pediatric patients greater than 1 year of age. <table> <tr> <td>Bronchospasm - Asthma - COPD - Allergic reaction </td><td>0.03ml/kg maxim 1 ml of fluid</td></tr> <tr> <td>Hyperkalemia</td><td>10mg</td></tr> </table> </td></tr> </table>	Adult	<table> <tr> <td>Bronchospasm - Asthma - COPD - Allergic reaction </td><td>2.5mg</td></tr> <tr> <td>Hyperkalemia</td><td>10mg</td></tr> </table>	Bronchospasm - Asthma - COPD - Allergic reaction	2.5mg	Hyperkalemia	10mg	Pedi	- Use adult dosage for pediatric patients greater than 1 year of age. <table> <tr> <td>Bronchospasm - Asthma - COPD - Allergic reaction </td><td>0.03ml/kg maxim 1 ml of fluid</td></tr> <tr> <td>Hyperkalemia</td><td>10mg</td></tr> </table>	Bronchospasm - Asthma - COPD - Allergic reaction	0.03ml/kg maxim 1 ml of fluid	Hyperkalemia	10mg
Adult	<table> <tr> <td>Bronchospasm - Asthma - COPD - Allergic reaction </td><td>2.5mg</td></tr> <tr> <td>Hyperkalemia</td><td>10mg</td></tr> </table>	Bronchospasm - Asthma - COPD - Allergic reaction	2.5mg	Hyperkalemia	10mg								
Bronchospasm - Asthma - COPD - Allergic reaction	2.5mg												
Hyperkalemia	10mg												
Pedi	- Use adult dosage for pediatric patients greater than 1 year of age. <table> <tr> <td>Bronchospasm - Asthma - COPD - Allergic reaction </td><td>0.03ml/kg maxim 1 ml of fluid</td></tr> <tr> <td>Hyperkalemia</td><td>10mg</td></tr> </table>	Bronchospasm - Asthma - COPD - Allergic reaction	0.03ml/kg maxim 1 ml of fluid	Hyperkalemia	10mg								
Bronchospasm - Asthma - COPD - Allergic reaction	0.03ml/kg maxim 1 ml of fluid												
Hyperkalemia	10mg												

Albuterol

NOTES	• Some physicians recommend albuterol updrafts in patients with possible pneumonia and/or CHF. When in doubt consult medical control. • Because albuterol increases heart rate, it should be used in caution in patients with tachycardia with signs and symptoms of AMI. • Albuterol can be given in continuous updrafts to both adult and pediatric patients with severe bronchospasm or reactive airway disease.
-------	---

Amiodarone

NAME	Amiodarone							
CLASS	Class III Antidysrhythmic							
ACTION	Amiodarone is a unique antidysrhythmic agent with multiple mechanisms of action. The drug prolongs duration of the action potential and effective refractory period.							
ONSET/DURATION	Onset: within minutes Duration: Variable							
INDICATIONS	Initial treatment and prophylaxis of frequent recurring VF and hemodynamically stable VT in patients refractory to other therapy.							
CONTRAINDICATIONS	● Sensitivity to Amiodarone ● Pulmonary congestion ● Cardiogenic shock ● Hypotension ● 2nd or 3rd degree block ● Sensitivity to iodine ● If the patient is taking other drugs that prolong the QT such as procainamide. ● QT interval greater than 0.46							
ADVERSE REACTIONS	● Hypotension ● Headache ● Dizziness ● Bradycardia ● AV conduction abnormalities ● Flushing ● Abnormal salivation							
DOSE AND ROUTE	<table><tr><td rowspan="2">Adult</td><td>Cardiac arrest</td><td>VF/VT: 300 mg IV/IO may repeat 150 mg IV/IO in 3-5 minutes</td></tr><tr><td>Stable Wide Complex Tachycardia with a pulse</td><td>Rapid Infusion: 150 mg IV over 10 minutes (mix 150 mg in 100ml of ND or D5W and run at 10ml/min (if available set pump to 600 ml per hour.) May repeat 150 mg in 10 minutes. (monitor BP closely and if it drops below 90 systolic withhold amiodarone until medical control is contacted.)</td></tr></table>			Adult	Cardiac arrest	VF/VT: 300 mg IV/IO may repeat 150 mg IV/IO in 3-5 minutes	Stable Wide Complex Tachycardia with a pulse	Rapid Infusion: 150 mg IV over 10 minutes (mix 150 mg in 100ml of ND or D5W and run at 10ml/min (if available set pump to 600 ml per hour.) May repeat 150 mg in 10 minutes. (monitor BP closely and if it drops below 90 systolic withhold amiodarone until medical control is contacted.)
Adult	Cardiac arrest	VF/VT: 300 mg IV/IO may repeat 150 mg IV/IO in 3-5 minutes						
	Stable Wide Complex Tachycardia with a pulse	Rapid Infusion: 150 mg IV over 10 minutes (mix 150 mg in 100ml of ND or D5W and run at 10ml/min (if available set pump to 600 ml per hour.) May repeat 150 mg in 10 minutes. (monitor BP closely and if it drops below 90 systolic withhold amiodarone until medical control is contacted.)						

Amiodarone

	<table><tr><td rowspan="2">Pedi</td><td>SVT, VT (with pulses)</td><td>5 mg/kg IV/IO load over 20 to 60 minutes. (max 300mg) may repeat, max 15mg/kg</td></tr><tr><td>V-fib/pulseless VT:</td><td>5 mg/kg IV/IO bolus (max 300mg) may repeat, max 15mg/kg</td></tr></table>	Pedi	SVT, VT (with pulses)	5 mg/kg IV/IO load over 20 to 60 minutes. (max 300mg) may repeat, max 15mg/kg	V-fib/pulseless VT:	5 mg/kg IV/IO bolus (max 300mg) may repeat, max 15mg/kg
Pedi	SVT, VT (with pulses)		5 mg/kg IV/IO load over 20 to 60 minutes. (max 300mg) may repeat, max 15mg/kg			
	V-fib/pulseless VT:	5 mg/kg IV/IO bolus (max 300mg) may repeat, max 15mg/kg				
NOTES	• Amiodarone may also be helpful for controlling the ventricular rate in rapid atrial dysrhythmias in patients with severely impaired left ventricular function. • If pregnancy is suspected the use of lidocaine is preferred over amiodarone.					

Aspirin (ASA)

NAME	Aspirin (ASA)				
CLASS	Analgesic, anti-inflammatory, antipyretic, antiplatelet				
ACTION	Aspirin blocks pain impulses in the CNS, dilates peripheral vessels and decreases platelet aggregation. The use of aspirin is strongly recommended for all acute MI patients.				
ONSET/DURATION	Onset: 15-30 minutes Duration: 4-6 Hours				
INDICATIONS	• AMI • Prevention of platelet aggregation in ischemia and thromboembolism. • Unstable angina/Chest Pain of cardiac Origin				
CONTRAINDICATIONS	• Hypersensitivity to salicylates • GI bleeding • Active ulcer disease • Stroke • Bleeding disorders • Children with flu like symptoms (19 Y/O and younger) • Possible dissecting thoracic aneurysm				
ADVERSE REACTIONS	• Stomach irritation • Heartburn or indigestion • Nausea and vomiting • Allergic reaction • Asthma induced by NSAIDS				
DOSE AND ROUTE	<table border="1"> <tr> <td>Adult</td><td>Acute Myocardial Infarction: 160-325mg PO (2-4 low dose ASA)</td></tr> <tr> <td>Pedi</td><td>Not recommended</td></tr> </table>	Adult	Acute Myocardial Infarction: 160-325mg PO (2-4 low dose ASA)	Pedi	Not recommended
Adult	Acute Myocardial Infarction: 160-325mg PO (2-4 low dose ASA)				
Pedi	Not recommended				

NOTES

- Adult patients should all be carefully assessed for signs and symptoms of thoracic aneurysm and/or GI bleeding prior to administration of aspirin in the prehospital setting.

<u>NAME</u>	ATROPINE SULFATE				
<u>CLASS</u>	Anticholinergic agent, Vagolytic				
<u>ACTION</u>	Atropine sulfate (a potent parasympatholytic), inhibits actions of acetylcholine at postganglionic parasympathetic (primarily muscarinic) receptor sites. In emergency care, it is primarily used to increase the heart rate in life-threatening or symptomatic bradycardia and to antagonize excess muscarinic receptor stimulation caused by organophosphate insecticides or chemical nerve agents.				
<u>ONSET/DURATION</u>	<table><tr><td>Onset: Rapid</td></tr><tr><td>Duration: 2-6 Hours</td></tr></table>	Onset: Rapid	Duration: 2-6 Hours		
Onset: Rapid					
Duration: 2-6 Hours					
<u>INDICATIONS</u>	• Hemodynamically significant bradycardia • Organophosphate or nerve gas poisoning				
<u>CONTRAINDICATIONS</u>	• Tachycardia • Hypersensitivity to atropine				
<u>ADVERSE REACTIONS</u>	• Tachycardia • Paradoxical bradycardia when pushed too slowly or when used at doses less than 0.5 mg (Adult) or 0.1 mg (Pediatric) • Palpitations • Dysrhythmias • Headache • Dizziness • Nausea and Vomiting • Flushed, hot, dry skin • Allergic reactions				
<u>DOSE AND ROUTE</u>	<table><tr><td>Adult</td><td><u>Rx: Symptomatic Bradycardia:</u> 1 mg IV push fast q 3-5 minutes; up to 3 mg total dose <u>Rx: Organophosphate or Carbamate Insecticide poisoning:</u> 2-5 mg IV/IO q 5 - 10 minutes until secretions dry or HR increases (No max dose in OPP)</td></tr><tr><td>Pedi</td><td><u>Rx: Bradydysrhythmias:</u> 0.02 mg/kg IV, IO. The minimum dose is 0.1 mg. May be repeated in 5 minutes for a maximum total dose of 1.0 mg for a child and 2.0 mg for an adolescent <u>Rx: Organophosphate or Carbamate Insecticide poisoning:</u> 0.05 mg/kg IV/IO q 5 min. Until secretions dry or HR increases. The minimum dose is 0.1 mg.,max dose is 5 mg</td></tr></table>	Adult	<u>Rx: Symptomatic Bradycardia:</u> 1 mg IV push fast q 3-5 minutes; up to 3 mg total dose <u>Rx: Organophosphate or Carbamate Insecticide poisoning:</u> 2-5 mg IV/IO q 5 - 10 minutes until secretions dry or HR increases (No max dose in OPP)	Pedi	<u>Rx: Bradydysrhythmias:</u> 0.02 mg/kg IV, IO. The minimum dose is 0.1 mg. May be repeated in 5 minutes for a maximum total dose of 1.0 mg for a child and 2.0 mg for an adolescent <u>Rx: Organophosphate or Carbamate Insecticide poisoning:</u> 0.05 mg/kg IV/IO q 5 min. Until secretions dry or HR increases. The minimum dose is 0.1 mg.,max dose is 5 mg
Adult	<u>Rx: Symptomatic Bradycardia:</u> 1 mg IV push fast q 3-5 minutes; up to 3 mg total dose <u>Rx: Organophosphate or Carbamate Insecticide poisoning:</u> 2-5 mg IV/IO q 5 - 10 minutes until secretions dry or HR increases (No max dose in OPP)				
Pedi	<u>Rx: Bradydysrhythmias:</u> 0.02 mg/kg IV, IO. The minimum dose is 0.1 mg. May be repeated in 5 minutes for a maximum total dose of 1.0 mg for a child and 2.0 mg for an adolescent <u>Rx: Organophosphate or Carbamate Insecticide poisoning:</u> 0.05 mg/kg IV/IO q 5 min. Until secretions dry or HR increases. The minimum dose is 0.1 mg.,max dose is 5 mg				

NOTES

- Atropine causes pupillary dilation rendering the pupils nonreactive.
- Atropine should be cautiously used in the presence of AMI because excessive increases in rate may worsen ischemia or increase the zone of infarction.
- Atropine may not be effective in high-degree heart block.
- Large or repeated doses may decrease respiratory drive.

NAME	CEFAZOLIN (Ancef)				
CLASS	Cephalosporins, 1st Generation				
ACTION	Bactericidal; inhibits cell wall mucopeptide synthesis				
ONSET/DURATION					
INDICATIONS	- Patients with evidence of an open fracture (i.e. deformity with a break in the overlying/adjacent skin) - Large significant trauma with anticipated contamination				
CONTRAINDICATIONS	Hypersensitive to drug/class/components				
ADVERSE REACTIONS	- Neutropenia - Thrombocytopenia - Anaphylaxis - Stevens-Johnson syndrome - Nephrotoxicity - Diarrhea - Rash - Vomiting				
DOSE AND ROUTE	<table border="1"> <tr> <td>Adult</td><td> Rx: Open Fracture: weight > 120 kg administer 3000mg weight > 20 – 120 kg administer 2000 mg </td></tr> <tr> <td>Pedi</td><td> Rx: weight < 20 kg give 50 mg/kg </td></tr> </table>	Adult	Rx: Open Fracture: weight > 120 kg administer 3000mg weight > 20 – 120 kg administer 2000 mg	Pedi	Rx: weight < 20 kg give 50 mg/kg
Adult	Rx: Open Fracture: weight > 120 kg administer 3000mg weight > 20 – 120 kg administer 2000 mg				
Pedi	Rx: weight < 20 kg give 50 mg/kg				

NOTES

- Reconstitute with 0.9% NS and inject in 100 ml 0.9% NS – infuse over 10 minutes
- Using a 100c bag, and a 10 gtts set – infuse at 100gtt/min

NAME	CETACAINE				
CLASS	Topical Anesthetic				
ACTION	Cetacaine is used for rapid, brief, superficial anesthesia of the nasal pharynx and oral pharynx.				
ONSET/DURATION	<table> <tr> <td>Onset</td><td>5-10 Minutes</td></tr> <tr> <td>Duration</td><td>Transient</td></tr> </table>	Onset	5-10 Minutes	Duration	Transient
Onset	5-10 Minutes				
Duration	Transient				
INDICATIONS	To provide surface anesthesia of the upper airway mucosa to reduce resistance during tracheal intubation.				
CONTRAINDICATIONS	Hypersensitivity to Cetacaine				
ADVERSE REACTIONS	• Burning or stinging sensation • Irritation				
DOSE AND ROUTE	<table> <tr> <td>Adult</td><td>Aid in oral intubation: Topical spray—aerosol – spray 1 to 2 seconds into the back of the throat to allow numbing effect and reduce gag reflex.</td></tr> <tr> <td>Pedi</td><td>Aid in oral intubation: Topical spray—aerosol – spray 1 to 2 seconds into the back of the throat to allow numbing effect and reduce gag reflex.</td></tr> </table>	Adult	Aid in oral intubation: Topical spray—aerosol – spray 1 to 2 seconds into the back of the throat to allow numbing effect and reduce gag reflex.	Pedi	Aid in oral intubation: Topical spray—aerosol – spray 1 to 2 seconds into the back of the throat to allow numbing effect and reduce gag reflex.
Adult	Aid in oral intubation: Topical spray—aerosol – spray 1 to 2 seconds into the back of the throat to allow numbing effect and reduce gag reflex.				
Pedi	Aid in oral intubation: Topical spray—aerosol – spray 1 to 2 seconds into the back of the throat to allow numbing effect and reduce gag reflex.				

NOTES
• Reassure the patient and inform them about the procedure for intubation.

NAME	DEXAMETHASONE (DECADRON, HEXADROL)					
CLASS	Glucocorticoid/Corticosteroid					
ACTION	Glucocorticoids cause varied metabolic effects. They modify the body's immune responses to diverse stimuli. Inflammatory cytokines are inhibited. Suppresses acute and chronic inflammation; immunosuppressive effects					
ONSET/DURATION	<table><tr><td>Onset</td><td>Unknown</td></tr><tr><td>Duration</td><td>Half Life 1.8 to 3.5 hours </td></tr></table>		Onset	Unknown	Duration	Half Life 1.8 to 3.5 hours
Onset	Unknown					
Duration	Half Life 1.8 to 3.5 hours					
INDICATIONS	- Severe Allergic Reaction - Asthma - Croup - COPD					
CONTRAINDICATIONS	- Systemic sepsis - Bacterial infection - Hypersensitivity to product					
ADVERSE REACTIONS	- Hypertension - Sodium and water retention - Gastrointestinal bleeding - TB - None from single dose					
DOSE AND ROUTE	<table><tr><td>Adult</td><td>Rx: Asthma, Stridor, Croup, COPD, Severe Allergic Reaction (Anaphylaxis): 10 mg IV/IO, Nebulized or IM</td></tr><tr><td>Pedi</td><td>Rx: Asthma, Stridor, Croup, COPD, Severe Allergic Reaction (Anaphylaxis): 0.5mg/kg IV/IO, Ne or IM Max dose 10mg</td></tr></table>		Adult	Rx: Asthma, Stridor, Croup, COPD, Severe Allergic Reaction (Anaphylaxis): 10 mg IV/IO, Nebulized or IM	Pedi	Rx: Asthma, Stridor, Croup, COPD, Severe Allergic Reaction (Anaphylaxis): 0.5mg/kg IV/IO, Ne or IM Max dose 10mg
Adult	Rx: Asthma, Stridor, Croup, COPD, Severe Allergic Reaction (Anaphylaxis): 10 mg IV/IO, Nebulized or IM					
Pedi	Rx: Asthma, Stridor, Croup, COPD, Severe Allergic Reaction (Anaphylaxis): 0.5mg/kg IV/IO, Ne or IM Max dose 10mg					
NOTES	Unknown safety in pregnancy					

NAME	DEXTROSE 10%						
CLASS	Carbohydrate, Hypertonic Solution						
ACTION	10% dextrose solution (D10) is used in emergency care to treat hypoglycemia and in the management of coma of unknown origin.						
ONSET/DURATION	<table border="1"> <tr> <td>Onset</td><td>1-3 Minutes</td></tr> <tr> <td>Duration</td><td>Depends on the degree of hypoglycemia</td></tr> </table>	Onset	1-3 Minutes	Duration	Depends on the degree of hypoglycemia		
Onset	1-3 Minutes						
Duration	Depends on the degree of hypoglycemia						
INDICATIONS	• Hypoglycemia • Altered level of consciousness • Coma of unknown etiology • Seizure of unknown etiology						
CONTRAINDICATIONS	• Intracranial hemorrhage • Increased intracranial pressure • Known or suspected CVA in the absence of hypoglycemia						
ADVERSE REACTIONS	• Warmth, pain, burning from medication infusion • Hyperglycemia • Thrombophlebitis						
DOSE AND ROUTE <u>DILUTION RESULTS</u> • 250cc Bag NS – remove 50 cc- then inject 50cc D50 into the bag and mix.- you now have 250cc of D10 solution	<table border="1"> <tr> <td>Adult</td><td>Rx: Diabetic Emergencies (Hypoglycemia): administer up to 2 ml/kg of D10 solution. May repeat if needed</td></tr> <tr> <td>Pedi</td><td>Rx: Hypoglycemia, Seizures if Blood glucose level unknown: 2.5 ml/kg of D10 solution. May repeat if needed</td></tr> <tr> <td>Neonate:</td><td>Rx: Hypoglycemia, 2-3 ml/kg of D10 solution. May repeat if needed</td></tr> </table>	Adult	Rx: Diabetic Emergencies (Hypoglycemia): administer up to 2 ml/kg of D10 solution. May repeat if needed	Pedi	Rx: Hypoglycemia, Seizures if Blood glucose level unknown: 2.5 ml/kg of D10 solution. May repeat if needed	Neonate:	Rx: Hypoglycemia, 2-3 ml/kg of D10 solution. May repeat if needed
Adult	Rx: Diabetic Emergencies (Hypoglycemia): administer up to 2 ml/kg of D10 solution. May repeat if needed						
Pedi	Rx: Hypoglycemia, Seizures if Blood glucose level unknown: 2.5 ml/kg of D10 solution. May repeat if needed						
Neonate:	Rx: Hypoglycemia, 2-3 ml/kg of D10 solution. May repeat if needed						

NOTES

- Normal blood glucose range = 60-110 mg/dl
- Infiltration of IV sites during the administration of D10 may produce tissue necrosis at the site but is much safer than D50.
- A blood glucose of < 40mg/dl indicates hypoglycemia in an infant.
- Administer appropriate doses and recheck blood sugar.

NAME	DIAZEPAM (Valium)				
CLASS	Benzodiazepine				
ACTION	Diazepam acts on the limbic, thalamic, and hypothalamic regions of the CNS to potentiate the effects of inhibitory neurotransmitters, raising the seizure threshold in the motor cortex.				
ONSET/DURATION	<table border="1"> <tr> <td>Onset</td><td>(IV) 1-5 min (IM) 15-30 min (PR) 4 – 10 min</td></tr> <tr> <td>Duration</td><td>(IV) 15 min- 1 hour (IM) 15 min – 1 hour</td></tr> </table>	Onset	(IV) 1-5 min (IM) 15-30 min (PR) 4 – 10 min	Duration	(IV) 15 min- 1 hour (IM) 15 min – 1 hour
Onset	(IV) 1-5 min (IM) 15-30 min (PR) 4 – 10 min				
Duration	(IV) 15 min- 1 hour (IM) 15 min – 1 hour				
INDICATIONS	• Acute anxiety states • Acute alcohol withdrawal • Skeletal muscle relaxation • Seizure Activity • Premedication prior to counter shock or TCP				
CONTRAINDICATIONS	• Hypersensitivity to the drug • Substance abuse (use with caution) • Coma (unless the patient has seizures, severe muscle or myoclonus) • Shock • CNS depression as a result of head injury • Respiratory depression				
ADVERSE REACTIONS	• Hypotension • Reflex tachycardia (rare) • Respiratory depression • Ataxia • Psychomotor impairment • Confusion • Nausea				

DOSE AND ROUTE	<table border="1"> <tr> <td data-bbox="721 226 878 583">Adult</td><td data-bbox="878 226 1539 583"> Rx: Status Generalized Motor Seizures, Overdose toxic exposure seizures, Eclampsia seizures, Skeletal Muscle Relaxation, Pain, Premedication prior to Cardioversion or Pacing: 5-10 mg IV, over 2 minutes may be given IM, or IO May repeat q 10-15 minutes prn up to total dose of 30 mg guided by B/P and Respiratory effort Acute Anxiety or Acute Alcohol – 2 -10 mg IV, IM, IO. </td></tr> <tr> <td data-bbox="721 583 878 842">Pedi</td><td data-bbox="878 583 1539 842"> ● Rx: Status Seizures: Infants-5 yrs of age: 0.2-0.5 mg/kg slow IV, IO q 2-5 minutes to a maximum dose of 5 mg. ● Children > 5 yrs: 1 mg IV slow q 2-5 minutes to a maximum dose of 10 mg ● Rectal Valium: (PR) Double recommended IV dose </td></tr> </table>	Adult	Rx: Status Generalized Motor Seizures, Overdose toxic exposure seizures, Eclampsia seizures, Skeletal Muscle Relaxation, Pain, Premedication prior to Cardioversion or Pacing: 5-10 mg IV, over 2 minutes may be given IM, or IO May repeat q 10-15 minutes prn up to total dose of 30 mg guided by B/P and Respiratory effort Acute Anxiety or Acute Alcohol – 2 -10 mg IV, IM, IO.	Pedi	● Rx: Status Seizures: Infants-5 yrs of age: 0.2-0.5 mg/kg slow IV, IO q 2-5 minutes to a maximum dose of 5 mg. ● Children > 5 yrs: 1 mg IV slow q 2-5 minutes to a maximum dose of 10 mg ● Rectal Valium: (PR) Double recommended IV dose
Adult	Rx: Status Generalized Motor Seizures, Overdose toxic exposure seizures, Eclampsia seizures, Skeletal Muscle Relaxation, Pain, Premedication prior to Cardioversion or Pacing: 5-10 mg IV, over 2 minutes may be given IM, or IO May repeat q 10-15 minutes prn up to total dose of 30 mg guided by B/P and Respiratory effort Acute Anxiety or Acute Alcohol – 2 -10 mg IV, IM, IO.				
Pedi	● Rx: Status Seizures: Infants-5 yrs of age: 0.2-0.5 mg/kg slow IV, IO q 2-5 minutes to a maximum dose of 5 mg. ● Children > 5 yrs: 1 mg IV slow q 2-5 minutes to a maximum dose of 10 mg ● Rectal Valium: (PR) Double recommended IV dose				
NOTES	● Respiratory depression, although a rare occurrence, should be anticipated when administering valium. Prepare to assist with ventilations.				

NAME	DILTIAZEM (Cardizem)					
CLASS	Slow Calcium Channel Blocker					
ACTION	Calcium channel blocking agent that slows cardiac cell conduction, increases refractoriness in AV node, and causes coronary and peripheral vasodilation. The drug is used to control ventricular response rates in patients with atrial fibrillation or flutter, multifocal atrial tachycardia, and SVT.					
ONSET/DURATION	<table><tr><td>Onset</td><td>2-5 Minutes</td></tr><tr><td>Duration</td><td>1-3 Hours</td></tr></table>		Onset	2-5 Minutes	Duration	1-3 Hours
Onset	2-5 Minutes					
Duration	1-3 Hours					
INDICATIONS	• SVT • Accelerated Atrial Fibrillation • Atrial Flutter					
CONTRAINDICATIONS	• WPW • 2nd or 3rd-degree block • Hypotension (< 90 mm Hg) • Cardiogenic shock • Hypersensitivity to drug • Ventricular Tachycardia • Wide complex tachycardia of unknown origin • AMI					
ADVERSE REACTIONS	• Hypotension • Bradycardia • 2nd and 3rd-degree Block • Syncope • Ventricular dysrhythmias • Nausea and Vomiting • Dyspnea • Chest Pain					
DOSE AND ROUTE	<table><tr><td>Adult</td><td>Rx: Supraventricular Tachycardia, Atrial fib, Atrial flutter: Bolus initial dose 10mg IV SLOW over 2 - 5 minutes (up to 0.25mg/kg). May be repeated in 15 minutes at 0.35 mg/kg IV over 2-5 minutes</td></tr><tr><td>Pedi</td><td>Is <u>NOT</u> indicated for use in pediatric patients.</td></tr></table>		Adult	Rx: Supraventricular Tachycardia, Atrial fib, Atrial flutter: Bolus initial dose 10mg IV SLOW over 2 - 5 minutes (up to 0.25mg/kg). May be repeated in 15 minutes at 0.35 mg/kg IV over 2-5 minutes	Pedi	Is <u>NOT</u> indicated for use in pediatric patients.
Adult	Rx: Supraventricular Tachycardia, Atrial fib, Atrial flutter: Bolus initial dose 10mg IV SLOW over 2 - 5 minutes (up to 0.25mg/kg). May be repeated in 15 minutes at 0.35 mg/kg IV over 2-5 minutes					
Pedi	Is <u>NOT</u> indicated for use in pediatric patients.					

NOTES	• Hypotension is the most common side effect (Manage with fluid bolus if lungs are clear) • PVCs are common on the conversion of SVT to sinus rhythm

NAME	DIPHENHYDRAMINE (Benadryl)
CLASS	Antihistamine
ACTION	Antihistamines prevent the physiologic actions of histamine by blocking H1 and H2 receptor sites. Antihistamines are indicated for conditions in which histamine excess is present (e.g., acute urticarial), but also are used as adjunctive therapy (with epinephrine, for example) in the treatment of anaphylactic shock.
ONSET/DURATION	Onset: 10 – 20 min Duration: 6 – 12 hours
INDICATIONS	• Moderate to severe allergic reactions (administer epinephrine first if severe) • Anaphylaxis • Acute extrapyramidal reactions (EPS) • Nausea /Vomiting
CONTRAINDICATIONS	• Hypersensitivity • Narrow angle glaucoma (relative) • Newborns and nursing mothers
ADVERSE REACTIONS	• Dose-related drowsiness • Sedation • Disturbed coordination • Hypotension • Palpitations • Tachycardia, bradycardia
DOSE AND ROUTE	Adult: Rx: Allergic Reactions, Anaphylaxis: 25-50mg (IV) or (Deep IM), may be repeated x1. Nausea/Vomiting – 12.5 mg IV or IM Pediatric: Rx: Allergic Reactions, Anaphylaxis: 1mg/kg IV (max initial dose 25mg) 2mg/kg IM (max initial dose 50mg)
NOTES	When used in anaphylaxis, will be in conjunction with epinephrine and / or corticosteroids Not used in infants or in pregnancy

NAME		DOPAMINE (Intropin)												
CLASS		Sympathomimetic												
ACTION		At low doses dopamine acts on dopaminergic receptors causing renal, mesenteric, and cerebral vascular dilation. At moderate doses (“cardiac doses”), dopamine stimulates beta adrenergic receptors causing enhanced myocardial contractility, increased cardiac output, and a rise in blood pressure. At high doses (“vasopressor doses”), dopamine has an alpha-adrenergic effect, producing peripheral arterial and venous constriction.												
ONSET/DURATION		Onset: 2-4 min Duration: 10- 15 min												
INDICATIONS		• Hemodynamically significant hypotension in the absence of hypovolemia. • Bradycardia if Atropine has been ineffective.												
CONTRAINDICATIONS		• Tachydysrhythmias • Ventricular fibrillation												
ADVERSE REACTIONS		• Dose-related tachydysrhythmias • Hypertension • Increased myocardial oxygen demand (e.g., ischemia) • Renal failure												
DOSE AND ROUTE		Adult: Rx: Hypotension, Bradycardia: 5-20 µg/kg/min, titrate to effect. Use premix or mix 400 mg in 250 ml D5W (1600 µg/ml) use 60 gtt set. Pediatric Rx: Hypotension: 2-20 µg/kg/min. Use premix or mix 400 mg in 250 ml D5W (1600 µg/ml) use 60 gtt set. Extravasation causes tissue necrosis												
NOTES			Patient Weight in Kg											
			2.5	5	10	20	30	40	50	60	70	80	90	100
	µg /Kg/Min	5µg		1	2	4	6	8	9	11	13	15	17	19
		10µg	1	2	4	8	11	15	19	23	26	30	34	38
		15µg	1	3	6	11	17	23	28	34	39	45	51	56
		20µg	2	4	8	15	23	30	38	45	53	60	68	75
			Microdrops per minute (or ml/hr)											

NAME	Droperidol (Inapsine)				
CLASS	Neuroleptic (tranquilizer)				
ACTION	Antidopaminergic, Antiemetic, Anti-Psychotic				
ONSET/DURATION	<table> <tr> <td>Onset</td><td>IM/IV 3-10 minutes; peak in efficacy in 30 minutes.</td></tr> <tr> <td>Duration</td><td>2-4 hours</td></tr> </table>	Onset	IM/IV 3-10 minutes; peak in efficacy in 30 minutes.	Duration	2-4 hours
Onset	IM/IV 3-10 minutes; peak in efficacy in 30 minutes.				
Duration	2-4 hours				
INDICATIONS	- Severe agitation - Nausea/Vomiting - Migraine headache				
CONTRAINDICATIONS	Droperidol is contraindicated in patients with known or suspected QT prolongation including patients with congenital long QT syndrome				
ADVERSE REACTIONS					
DOSE AND ROUTE	<table> <tr> <td>Adult</td><td> - Agitation / Excited Delirium: 2.5 mg increments IV, IO or IM, titrate to the desired effect, maximum dose of 10 mg - Antiemetic: 0.625 – 1.25mg IV or I </td></tr> <tr> <td>Pedi</td><td>Pediatric dose (8-14 y/o only): 0.05 mg/kg slow IV push or IM</td></tr> </table>	Adult	- Agitation / Excited Delirium: 2.5 mg increments IV, IO or IM, titrate to the desired effect, maximum dose of 10 mg - Antiemetic: 0.625 – 1.25mg IV or I	Pedi	Pediatric dose (8-14 y/o only): 0.05 mg/kg slow IV push or IM
Adult	- Agitation / Excited Delirium: 2.5 mg increments IV, IO or IM, titrate to the desired effect, maximum dose of 10 mg - Antiemetic: 0.625 – 1.25mg IV or I				
Pedi	Pediatric dose (8-14 y/o only): 0.05 mg/kg slow IV push or IM				
NOTES	The onset of action of single intramuscular and intravenous doses is from three to ten minutes following administration, although the peak effect may not be apparent for up to thirty minutes				

NAME	EPINEPHRINE (Adrenalin)
CLASS	Sympathomimetic
ACTION	Epinephrine is an endogenous catecholamine that directly stimulates alpha, beta1 and beta2 adrenergic receptors in dose-related fashion. It is the initial drug of choice for treating bronchoconstriction and hypotension resulting from anaphylaxis as well as all forms of cardiac arrest.
ONSET/DURATION	Onset: (IM) 1-5 min (IV) 1-2 min Duration: 5-10 min
INDICATIONS	• Bronchial asthma • Upper Airway Edema • Hypotension • Non hypovolemic shock • Acute allergic reaction (anaphylaxis) • Cardiac arrest • Profound symptomatic bradycardia
CONTRAINDICATIONS	• Hypersensitivity (not an issue especially in emergencies- the dose should be lowered or given slowly in non-cardiac arrest patients with heart disease) • Hypovolemic shock (as with other catecholamines, correct hypovolemia prior to use) • Coronary insufficiency (use with caution)
ADVERSE REACTIONS	• Headache • Nausea • Restlessness/Agitation • Weakness • Dysrhythmia, including ventricular tachycardia and ventricular fibrillation • Hypertension • Precipitation of angina pectoris • Tachycardia
DOSE AND ROUTE	Adult: Rx: Cardiac Arrest: 1 mg IV/IO q 3-5 minutes. 1:10,000 IV or IO Rx: Upper Airway Edema/ Severe Asthma: 0.3-0.5 mg IM or Nebulized (0.3-0.5 ml 1:1000 in 3cc NS) Rx: Anaphylaxis with hypoperfusion: 0.3 – 0.5mg:1000 IM may repeat and/or 0.3-0.5 mg slow IV (3-5 ml 1:10,000) Rx: Bradycardia: 2 –10 µg/min IV infusion Rx: Shock - (PDP) Push dose presser administration – 1ml of 1:100,000 every minute until target HR and or B/P is achieved (see PDP under medication administration- located in guidelines) Pediatric: Rx: Cardiac Arrest: 0.01 mg/kg IV/IO q 3-5 minutes. 1:10,000 (0.1ml/kg) Rx: Allergic Reaction/ Severe Asthma/ upper airway edema: 0.01mg/kg 1:1000 IM or Nebulized – mix with 3cc NS Rx: Anaphylaxis with hypoperfusion 0.01mg/kg slow IV (1:10,000) Rx: Bradycardia/hypotension: 0.01 - .03 mg/kg IV, IO (1:10,000) Q 3 – 5 minutes Rx: Child hypoperfusion- 0.1mcg/kg/min drip up to 1 mcg/kg/min titrate to effect Neonatal: 0.01 -0.03 mg/kg (0.1 – 0.3 ml/kg) (IV, IO, UV) All 1:10,000. Rx: Croup/Stridor – 0.5 mg/kg of 1:1000 epi Nebulized, (Max of 5 ml dilute in 3 ml NS)
NOTES	• Do not administer Epinephrine in the buttocks on Children, use the Vastus Lateralis area of leg. Do not administer more than once in any given injection site. ET Administration is permissible use 2 – 2.5 X normal amount. • IV Epi can be diluted to 1:100,000

NAME	ETOMIDATE (Amidate)
CLASS	Sedative, hypnotic
ACTION	Exact action is unknown. May have GABA like effects, depresses brain stem reticular formation activity and produces hypnosis Creates an ultra-short acting sedative/hypnotic effect It decreases Intra Cranial Pressure (ICP) and Decreases cerebral blood flow thus resulting in decreased basal metabolic rate.
ONSET/DURATION	Onset: 30 – 60 seconds Duration: 10 – 15 minutes
INDICATIONS	• Sedative for Pharmacological Assisted Intubation (PAI) • Sedation / hypnotic
CONTRAINDICATIONS	• Patient < 8 year-old • Marked Hypotension • Pregnancy • Immunosuppression • Sepsis • Transplant patient
ADVERSE REACTIONS	• Apnea • Bradycardia • Hypotension • Dysrhythmias • N & V • Laryngospasm
DOSE AND ROUTE	Adult Rx: PAI: 0.3 mg/kg IV slowly Sedation: 0.2 mg/kg SLOWLY Pediatric: Sedation: Titrate up to 0.3mg/kg IV slowly until sedated
NOTES	Always use IV fluid flowing during administration to reduce pain at injection site.

NAME	FENTANYL CITRATE (SUBLIMAZE)
CLASS	Narcotic Analgesic
ACTION	Therapeutic values are analgesic and sedative Fentanyl is 50 – 100 times more potent than morphine. It has a rapid onset but its duration of action is shorter than that of meperidine or morphine.. Fentanyl has less emetic activity than other narcotics. The respiratory effect in slowing rate and alveolar ventilation may last longer than the analgesic effect.
ONSET/DURATION	Onset – IV or IO Immediate IN 2 – 10 Minutes Peak Effects: 3 -5 minutes (IV) Duration: 15 -30 minutes Half-Life: 6-8 hours
INDICATIONS	• Used for maintenance of analgesia, as an adjunct PAI • Pain /Analgesia
CONTRAINDICATIONS	• Severe hemorrhage • Shock • known hypersensitivity
ADVERSE REACTIONS	As seen with all narcotic analgesics: • May cause prolonged Q-T interval • Respiratory depression • Apnea • Chest wall muscle rigidity • Bradycardia • Hypotension
DOSE AND ROUTE	Adult: Rx: Analgesia / PAI: – 0.5 – 1mcg/kg – May repeat at 2 – 3 minutes to max of 2 mcg/kg May be given IV, IM, or IN Pediatric: Rx: Analgesia: 0.5 – 1mcg/kg IV or IM – may repeat one time Intra Nasal administration 2 mcg/kg – ½ in each nostril
NOTES	

NAME	FUROSEMIDE (Lasix)
CLASS	Loop diuretic
ACTION	Furosemide is a potent diuretic that inhibits the reabsorption of sodium and chloride in the proximal tubule and loop of Henle. Intravenous doses can also reduce cardiac preload by increasing venous capacitance.
ONSET/DURATION	Onset: (IV) diuretic effects within 15-20 min; vascular effects within 5 min Duration: 2 hours
INDICATIONS	• Pulmonary edema associated with CHF, hepatic or renal disease • Isolated closed head trauma with signs and symptoms of herniation • Severe Hyperkalemia
CONTRAINDICATIONS	• Hypersensitivity • Hypovolemia/dehydration • Known hypersensitivity to sulfonamides • Severe electrolyte depletion (hypokalemia) • Sepsis • Pneumonia
ADVERSE REACTIONS	• Hypotension • ECG changes associated with electrolyte disturbances • Dry mouth • Hypochloremia • Hypokalemia • Hyponatremia • Hypercalcemia • Hyperglycemia • Hearing loss can rarely occur after too rapid infusion of large doses especially in patients with renal impairment.
DOSE AND ROUTE	Adult: Rx: CHF with Pulmonary Edema, Hyperkalemia, Isolated Closed Head Trauma: 0.5-1.0 mg/kg Slow IV. Maximum dose: 2 mg/kg Pediatric: Rx: CHF with Pulmonary Edema 1 mg/kg IV, IO slowly
NOTES	Hypotension is a common side effect that often results when Lasix is given too rapidly. As the diuretic effect of Lasix usually does not begin for 15-20 minutes after the drug is given, the primary effect of this drug when given in the prehospital environment is to dilate the venous system and reduce preload in patients with bi-ventricular failure.

NAME	GEODON /ZIPRASIDONE
CLASS	Anti-psychotic and tranquilizing agent
ACTION	When introduced by intramuscular injection Geodon enters the body and acts as an antagonist at dopamine and serotonin receptors. It also moderately inhibits re-uptake of norepinephrine and serotonin. Geodon works as an anti-histamine.
ONSET/DURATION	Onset: IM 15 – 20 Minutes Duration: 2 hours for 10 mg and 4 hours for 20 mg.
INDICATIONS	• Agitated Delirium • Schizophrenia • Bipolar Disorders • Manic Disorders
CONTRAINDICATIONS	• Not to be administered to patients with Dementia-related psychosis • History of prolonged QT interval or medications that prolongs QT interval • Hypersensitivity to Geodon • Recent MI or uncontrolled CHF • Current use of antibiotics • Current use of antidepressants • Use of cancer medications
ADVERSE REACTIONS	• Dystonic reactions • Somnolence (sleepiness) • Dizziness • Headache • Orthostatic hypotension
DOSE AND ROUTE	Adult: Rx: Agitated Delirium, Schizophrenia, Bipolar disorder, Manic Disorders: 10 – 20 mg IM Pediatric: Rx: Not recommended
NOTES	

NAME	GLUCAGON
CLASS	Pancreatic hormone, insulin antagonist
ACTION	Glucagon is a protein secreted by the alpha cells of the pancreas. When released, it results in blood glucose elevation by increasing the breakdown of glycogen to glucose (glycogenolysis) and stimulating glucose synthesis (gluconeogenesis). The drug is only effective in treating hypoglycemia if liver glycogen is available, and may therefore be ineffective in chronic states of hypoglycemia, starvation, and adrenal insufficiency.
ONSET/DURATION	Onset: With in 1 minute Duration: 60 – 90 minutes
INDICATIONS	• Hypoglycemia when IV access is not obtainable or D₁₀ is contraindicated • Beta Blocker Overdose • Refractory Hypotension with Anaphylaxis
CONTRAINDICATIONS	• Hypersensitivity (allergy to proteins)
ADVERSE REACTIONS	• Tachycardia • Hypotension • Nausea, vomiting • Urticaria
DOSE AND ROUTE	Adult: Rx: Hypoglycemia: 0.5-1.0 mg (or unit) IM, IV (IN – use 2-3 mg) Rx: Beta Blocker OD: 3-10 mg IV (50-100 µg/kg), followed by drip: 1-5 mg/hour Rx: Anaphylaxis:: 1 – 2 mg IV Pediatric < 20 kg: Rx: Hypoglycemia 0.5 – 1.0 mg IV, IO, IM
NOTES	

NAME	HALOPERIDOL LACTATE (Haldol)
CLASS	Antipsychotic/Neuroleptic
ACTION	Haloperidol has pharmacologic properties similar to those of phenothiazines. The drug is thought to block dopamine (type 2) receptors in the brain, altering mood and behavior. In emergency care, haloperidol usually is given IM, but may also be given IV.
ONSET/DURATION	Onset: (IM) 30 – 60 minutes Duration: 12 – 24 hours
INDICATIONS	• Acute psychotic episodes • Emergency sedation of severely agitated or delirious patients • Excited Delirium
CONTRAINDICATIONS	• CNS depression • Coma • Hypersensitivity • Pregnancy • Severe liver or cardiac disease • Prolonged Q-T interval
ADVERSE REACTIONS	• Dose-related extrapyramidal reactions • Hypotension • Orthostatic hypotension • Nausea, vomiting • Allergic reactions • Blurred vision
DOSE AND ROUTE	Adult: Rx: Acute Psychotic episodes, Sedations, Severe Agitation/ Excited Delirium : 2-5 mg IM IV, or IN Pediatric: Not Recommended
NOTES	• Some patients may have prolonged reaction to Haldol • May significantly suppress CNS in patients with alcohol ingestion (monitor vital signs and prepare to intervene if respiratory depression occur)

NAME	HEPARIN SODIUM
CLASS	Anticoagulant
ACTION	Heparin inhibits the clotting cascade by activating specific plasma proteins. The drug is used in the prevention and treatment of all types of thrombosis and emboli, DIC, arterial occlusion and thrombophlebitis, and prophylactically to prevent clotting before and after surgery. Heparin is also considered part of the “thrombolytic package” administered to patients with acute myocardial infarction (along with aspirin and thrombolytic agents) and acute coronary syndromes including unstable angina and non-Q wave myocardial infarction.
ONSET/DURATION	Onset: (IV) Immediate (SQ) 20 – 60 min Duration: 4 –8 hours
INDICATIONS	• Acute myocardial infarction • Prophylaxis and treatment of thrombotic disorders (e.g., pulmonary emboli, DVT)
CONTRAINDICATIONS	• Hypersensitivity • Active bleeding • Recent intracranial, intraspinal, or eye surgery • Severe hypertension • Bleeding tendencies • Severe thrombocytopenia
ADVERSE REACTIONS	• Allergic reaction (chills, fever, back pain) • Thrombocytopenia • Hemorrhage • Bruising
DOSE AND ROUTE	Adult Follow your Specific EMS service protocol for Heparin Administration Authorization from Medical Control is mandatory. 5000 units followed by maintenance infusion of 100 units/hour.
NOTES	Protamine sulfate is a Heparin antagonist and 1 mg neutralizes approx. 100 IU heparin

NAME	HYDROXYZINE (Vistaril, Atarax, Apresoline)
CLASS	Antihistamine
ACTION	Hydroxyzine is a histamine-1 receptor antagonist that is used to treat allergy-induced pruritus, and is used for its antiemetic and sedative properties. It is effective for treatment of anxiety and tension associated with neuroses and alcohol withdrawal. Concomitant use with analgesics may potentiate the effects.
ONSET/DURATION	Onset: (IM) 15 – 30 min Duration: 4 – 6 hr
INDICATIONS	• Nausea and vomiting
CONTRAINDICATIONS	• Hypersensitivity • Prolonged Q-T • Pregnancy • Hyper sensitivity to Zertech
ADVERSE REACTIONS	• Dry mouth • Drowsiness • N & V • Tachycardia • Diarrhea
DOSE AND ROUTE	Adult: Rx: Nausea and vomiting: 25-100 mg deep IM Pediatric: Rx: Nausea and vomiting: 0.5 - 1.0 mg/kg deep IM
NOTES	Localized burning at injection site is common.

NAME	Ibuprofen / Motrin
CLASS	Non-opioid analgesic, anti-inflammatory, antipyretic, NSAID
ACTION	Ibuprofen works to inhibit cyclooxygenase which prevents prostaglandin formation. Ibuprofen also inhibits chemotaxis and neutrophil formation, alters lymphocyte activity, and decreases cytokine levels. May also exert some effects on the hypothalamus
ONSET/DURATION	Onset: 30-60 minutes Duration: 6-8 hours, with maximum effect at 2-4 hours
INDICATIONS	Mild to moderate pain; Fever over 38.0 Celsius taken orally or rectally for those under 2 years of age; Febrile seizure
CONTRAINDICATIONS	Hypersensitivity, history of renal or hepatic disease, actual or suspected hematologic/oncologic, bleeding disorders, history of ulcers, concern low platelet levels, $\leq$ six (6) months of age, has had ibuprofen during the previous 6 hours
ADVERSE REACTIONS	Possible GI irritation, bleeding, anaphylaxis, heartburn, dizziness, skin rashes, kidney, or liver problems, hypertension, and ringing in ears

DOSE AND ROUTE	<table border="1"> <tr> <td data-bbox="837 275 1040 680">Adult</td><td data-bbox="1040 275 1412 680"> Tablet Dosage (for patients able to swallow pills): - o 20 – 40 kg: 200mg by mouth - o 40 – 60 kg: 400mg by mouth - o 60 – 80 kg: 600mg by mouth >80 kg: 800mg by mouth </td></tr> <tr> <td data-bbox="837 680 1040 852">Pedi</td><td data-bbox="1040 680 1412 852"> 10 ml/kg/dose every 6-8 hours. Max liquid dose of 800 mg. Round to the nearest ML </td></tr> </table>	Adult	Tablet Dosage (for patients able to swallow pills): - o 20 – 40 kg: 200mg by mouth - o 40 – 60 kg: 400mg by mouth - o 60 – 80 kg: 600mg by mouth >80 kg: 800mg by mouth	Pedi	10 ml/kg/dose every 6-8 hours. Max liquid dose of 800 mg. Round to the nearest ML
Adult	Tablet Dosage (for patients able to swallow pills): - o 20 – 40 kg: 200mg by mouth - o 40 – 60 kg: 400mg by mouth - o 60 – 80 kg: 600mg by mouth >80 kg: 800mg by mouth				
Pedi	10 ml/kg/dose every 6-8 hours. Max liquid dose of 800 mg. Round to the nearest ML				
NOTES	Weight dosing is preferred, however, if not available, may use the table provided or dosing by age on product directions. Pay close attention to packaging and concentration as there is a wide variety available. Check concentration prior to administration, max dosing of 2,400mg or 40mg/kg/day or 4 total doses whichever occurs first				

NAME	IPRATROPIUM (Atrovent)
CLASS	Anticholinergic
ACTION	Ipratropium is an anticholinergic (parasympatholytic) bronchodilator that is chemically related to atropine.
ONSET/DURATION	Onset: Varies Duration: 4-6 hours
INDICATION	• Bronchial Asthma • Chronic Bronchitis • Emphysema
CONTRAINDICATIONS	• Hypersensitivity to the drug • Allergies Atropine
ADVERSE REACTIONS	• Palpitations • Anxiety • Rash • Nausea • Vomiting • Nervousness • Dizziness
DOSE AND ROUTE	Adult: Rx: Bronchial Asthma, Chronic Bronchitis, Emphysema: 0.5mg nebulized – max 3 doses Can be administered with a B_2 agonist in a nebulized treatment Pediatric: Rx: Bronchial Asthma, Chronic Bronchitis: 0.5mg nebulized – max 3 doses Can be administered with a B_2 agonist in a nebulized treatment
NOTES	

NAME	KETAMINE (Ketalar)
CLASS	Anesthetic Analgesic
ACTION	Causes disassociation. Ketamine acts on the limbic system and cortex to block afferent transmission of impulses associated with pain perception. It produces short-acting amnesia without muscular relaxation.
ONSET/DURATION	Onset: 30 seconds IV Duration: IV 5-10 minutes IM up to 30 minutes IM
INDICATIONS	• PAI (Pharmacological assisted intubation) • Pain • Agitated Delirium • Chemical Sedation of Violent Patient • Post resuscitation when Versed is contraindicated
CONTRAINDICATIONS	• Hypertensive Patients • Patients with increased intracranial pressure • Glaucoma • Hypersensitivity to drug
ADVERSE REACTIONS	• Increased salivation • Elevated blood pressure • Elevated heart rate • Altered mental status
DOSE AND ROUTE	Adult: Rx: Pain Control: 0.1 – 0.3 mg/kg IV over 1 minute – may repeat 1 time or 0.2 – 0.6 mg/kg IM/IN (no more than 1cc per nostril) Rx:: PAI, Post Resuscitation, Violent Patient: 0.5 - 1 mg/kg IV over 1 minute or 1-2 mg/kg IM/IN Pediatric > 2years old: Rx: Pain: 0.5 mg/kg IV over 1 minute may repeat one time max 25mg
NOTES	Consider Atropine in Children at 0.1mg/kg Consider lower doses for sedation (0.5mg/kg) if patient is under the influence of opiates. Use of capnography and pulse ox for all patient receiving Ketamine

	KETOROLAC TROMETHAMINE (Toradol)
CLASS	Non-steroidal anti-inflammatory
ACTION	Ketorolac Tromethamine is an anti-inflammatory drug that also exhibits peripherally acting non-narcotic analgesic activity by inhibiting prostaglandin synthesis.
ONSET/DURATION	Onset: Within 10 min Duration: 6-8 hours
INDICATION	Short-term management of moderate to severe pain
CONTRAINDICATIONS	• Hypersensitivity to the drug • Pregnancy • Pain associated with significant trauma/bleeding • Patients with allergies to aspirin or other nonsteroidal anti-inflammatory drugs • Bleeding disorders • Renal failure • Active peptic ulcer disease • Use of blood thinners: Coumadin, Plavix, etc.
ADVERSE REACTIONS	• Anaphylaxis from hypersensitivity • Edema • Sedation • Bleeding disorders • Rash • Nausea • Headache
DOSE AND ROUTE	Adult: Rx: Analgesia: 15-30 mg IV or 30-60 mg IM Pediatric: Typically Not Recommended
NOTES	Toradol (30mg) usually provides analgesia comparable to 12 mg Morphine or 100 mg Demerol.

NAME	LABETALOL (Normodyne, Trandate)
CLASS	Alpha and beta adrenergic blocker
ACTION	Labetalol is a competitive alpha, receptor blocker as well as a nonselective beta receptor blocker that is used for lowering blood pressure in hypertensive crisis.
ONSET/DURATION	Onset: Within 5 min Duration: 3-6 hours
INDICATION	Hypertensive emergencies: Systolic B/P of greater than 230 Diastolic B/P of greater than 120 (Goal is to drop B/P no less than 170 systolic / 90 diastolic)
CONTRAINDICATIONS	• Known or suspected CVA without medical control approval • Bronchial asthma (relative) • Uncompensated CHF • Second and third degree heart block • Bradycardia • Cardiogenic shock • Pulmonary edema
ADVERSE REACTIONS	• Headache • Dizziness • Dose related orthostatic hypotension • Fatigue • Vertigo • Ventricular dysrhythmias • Dyspnea • Allergic reaction • Facial flushing • Diaphoresis
DOSE AND ROUTE	Adult: Rx: Hypertensive Crisis: 10-20 mg IV over 1-2 minutes. May repeat or double dose q 10 minutes until a total dose of 150 mg OR start infusion at 2 mg/min. Drip: Start at 2 mg/min.
NOTES	Bronchodilator effects of Albuterol may be blunted by Labetalol With infusion: Usually do not want to drop BP by more than 10 mmHg over 2 minutes

NAME	LEVALBUTEROL (XOPENEX)
CLASS	Sympathetic agonist
ACTION	Levalbuterol is a selective B_2 -adrenergic agonist that causes relaxation of bronchial smooth muscle, thus decreasing airway resistance and increasing vital capacity. Levalbuterol is a chemical variant of albuterol with greater affinity for the B_2 -adrenergic receptors.
ONSET/DURATION	Onset: 5 -15 minutes Duration: 3 - 6 hours
INDICATIONS	• Asthma • Chronic bronchitis • Emphysema • Reversible obstructive airway disease
CONTRAINDICATIONS	• Known hypersensitivity to the drug
ADVERSE REACTIONS	• Tremors • Anxiety • Dizziness • Headache • Insomnia • Nausea • Palpitations • Tachycardia • Hypertension
DOSE AND ROUTE	Adult: Rx: Asthma or Allergic Reaction: 0.63mg in 3.0 mL normal saline Pediatric: Rx: Asthma or Allergic Reaction: < 12 year-of-age 0.31 mg nebulized / mix in 3.0 ml normal saline

	LIDOCAINE (Xylocaine)																				
CLASS	Antidysrhythmic (Class I-B) Local anesthetic																				
ACTION	Lidocaine decreases phase 4 diastolic depolarization (which decreases automaticity), and has been shown to be effective in suppressing premature ventricular complexes. In addition it is used to treat ventricular tachycardia. Lidocaine also raises the ventricular fibrillation threshold.																				
ONSET/DURATION	Onset: 30-90 sec Duration: 10-20 min																				
INDICATIONS	• Pain from Intraosseous Infusion Pressure • Used if Amiodarone is not available for: ◦ Ventricular tachycardia ◦ Ventricular fibrillation ◦ Wide-complex tachycardia of uncertain origin ◦ Significant ventricular ectopy in the setting of myocardial ischemia/infarction																				
CONTRAINDICATIONS	• Hypersensitivity • Second or third degree heart block • Relative Contraindication: Bradycardic rhythms with escape ectopy																				
ADVERSE REACTIONS	• Lightheadedness • Confusion • Blurred vision • Slurred speech • Hypotension • Bradycardia • Altered level of consciousness, irritability, muscle twitching, seizures with high doses																				
DOSE AND ROUTE	Adult: Rx: Cardiac Arrest VT/VF: 1-1.5 mg/kg IVP. (ET dose 2-4 mg/kg)May repeat with 0.5-0.75 mg/kg IVP q 5-10 minutes. Maximum: 3 mg/kg. If effective conversion start drip ASAP (2-4 mg/min) Rx: VT with Pulse: 1 – 1.5 mg/kg IVP; then 0.5 – 0.75 mg/kg q 5 – 10 minutes up to 3 mg/kg. Start Drip ASAP (2 – 4 mg/min) Rx: AICD firing, and/or Frequent PVC's with cardiac symptomology: 0.5 – 1.5 mg/kg IV. May repeat as above up to 3 mg/kg. Start Drip ASAP (2 – 4 mg/min) Rx: Intraosseous Infusion Pain control: 40mg and then wait at least 1- 2 minutes prior to flushing with 10cc NS. May use a subsequent dose of 10 – 20mg if pain returns or is persistent. Pediatric: Rx: VF/VT: 1 mg/kg IV, IO. Followed by drip of 20 – 50 µg/kg/min (See Length Based Tape, Pedi-Wheel, or EMS Field Guide for Pediatric Infusions of Lidocaine) Rx- IO Pain: 0.5mg/kg up to max of 40 mg <table><tr><th colspan="5">Drip: 1-4 mg/min. Use premix of mix 1 Gm in 250 ml D₅W & run at:</th></tr><tr><th>Lidocaine Drip (4 mg/ml)</th><th>1 mg</th><th>2 mg</th><th>3 mg</th><th>4 mg</th></tr><tr><th>Micro drops/min (ml/hr)</th><td>15gtts0</td><td>30gtts</td><td>45gtts</td><td>60gtts</td></tr><tr><th>If using Premix (8 mg/ml) run at</th><td>7gtts</td><td>15gtts</td><td>30gtts</td><td>45gtts</td></tr></table> Microdrops per minute (or ml/hr) Reduce maintenance infusion by 50% if patient is >70 YO, has liver disease, or is in CHF or Shock	Drip: 1-4 mg/min. Use premix of mix 1 Gm in 250 ml D ₅ W & run at:					Lidocaine Drip (4 mg/ml)	1 mg	2 mg	3 mg	4 mg	Micro drops/min (ml/hr)	15gtts0	30gtts	45gtts	60gtts	If using Premix (8 mg/ml) run at	7gtts	15gtts	30gtts	45gtts
Drip: 1-4 mg/min. Use premix of mix 1 Gm in 250 ml D ₅ W & run at:																					
Lidocaine Drip (4 mg/ml)	1 mg	2 mg	3 mg	4 mg																	
Micro drops/min (ml/hr)	15gtts0	30gtts	45gtts	60gtts																	
If using Premix (8 mg/ml) run at	7gtts	15gtts	30gtts	45gtts																	

	LORAZEPAM (Ativan)
CLASS	Antianxiety, Sedative
ACTION	Anxiolytic effect with skeletal muscle relaxation.
ONSET/DURATION	Onset: (IV) 1-5 minutes IM 15 – 30 minutes Duration: 12-24 hours
INDICATIONS	• Anxiety Disorder • Status Epilepticus • Skeletal muscle spasms • Sedation Prior to cardioversion or RSI • Post PAI sedation
CONTRAINDICATIONS	• Acute narrow angle Glaucoma • Acute Alcohol Intoxication • Shock
ADVERSE REACTIONS	• Nausea/Vomiting • Hypotension • Respiratory/CNS depression
DOSE AND ROUTE	Adult: Rx: Seizure, Anxiety Disorder: 2-4 mg slow IV (no faster than 2 mg/min) or IM May repeat every 15 – 20 minutes Pediatric: Rx: Seizure: 0.1 mg/kg slow IV or IO (Over 2-5 min) up to a maximum of 4 mg. May repeat at 0.5 mg/kg in 10-15 min.
NOTES	• Lorazepam can be administered rectally in the pediatric patient at double the recommended IV dose. • Attempt to use large secure veins as venous irritation is a common side effect

NAME	MAGNESIUM SULFATE
CLASS	Electrolyte, Anticonvulsant
ACTION	Magnesium sulfate reduces striated muscle contractions and blocks peripheral neuromuscular transmission by reducing acetylcholine release at the myoneural junction. In emergency care, magnesium sulfate is used in the management of seizures associated with toxemia of pregnancy. Other uses of magnesium sulfate include uterine relaxation (to inhibit contractions of premature labor), as a bronchodilator after beta agonist and anticholinergic agents have been used, replacement therapy for magnesium deficiency. Magnesium sulfate is gaining popularity as an initial treatment in the management of various dysrhythmias, particularly torsades de pointes, and dysrhythmias secondary to TCA overdose or digitalis toxicity.
ONSET/DURATION	Onset: (IV) Immediate Duration: (IV) 30 min
INDICATIONS	• Seizures of eclampsia (toxemia of pregnancy) • Torsades de pointes • Suspected hypomagnesemia • Refractory ventricular fibrillation • Asthma (must contact med control)
CONTRAINDICATIONS	• Heart block or myocardial damage • Hypermagnesia
ADVERSE REACTIONS	• Diaphoresis • Facial flushing • Hypotension • Depressed reflexes • Hypothermia • Reduced Heart rate • Respiratory depression • Diarrhea
DOSE AND ROUTE	Adult: Rx: Cardiac Arrest (Torsades, Hypomagnesemia, Refractory VF/VT): 1-2 gm IVP (5-10 gm may be needed) Rx: Torsades with a pulse: 1-2 gm IV over 5-60 min (mix in 50 ml of D₅W) Rx: Seizures secondary to Eclampsia: 1-4 gm IV slowly Rx: Asthma – 2 gm nebulized (after Med Control approval) Pediatric: Rx: Asthma, Cardiac Arrest (Torsades, Hypomagnesemia, Refractory VF/VT): 25—50 mg/kg IV, IO, Nebulized over 15-30 minutes. Maximum: 2 gm
NOTES	IV calcium chloride or calcium gluconate is an antagonist to magnesium if needed.

NAME	METHYLPREDNISOLONE (SOLU-MEDROL)
CLASS	Corticosteroid and anti-inflammatory
ACTION	Corticosteroids have multiple actions in the body. They have potent anti-inflammatory properties and inhibit many of the substances that cause inflammation.
ONSET/DURATION	Onset – Varies Duration- 1 – 5 weeks Half-life – 3.5 hours
INDICATIONS	• Moderate to Severe Anaphylaxis • Asthma
CONTRAINDICATIONS	• Known hypersensitivity • Systemic Fungal Infections
ADVERSE REACTIONS	• Fluid Retention • CHF • Hypertension • ABD distention • Vertigo • Head Ache • Nausea • Malaise • Hiccups
DOSE AND ROUTE	Adult: Rx: Anaphylaxis and Asthma: 125 mg IV or IO, IM Pediatric: Rx: Anaphylaxis and Asthma: 2 mg/kg IV, IO or IM
NOTES	

NAME	MIDAZOLAM HYDROCHLORIDE (Versed)
CLASS	Short-acting benzodiazepine
ACTION	Midazolam HCl is a water-soluble benzodiazepine that may be administered for sedation to relieve apprehension or impair memory prior to tracheal intubation or cardioversion. It may also be used in the setting of status seizure activity.
ONSET/DURATION	Onset: (IV) 1-3 min; dose dependent Duration: 2-6 hours; dose dependent
INDICATIONS	• Premedication for tracheal intubation or cardioversion • Seizures • Sedation for Pharmacological Assisted Intubation (PAI) • Traumatic Injury • Agitated Delirium
CONTRAINDICATIONS	• Hypersensitivity to midazolam • Glaucoma (relative) • Shock • Depressed vital signs • Hypotension
ADVERSE REACTIONS	• Respiratory depression • Hiccough • Cough • Over sedation • Nausea and vomiting • Fluctuations in vital signs • Hypotension
DOSE AND ROUTE	Adult: Rx: Sedation & Traumatic Injury: 0.1 mg/kg slow IV, or IN. Maximum single dose 5 mg – May repeat 1 time. Rx: Seizures & Agitated Delirium: 2.5 mg slow IV or IN. If unable to start IV, may give 5 mg IM, or IN Rx: PAI: 2.5mg Q 2 minutes up to max dose of 20 mg until sedated or B/P falls below 90 systolic Pediatric: Rx: Seizures: 0.1 mg/kg IV. Maximum 2.5 mg. If unable to start IV may give 0.2 mg/kg IM. Or 0.1mg/kg IN ½ of dose in each nostril (Maximum IM dose: 5 mg)
NOTES	• Sedative effects of Versed may be enhanced by patient use of barbiturates, alcohol or narcotics. • ECG monitor, blood pressure, and pulse oximetry should be monitored throughout administration of Versed and during transport. • Resuscitation equipment should be readily available.

NAME	MORPHINE SULFATE
CLASS	Opioid analgesic
ACTION	Morphine sulfate is a natural opium alkaloid that has a primary effect of analgesia. It also increases peripheral venous capacitance and decreases venous return ("chemical phlebotomy"). In addition, because morphine decreases both preload and afterload, it may decrease myocardial oxygen demand.
ONSET/DURATION	Onset: 1-2 min after administration Duration: 2-7 hours
INDICATIONS	• Chest pain associated with myocardial infarction. • Pulmonary edema, with or without associated pain. • Moderate to severe acute and chronic pain.
CONTRAINDICATIONS	• Hypersensitivity to narcotics • Hypovolemia • Hypotension • Head injury • Increased ICP • Severe respiratory depression, exacerbated COPD • Decreased LOC • Acute/severe bronchial asthma
ADVERSE REACTIONS	• Hypotension • Tachycardia • Bradycardia • Palpitations • Syncope • Respiratory Depression • Euphoria • Bronchospasm • Dry mouth • Allergic reaction
DOSE AND ROUTE	Adult: Rx: Analgesia, pulmonary edema: 2-5 mg IV, IM, SQ. May repeat q 5minutes up to 20 mg Pediatric: Rx: Analgesia, pulmonary edema 0.1-0.2 mg/kg IV, SQ, IO, IM
NOTES	• Narcan should be readily available • Anticipate respiratory depression and prepare to intervene • May worsen bradycardia or heart block in inferior M.I. (vagotonic effect) • Use with caution in patients under the influence of Alcohol or narcotics

NAME	NALOXONE (Narcan)
CLASS	Narcotic Antagonist
ACTION	Naloxone is a competitive narcotic antagonist that is used in the management of known or overdose caused by narcotics. Naloxone antagonizes all actions of morphine.
ONSET/DURATION	Onset: Within 2 min Duration: 30-60 min
INDICATIONS	For the complete or partial reversal of CNS and respiratory depression induced by opioids including but not limited to the following: • Decreased level of consciousness • Coma of unknown origin. • Clonidine
CONTRAINDICATIONS	• Hypersensitivity
ADVERSE REACTIONS	• Tachycardia • Hypertension • Dysrhythmias • Nausea and vomiting • Diaphoresis • Blurred vision • Withdrawal (opiate)
DOSE AND ROUTE	Adult: Rx: Narcotic OD, Clonidine, Coma, Decreased LOC: 0.4-2 mg IV, IM, SQ, SL, IN (or ET diluted) May repeat in 5 minute intervals up to 10 mg Pediatric: Rx: Narcotic OD, Coma, Decreased LOC: 0.1 mg/kg IV, IO, ET, IM, SQ, IN (see Broselow Tape®, Pedi-wheel®, or EMS field Guide)
NOTES	• Use with caution in narcotic dependent patients who may experience withdrawal syndrome (including neonates of narcotic-dependent mothers) • May precipitate narcotic withdrawal with hypertension, tachycardia, and violent behavior. • May not reverse hypotension • May precipitate seizures or violent behavior • Use of capnography is a prudent step • Vomiting may occur if administered too rapidly

NAME	NITROGLYCERIN (Nitrostat and others)
CLASS	Vasodilator
ACTION	Nitrates and nitrites dilate arterioles and veins in the periphery (and coronary arteries in high doses). The resultant reduction in preload, and to a lesser extent in afterload, decreases the workload of the heart and lowers myocardial oxygen demand.
ONSET/DURATION	Onset: 1-3 min Duration: 30-60 min
INDICATIONS	• Ischemic chest pain • Pulmonary hypertension • CHF • Hypertensive emergencies
CONTRAINDICATIONS	• Hypersensitivity • Hypotension • Head injury • Cerebral hemorrhage (CVA) • Use with caution in setting of inferior AMI.
ADVERSE REACTIONS	• Transient headache • Reflex tachycardia • Hypotension • Nausea and vomiting • Postural syncope • Diaphoresis
DOSE AND ROUTE	Adult: Rx: Angina, Hypertensive Crisis, Pulmonary Edema: 0.4 mg SL spray or tablet. May repeat in 3-5 minutes prn pain without hypotension Rx: Unstable Angina and CHF (IV Drip) 5 – 200 mcg/min titrated to blood pressure and symptoms Pediatric: Not recommended
NOTES	• Assure IV access prior to administration of NTG in setting of inferior AMI • Have IV fluid hanging in anticipation of syncope and/or hypotension. • NTG can also be administered in CHF patients with altered LOC when presenting with hypertension and pulmonary edema.

NAME	ONDANSETRON HYDROCHLORIDE - ZOFRAN
CLASS	Antiemetic
ACTION	Selectively antagonizes serotonin 5-HT ₃ receptors.
ONSET/DURATION	Onset: Rapid. Peaks in 15-30 minutes Duration: 4-8 hours
INDICATIONS	• Nausea • Vomiting
CONTRAINDICATIONS	• Hypersensitivity • 1st Trimester Pregnancy
ADVERSE REACTIONS	• Bronchospasm • Anaphylaxis • Transient Blindness • Dizziness • Abdominal Pain • Fatigue • Drowsiness • ECG changes including prolonged Q-T interval
DOSE AND ROUTE	Adult: Rx: Nausea: 4-8mg IV Slow (over 1 minute) or IM. May repeat once up to max dose of 8mg. SL Tablet 4-8mg Pediatric: Rx: Nausea: 0.15mg/kg up to 4mg per dose IV Slow (over 1 minute). May repeat once up to max of 8mg
NOTES	• Side effects are rare • May be used for patients with known head trauma

NAME	OXYMETAZOLINE HYDROCHLORIDE (Afrin)
CLASS	Nasal decongestant; sympathomimetic-alpha agonist.
ACTION	It acts directly on alpha receptors of the sympathetic nervous system to constrict smaller arterioles in nasal passages and prolong decongestant effect. It has no effect on beta receptors.
ONSET/DURATION	Onset: 5-10 minutes Duration: 6-10 hours
INDICATION	Vasoconstriction of surface vessels in nasal passage to decrease bleeding and during nasal intubation.
CONTRAINDICATIONS	• Children under 6 years old • Pregnancy
ADVERSE REACTIONS	Burning, stinging, dryness of nasal mucosa
DOSE AND ROUTE	Intranasal Spray Adult: Rx: Nasal Intubation/Bleeding: 0.05% Solution; 2-3 sprays of drops in each nostril Pediatric: Rx: Nasal Intubation : > 6 years old: 0.025% solution; 2-3 sprays in each nostril
NOTES	• May reduce the incidence of epistaxis with nasal tracheal intubation. • Reassure the patient and inform them about the procedure for nasal intubation.

NAME	OXYTOCIN (Pitocin)
CLASS	Pituitary hormone
ACTION	It stimulates uterine smooth muscle contractions, and helps expedite the normal contractions of a spontaneous labor. The drug is administered in the prehospital setting to control postpartum bleeding.
ONSET/DURATION	Onset: (IV) Immediate (IM) within 3-5 min Duration: (IV) 20 minutes after the infusion is stopped (IM) 30-60 min
INDICATIONS	• Postpartum hemorrhage after infant and placental delivery
CONTRAINDICATIONS	• Hypertonic or hyperactive uterus • Presence of a second fetus • Fetal distress
ADVERSE REACTIONS	• Hypotension • Tachycardia • Hypertension • Dysrhythmias • Angina pectoris • Anxiety • Seizure • Nausea and vomiting • Allergic reaction • Uterine rupture (from excessive administration)
DOSE AND ROUTE	Adult: Rx: Postpartum Hemorrhage after infant and placental delivery: IV, mix 10 units in 1000 ml of NS or LR and infuse at 20 – 30 drops per minute via microdrop tubing, titrated to severity of bleeding and uterine response or 3 – 10 units IM following delivery of placenta
NOTES	Fundal massage along with allowing neonate to nurse can also aid in controlling postpartum hemorrhage

NAME	PROCAINAMIDE (Pronestyl)								
CLASS	Antidysrhythmic (Class 1-A)								
ACTION	Procainamide suppresses phase 4 depolarization in normal ventricular muscle and Perkinge fibers, reducing the automaticity of ectopic pacemakers. It also suppresses reentry dysrhythmias by slowing intraventricular conduction. Procainamide may be effective in treating PVC's and recurrent ventricular tachycardia that cannot be controlled with lidocaine.								
ONSET/DURATION	Onset: 10-30 min Duration: 3-6 hours								
INDICATIONS	• Suppressing PVCs refractory to lidocaine or Amioderone • Suppressing VT (with a pulse) refractory to Amioderone or lidocaine • Suppressing VF refractory to Amioderone or lidocaine • PSVTs with wide complex tachycardia of unknown origin								
CONTRAINDICATIONS	• Second and third-degree AV block • Digitalis toxicity • Torsades de pointes • Complete heart block • Tricyclic antidepressant toxicity								
ADVERSE REACTIONS	• Hypotension • Bradycardia • Reflex tachycardia • AV block • Widened QRS • Prolonged P-R or Q-T • Confusion • Seizure								
DOSE AND ROUTE	<u>Adult:</u> Rx: Cardiac Arrest VF/VT: 100 mg IVP q 5 minutes, or: 20 mg/min IV drip (maximum dose 17 mg/kg) Start Drip ASAP if successful conversion Rx: A-fib, VT; PSVT with WPW: 20 mg/min IV until dysrhythmia is converted, hypotension or QRS/QT widening develops, or 17 mg/kg has been given <u>Pediatric:</u> Not recommended in prehospital setting								
NOTES	Procainamide Drip: 1-4 mg/min: Mix 1 gm in 250 of D₅W & 60 gtt set run at: <table><tr><td>15gtt/min</td><td>30 gtt/min</td><td>45 gtt/min</td><td>60gtt/min</td></tr><tr><td>1 mg /min</td><td>2 mg/min</td><td>3 mg/ min</td><td>4 mg / min</td></tr></table>	15gtt/min	30 gtt/min	45 gtt/min	60gtt/min	1 mg /min	2 mg/min	3 mg/ min	4 mg / min
15gtt/min	30 gtt/min	45 gtt/min	60gtt/min						
1 mg /min	2 mg/min	3 mg/ min	4 mg / min						

NAME	PROMETHAZINE (Phenergan)
CLASS	Phenothiazine, Antihistamine
ACTION	Promethazine is an H ₁ receptor antagonist that blocks the actions of histamine by competitive antagonism at the H ₁ level. In addition to antihistaminic effects, promethazine also possesses sedative, antiemetic, and considerable anticholinergic activity.
ONSET/DURATION	Onset: (IM) (rapid) Duration: 4-6 hours
INDICATIONS	• Nausea and vomiting • Motion sickness • To potentiate the effects of analgesics • Allergic reactions
CONTRAINDICATIONS	• Hypersensitivity • Comatose states • CNS depression from alcohol, barbiturates, or narcotics. • Vomiting of unknown etiology in children. • Acutely ill dehydrated children.
ADVERSE REACTIONS	• Sedation • Dizziness • Allergic reactions • Dysrhythmias • Hyperexcitability • Dystonias • Burning at administration site • Use with caution in head injured patient's
DOSE AND ROUTE	Adult: Rx: Nausea/Vomiting, Potentiate effects of analgesics: (25 mg or deep IM) (This drug has not been FDA approved for IV administration and is not recommended by the drug manufacturer to be administered IV) Pediatric: Because of potential adverse reactions consider consulting medical control prior to administration in children (if administered: 0.5mg/kg IM)
NOTES	• Generally considered safe for use in pregnancy and during labor • Anticipate sedative effect and monitor airway and respiratory status

NAME	RACEMIC EPINEPHRINE
CLASS	Sympathomimetic
ACTION	Stimulate alpha and beta adrenergic receptors, reducing mucosal secretions and relaxing bronchial smooth muscles.
ONSET/DURATION	Onset: Immediate 1-5 min Duration: 1 -3 hours
INDICATIONS	• Bronchospasm (upper Airway) • Croup
CONTRAINDICATIONS	• Hypersensitivity to drug • Hypersensitivity to sulfites • Cardiac disease • Hypertension • Diabetes • Glaucoma angle-closure • BPH
ADVERSE REACTIONS	• Arrhythmias • Bronchospasm- paradoxical • Dizziness • Headache • Nervousness • Tremor • Insomnia • Nausea • Tachycardia
DOSE AND ROUTE	Adult: Rx: Bronchospasm/Croup- 0.5 mL NEB q3-4h of 2.25% solution Pediatric: Rx: Bronchospasm/Croup - < 4 years old – 0.25 mL of 2.25% Solution NEB q2-4h; Max 0.5 mL/dose q 1-2h Bronchospasm- > 4 Year old- 0.5 mL of 2.25% solution NEB Q 3- 4h
NOTES	

NAME	ROCURONIUM BROMIDE (Zemuron)
CLASS	Neuromuscular blocker
ACTION	Antagonizes motor endplate acetylcholine receptors (non-depolarizing neuromuscular blocker)
ONSET/DURATION	Onset: < 2 minutes Duration: 25% recovery in 31 minutes (short – intermediate-acting)
INDICATIONS	PAI (Pharmacological Assisted Intubation)
CONTRAINDICATIONS	Allergy or hypersensitivity to drug or drug class
ADVERSE REACTIONS	- Prolonged paralysis - Respiratory depression - Apnea - Anaphylactic reaction - Bronchospasm - Arrhythmias
DOSE AND ROUTE	Adult Rx: PAI: 1 mg/kg IV may not repeat
NOTES	

NAME	SODIUM BICARBONATE 8.4%
CLASS	Buffer, Alkalinizing agent, Electrolyte supplement
ACTION	Sodium bicarbonate reacts with hydrogen ions (H ⁺) to form water and carbon dioxide and thereby can act to buffer metabolic acidosis.
ONSET/DURATION	Onset: 2-10 min Duration: 30-60 min
INDICATIONS	• Known pre-existing bicarbonate responsive acidosis • Intubated patient with continued long arrest interval. PEA • Upon return of spontaneous circulation after long arrest interval • Tricyclic antidepressant overdose • Alkalinization for treatment of specific intoxications • Management of metabolic acidosis • DKA
CONTRAINDICATIONS	• In patients with chloride loss from vomiting and GI suction • Metabolic and respiratory alkalosis • Severe pulmonary edema • Abdominal pain of unknown origin • Hypocalcemia • Hypokalemia • Hypernatremia • When administration of sodium could be detrimental.
ADVERSE REACTIONS	• Metabolic alkalosis • Hypoxia • Rise in intracellular PCO₂ and increased tissue acidosis • Electrolyte imbalance (Hypernatremia) • Seizures • Tissue sloughing at injection site
DOSE AND ROUTE	Adult: Rx: Prolonged Cardiac Arrest with good ventilation: 1 mEq/kg IV (1 ml/kg) followed by 0.5 mEq/kg q 10 minutes Rx: Hyperkalemia, OD from Tricyclics, ASA, Phenobarbital, Cocaine, Benadryl: 1 mEq/kg IV Pediatric: Rx: Prolonged Cardiac Arrest with good ventilation: (1 mEq/kg) infuse slowly through good vein and only if ventilations are adequate (See Broselow Tape, Pedi wheel, or EMS Field Guide) Use 4.2% solution in neonates
NOTES	• Must flush IV lines before and after administration. • Must ventilate patient after administration. • Do not administer down ET. • When possible, arterial blood gas analysis should guide bicarbonate administration.

NAME	SUCCINYLCHOLINE (Anectine)
CLASS	Neuromuscular blocker (depolarizing)
ACTION	Succinylcholine has the quickest onset and briefest duration of action of all neuromuscular blocking drugs, making it a drug of choice for such procedures as endotracheal intubation, electroconvulsive shock therapy, and terminating laryngospasm.
ONSET/DURATION	Onset: Less than 1 min Duration: 5-10 min after single IV dose
INDICATIONS	• PAI (Pharmacological Assisted Intubation) • Terminating laryngospasm • Muscle relaxation
CONTRAINDICATIONS	• Acute injuries • Hypersensitivity • Skeletal muscle myopathies • Inability to control airway and or support ventilation with oxygen and positive pressure. • history of malignant hyperthermia • Acute rhabdomyolysis • Burns > 8 hours • Massive crush injury
ADVERSE REACTIONS	• Hypotension • Respiratory depression/apnea • Bradycardias • Dysrhythmias • Initial muscle fasciculation • Excessive salivation • Malignant hyperthermia • Allergic reaction
DOSE AND ROUTE	Adult: Rx: PAI: 2 mg/kg IV (onset 1 minute, recovery 4-6 minutes) IM dose: 3-4 mg/kg: onset 2-3 minutes with Max dose of 150 mg Pediatric: (Age 9 to puberty) Rx: PAI: 2 mg/kg IV -
NOTES	• All patients undergoing PAI should be appropriately sedated prior to receiving a paralytic agent as paralytic agents do not alter the patients LOC, hearing, memory, or feeling. • Neuromuscular agents produce respiratory paralysis: thus intubation, alternative airway adjuncts and resuscitative equipment should be readily available prior to administration.

NAME	TETRACAINE (Pontacaine)
CLASS	Topical ophthalmic anesthetic
ACTION	Tetracaine is used for rapid, brief, superficial anesthesia. The agent inhibits conduction of nerve impulses from sensory nerves.
ONSET/DURATION	Onset: Within 30 seconds Duration: 10-15 min
INDICATIONS	• Short-term relief from eye pain or irritation • Patient comfort before eye irrigation
CONTRAINDICATIONS	• Hypersensitivity to Tetracaine • Open injury to the eye
ADVERSE REACTIONS	• Burning or stinging sensation • Irritation
DOSE AND ROUTE	Adult: Rx: Eye pain: 1-2 gtt in affected eye Pediatric: Same as adult
NOTES	

NAME	THIAMINE (Betaxin)
CLASS	Vitamin (B1)
ACTION	Thiamine combines with adenosine triphosphate (ATP) to form thiamine pyrophosphate coenzyme, a necessary component for carbohydrate metabolism. Most vitamins required by the body are obtained through the diet. However, certain states such as alcoholism, malnourishment, and chemotherapy may affect the intake, absorption, and utilization of thiamine. The brain is extremely sensitive to thiamine deficiency.
ONSET/DURATION	Onset: Rapid Duration: Depends on degree of deficiency
INDICATIONS	• Coma of unknown origin (prior to or along with administration of D10 or naloxone) • Delirium tremens • Wernicke's encephalopathy
CONTRAINDICATIONS	None significant
ADVERSE REACTIONS	• Hypotension (from rapid injection) • Anxiety • Diaphoresis • Nausea/vomiting • Allergic reaction
DOSE AND ROUTE	Adult: Rx: 100 mg slow IV or deep IM Pediatric: Not recommended in the prehospital setting
NOTES	Anaphylactic reactions are possible For Coma of unknown Origin D10, Narcan, and Thiamine may be administered in succession if needed.

NAME	TRANEXAMIC ACID (TXA)
CLASS	Antifibrinolytic
ACTION	Competitively binds with Plasminogen to prevent clot breakdown thus preventing or stabilizing coagulopathy
ONSET/DURATION	Onset : Almost immediate when administered IV Duration : 2 – 4 hours
INDICATIONS	All three of below indications must be met: • Blunt or Penetrating trauma suggesting potential major hemorrhage (examples: multiple long bone fractures, flail chest, major abdominal injury, pelvic fracture, amputation) • Objective signs of hemorrhagic shock associated with trauma (B/P < 90 or HR > 115) • Known time of injury < 3 hours
CONTRAINDICATIONS	• Age less than 16 • Renal failure • Allergy to TXA • History of Thromboembolism • Known aneurismal Sub arachnoid hemorrhage • Injury greater than 3 hours old
ADVERSE REACTIONS	• Possible risk of thrombosis
DOSE AND ROUTE	Adult: Rx: Hemorrhage: 1 gram mixed in 100cc bag of NS and infuse IV or IO over 10 minutes. If possible – follow initial bolus with maintenance drip by reconstituting 1 gram in 250cc NS and infuse over the next 8 hours (31ml/hr.) Pediatric: Not indicated
NOTES	Hypotension may ensue if administered too quickly. Procedure: Reconstitute 1 (one) gram in 100ml NS and infuse over 10 minute. If possible, follow initial bolus with maintenance drip by reconstituting 1 gram in 250 ml NS and infuse over the next 8 hours (31 ml/hr). Should not be given in same line as blood products Administration of TXA should not alter any other treatments (fluids, medications, etc>)

NAME	VECURONIUM (Norcuron)
CLASS	Neuromuscular-blocking agent (non-depolarizing)
ACTION	Antagonizes motor endplate acetylcholine receptors (non-depolarizing neuromuscular blocker) resulting in paralysis
ONSET/DURATION	Onset: 75 – 90 seconds Duration: 25 – 40 min
INDICATIONS	PAI (Pharmacological Assisted Intubation) when Rocuronium is not available
CONTRAINDICATIONS	Hypersensitivity
ADVERSE REACTIONS	Prolonged paralysis
DOSE AND ROUTE	Adult: PAI Rx: 0.1 mg/kg IV Pediatric: (greater than 9 or puberty) PAI Rx: 0.08 – 0.1 mg/kg IV
NOTES	Should not be administered unless persons skilled in endotracheal intubation are present. Endotracheal intubation equipment must be available. Oxygen equipment and emergency resuscitative drugs must be available. Paralysis occurs after 1 minute and lasts for approximately 30 minutes.

NAME	VERAPAMIL (Isoptin)
CLASS	Calcium channel blocker (Class IV antiarrhythmic)
ACTION	Verapamil is used as an antidysrhythmic, antianginal, and antihypertensive agent. It works by inhibiting the movement of calcium ions across cell membranes. Verapamil decreases atrial automaticity, reduces AV conduction velocity, and prolongs the AV nodal refractory period. In addition, verapamil depresses myocardial contractility, reduces vascular smooth muscle tone, and dilates coronary arteries and arterioles in normal and ischemic tissues.
ONSET/DURATION	Onset: 2-5 min Duration: 30-60 min (up to 4 hr is possible)
INDICATIONS	• Only consider verapamil if Cardizem is unavailable • SVT • Atrial flutter with a rapid ventricular response • Atrial fibrillation with a rapid ventricular response
CONTRAINDICATIONS	• Hypersensitivity • Second-or-third degree heart block • Sinus bradycardia • Hypotension • Cardiogenic shock • Severe CHF • WPW with atrial fibrillation or flutter • Patients receiving intravenous beta blockers • Wide complex tachycardias (ventricular tachycardia can deteriorate into ventricular fibrillation when calcium channel blockers are given)
ADVERSE REACTIONS	• Dizziness • Headache • Nausea and vomiting • Hypotension • Bradycardia • Complete AV block
DOSE AND ROUTE	Adult: Rx: SVT, Accelerated A-Fib, A-Flutter: 2.5-5.0 mg IV SLOW over 3 minutes. May repeat 5 mg in 15 minutes. Maximum total dose: 20 mg. Pediatric: Not recommended in prehospital setting
NOTES	• Anticipate hypotension after administration. • Anticipate bradycardia. • Have resuscitation equipment readily available. • Some physicians recommend slow IV administration of 500 mg Calcium Chloride before Verapamil to minimize hypotension and bradycardia.

The Guideline Section
is included in the NW Protocol
to Function as a supplement to the protocols.

Guidelines are not meant to serve as protocols and
are to be used only when specifically indicated in the
Protocol Section of this manual

Guidelines will assist in performing specific
procedures. They are not meant to be restrictive.
Other techniques in performing a procedure
may be acceptable

Right Sided And Posterior 12 Lead Ecg

- The 12-lead ECG electrode placement is essential for paramedics and EMTs in prehospital settings as incorrect placement can lead to a false diagnosis of infarction or negatively change the EKG.
- Aside from a 12-lead ECG placement, there's something known as a 15-lead placement which includes placing leads V4-V6 on the posterior side of the patient below their left scapula (see below).
- The following are a few guidelines that are very helpful to reduce artifacts while performing ECGs.
- Place the patient in a supine position if the patient will tolerate it.
- Place the patient's arms down by their side to relax their shoulders.
- The patient's legs should be uncrossed.
- Electrical devices such as mobile phones should be away from the patient as these devices may interfere with the machine.
- Dry the skin if it is diaphoretic or moist.
- Shave any hair that can interfere with electrode placement.
- Electrode gel should be moist.
- Electrodes should not be placed over bones and over areas where there is a lot of muscle movement.
- Sometimes an abrasive material such as a washcloth may need to be used to remove dead skin cells.





Indications

- Suspected Right Ventricle Infarct
- Suspected Posterior Infarct
- Patients who have clinical findings highly suggestive of acute coronary ischemia but have ECG findings that are either normal or non-diagnostic.
- Patients with inferior lead changes or the reciprocal changes of ST-segment depression for a posterior infarction.
- ST-segment elevation suggestive of an inferior wall MI (II, III, AVF)
- Isolated ST-segment elevation in V1 or ST-segment elevation greater in V1 than in V2
Borderline ST-segment elevation in V5 and V6 or in V1 to V3
- ST-segment depression or suspicious isoelectric ST segments in V1 to V3.
- Because ST-segment elevation in V4R to V6R may resolve within 12 to 18 hours, the benefit of a 15- or 18-lead ECG is primarily soon after the patient's presentation. In addition, the presence of anterior wall infarction obscures the changes in the right precordial leads.
- 15- or 18-lead ECGs should be used soon after a presentation in patients with chest pain suggestive of acute coronary ischemia that does not have clear evidence of anterior ischemia or infarction.

Right Sided And Posterior 12 Lead Ecg

Procedure

1. Perform initial 12 Lead EKG procedures
2. Right Sided (Right view) a. Using new electrodes; position as follows and relocate leads V1, V2, and V3 as follows V4R: right midclavicular line, fifth intercostal space (use V3 lead)
3. Posterior (Posterior view) a. Using new electrodes; position as follows and relocate leads V4, V5, and V6 as follows V7: left posterior axillary line, straight line from V6 (use V4 lead) V8: left mid-scapular line, straight line from V7 (use V5 lead) V9: left paraspinal line, straight line from V8 (use V6 lead).
4. Label modified leads appropriately on all printouts
5. Notify receiving hospital of modifications when transmitted wirelessly



Right Sided And Posterior 12 Lead Ecg

CERTIFICATION REQUIREMENTS

EMT (apply)

ADVANCED EMT (apply)

PARAMEDIC (apply and interpret)

NOTES

- A right-sided 12-lead ECG placement allows you to detect a right-sided infarct. At a minimum, lead V4 should be placed on the 5th intercostal, mid-clavicular (exact opposite of the regular left side placement) if an inferior infarct was originally seen in leads II, III, and AVF.
- A Posterior view should consist of a minimum of V8 and V9.
- Electrode Misplacement
 - Up to 50 percent of cases have V1 and V2 in too high of a location which can mimic an anterior MI and cause T wave inversion.
 - Up to 33 percent of the cases have the precordial electrodes (V1-V6) lower or laterally misplaced which also leads to misdiagnosis.

APGAR Scoring System

Apgar Sign	2	1	0
Appearance (skin color)	Normal color all over (hands and feet are pink)	Normal color (but hands and feet are bluish)	Bluish-gray or pale all over
Pulse (heart rate)	Normal (above 100 beats per minute)	Below 100 beats per minute	Absent (no pulse)
Grimace ("reflex irritability")	Pulls away, sneezes, coughs, or cries with stimulation	Facial movement only (grimace) with stimulation	Absent (no response to stimulation)
Activity (muscle tone)	Active, spontaneous movement	Arms and legs flexed with little movement	No movement, "floppy" tone
Respiration (breathing rate and effort)	Normal rate and effort, good cry	Slow or irregular breathing, weak cry	Absent (no breathing)

What Is the Apgar Score?

The Apgar score is a test given to [newborns](#) soon after birth. This test checks a baby's heart rate, muscle tone, and other signs to see if extra medical care or emergency care is needed.

Babies usually get the test twice: 1 minute after birth, and again 5 minutes after they're born. If there are concerns, a baby may get the test again.

What Does It Check?

The Apgar score measures five things to check a baby's health. Each is scored on a scale of 0 to 2, with 2 being the best score:

1. **Appearance** (skin color)
2. **Pulse** (heart rate)
3. **Grimace** response (reflexes)
4. **Activity** (muscle tone)
5. **Respiration** (breathing rate and effort)

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while [Blue font identifies an ALS skill/treatment or consideration.](#)

APGAR Scoring System

Providers add up these five factors for the Apgar score. Scores are between 10 and 0. Ten is the highest score possible, but few babies get it. That's because most babies' hands and feet remain blue until they have warmed up.

What Does The Score Mean?

A baby who scores a 7 or above on the test is considered in good health. A lower score does not mean that the baby is unhealthy. It means that the baby may need some immediate medical care, such as suctioning of the airways or oxygen to help him or her breathe better. Perfectly healthy babies sometimes have a lower-than-usual score, especially in the first few minutes after birth.

Sources cited: <https://kidshealth.org/en/parents/apgar0.html>

Documentation/Important Highlights

- Document at 1 and 5 minutes.
- Infants with a score of 7-10 usually require supportive care only.
- A score of 4-6 indicates moderate depression.
- Infants with a score of 3 or less require aggressive resuscitation.

Protocols read top to bottom, left to right. Black font identifies BLS skills/treatments or considerations, while Blue font identifies an ALS skill/treatment or consideration.

Blood Draw In Reference To Arkansas Code § 5-65-204

INDICATION

Following Arkansas Code § 5-65-204 (c-1), when presented with documentation by police:

- (1) Blood may be drawn by a person who is licensed, certified, or otherwise authorized by law to perform venous blood draws when a person consents to the procedure or when a warrant or court order has been issued to take a sample of the person's blood.
- (2) When a blood sample is taken at the request of a law enforcement officer based on exigent circumstances, the blood may only be drawn by a physician or a licensed nurse.

(e) Upon the request of the person who submits to a chemical test at the request of a law enforcement officer or because a warrant has been issued to take a sample of the person's blood, full information concerning the chemical test shall be made available to the person or to his or her attorney.

(f) A person who is licensed, certified, or otherwise authorized by law to perform a venous blood draw and any institution or entity employing or using the services of the person is not liable for violating a criminal law of this state in connection with withdrawing blood at the request of a law enforcement officer under this chapter.

(2) A person who is licensed, certified, or otherwise authorized by law to perform a venous blood draw, and any institution or entity employing or using the services of the person is immune from civil and regulatory liability in connection with withdrawing blood at the request of a law enforcement officer under this chapter unless the person is negligent in connection with the withdrawal of the blood.

(3) The immunity granted under this subsection is not conditioned upon the existence of express consent, probable cause, a search warrant, or a court order.

PROCEDURE

Follow the NWAMFCA Blood Draw Guideline

BLOOD DRAW

INDICATION

Per your department's protocol.

PROCEDURE

1. Use universal precautions.
2. Discuss this procedure with the patient as per guidelines and answer all of the patient's questions.
3. Obtain consent.
4. Select the vein and prep as usual.
5. Select appropriate blood-drawing devices.
6. Draw appropriate tubes of blood for lab test
7. Place blood into blood tubes immediately.
8. Label all blood with a minimum of the Patient's name, Date, Time of Draw & Initial
9. Assure that the blood samples are labeled with the correct information.
10. Deliver the labeled blood tubes to the appropriate individual at the hospital.

CONSIDERATIONS

1. The size of the IV you are drawing from affects the blood as it is drawn. Smaller catheters (smaller than 20 – 22 gauge) should be avoided
2. If drawing blood with a syringe – do not pull back firmly because it will cause turbulence in the blood flow into the syringe and rupture the cell membrane – thus destroying your blood sample
3. If you are unable to reach the hospital in a timely manner – less than an hour – your blood sample will not be useable by the lab.
4. Never shake the tubes. This will rupture the cell membrane and destroy your sample. All tubes except the red top should be immediately - gently agitated a few times to mix the chemicals in the tube with the blood to preserve it for testing.
5. The blood tubes are vacuum sealed to draw in the proper amount of blood to mix with the chemical agent. A minimum of 2cc per tube is required. – Red top tubes do not have chemicals and should be filled last with whatever blood is remaining – no minimum.

Reference/Guide/Aids

- [Tube-Collection-guide](#)

CERTIFICATION REQUIREMENTS

Advanced EMT

PARAMEDIC

Blood Glucose Analysis

Indications

- Patients with suspected hypoglycemia (diabetic emergencies, change in mental status, bizarre behavior, etc.)
- Altered mental status

Procedure

1. Gather and prepare equipment.
2. The procedure can be performed simultaneously with intravenous access or a finger stick.
3. Place the correct amount of blood on the reagent strip or site on the glucometer per the manufacturer's instructions.
4. Time the analysis as instructed by the manufacturer.
5. Document the glucometer reading and treat the patient as indicated by the analysis and protocol.
6. Repeat glucose analysis as indicated for reassessment after treatment and as per protocol.

CERTIFICATION REQUIREMENTS

- EMT
- ADVANCED EMT
- PARAMEDIC

NOTES

BLOOD PRODUCTS MONITORING AND RE-INITIATING

Purpose

- Blood products must be initiated prior to the transport of a patient and started by the hospital staff. If a patient requires the administration of blood products during transport a paramedic can continue the administration. In the event the blood product requires replacement, the paramedic can hang another bag of product. If infiltration occurs a paramedic can re-initiate an IV and restart the blood product.
- Vital signs should be taken and recorded at least every 5 minutes. The time of transfusion should be documented. This is the time when the blood actually enters the vein.

Procedure

1. **Continued monitoring of already established blood products requires the following:**
 - a. An order for the transfusion with the flow rate documented
 - b. Verify IV access patency. Must be 20 gauge or larger.
 - c. Blood tubing must be a dedicated line. It may not be piggybacked into existing lines. No medications or solutions other than 0.9% Normal Saline may be mixed with or run concurrently with blood.
 - d. No more than 2 units may be infused through the same blood tubing. The Saline must be changed when the tubing is changed. If a leukocyte reduction filter is used, only one unit of blood may be infused through the tubing and filter.
2. **Replacing blood products:**
 - a. Review the orders from the facility.
 - b. Be sure you do not need to replace the blood-Y tubing. Do not use a 6" or 7" extension set.
 - c. Turn blood slowly end-over-end to mix blood (do not shake) and observe contents for change in color, consistency, or presence of unusual particulate matter.
 - d. Spike the bag containing the blood and hang with the ordered flow rate.
3. **Reinitiating infiltrated IV:**
 - a. Monitor the IV insertion site as usual. If there are signs of infiltration, reinitiate as per IV access protocol.
 - b. Remember a 20 gauge or larger must be used.

BLOOD PRODUCTS MONITORING AND RE-INITIATING

Precautions:

- Several types of blood transfusion reactions can occur during or up to 96 hours after infusion. Symptoms range from mild fever to life-threatening anaphylactic shock. If a reaction is suspected, stop the transfusion immediately and contact medical control.

Transfusion Reactions

- Hemolytic reactions occur when the recipient's serum contains antibodies directed against the corresponding antigen found on donor red blood cells. This can be an ABO incompatibility or an incompatibility related to a different blood group antigen.
- Disseminated intravascular coagulation (DIC), renal failure, and death are not uncommon following this type of reaction. The most common cause for a major hemolytic transfusion reaction is a clerical error, such as a mislabeled specimen sent to the blood bank, or not properly identifying the patient to whom you are giving the blood.
- **DO NOT ASSUME IT IS SOMEONE ELSE'S RESPONSIBILITY TO CHECK!**

Allergic Reactions:

- Allergic reactions to plasma proteins can range from complaints of hives and itching to anaphylaxis, and febrile reactions. White blood cell reactions (febrile reactions) are caused by patient antibodies directed against antigens present on transfused lymphocytes or granulocytes.
- Symptoms usually consist of chills and a temperature rise > 1 degree C.

Transfusion-related acute lung injury (TRALI):

- TRALI is caused when plasma contains HLA or granulocyte-specific antibodies which correspond to antigens found on donor WBCs.
- Granulocyte enzymes are released, increasing capillary permeability and resulting in sudden pulmonary edema.
- Most often occurs with the administration of blood products with plasma, such as FFP.

Bacterial Contamination:

- Bacterial contamination of blood can occur during collection. Bacteria can grow during storage at room temperature and during refrigeration (psychrophilic organisms). Transfusing a contaminated unit can result in septic shock and death.

Circulatory Overload:

- Circulatory overload can occur with the administration of blood or any intravenous fluid, particularly in patients with diminished cardiac function.

CERTIFICATION REQUIREMENTS

- EMT-P

BLOOD PRODUCTS MONITORING AND RE-INITIATING

NOTES

- Transfusion Associated Circulatory Overload (TACO) is essentially a CHF exacerbation from the acute increase in intravascular volume. At-risk populations include the elderly and those with pre-existing cardiac disease. Preventative strategies could include transfusing units over a longer period or administering concomitantly with diuretics.
- Transfusion Related Acute Lung Injury (TRALI) is described as a non-cardiogenic pulmonary edema thought to be secondary to increased vascular permeability. Patient may require intensive airway support, including intubation, until it does. Diuresis is generally less effective in cases of TRALI.

CAPNOGRAPHY - ELECTRONIC & Colorimetric / ETCO2

Indications

- Any Intubated Person
- Altered LOC
- Respiratory Distress
- Congestive Heart Failure
- COPD
- Suspected Traumatic Brain Injury
- Allergic Reactions
- Burn Victims
- Pharmacologically altered persons
- Sepsis


Procedure

Electronic

- Monitors equipped with this capability will have one disposable device for ET tube placement and a Nasal Cannula like device for monitoring naturally expired air. ETT Versions should be used on all intubated patients to confirm placement. Normal range for CO2 is 35-45mmHg. A square waveform (see below) indicates no bronchospasms treatable with bronchodilator medications. Audible wheezing with a square waveform may be considered Cardiac Asthma. Care should be taken to ensure readings are accurate by checking placement and fouled or faulty sensors.
 1. Allow time for the unit to calibrate and give readings
 2. Apply the device to Patient
 3. Change waveform to view on monitor
 4. Note values and waveform in both Pre and Post airway management
 - a. Breakthrough breathing during intermittent positive pressure ventilation (IPPV) makes a cleft in the plateau phase of the capnogram that is known as curare cleft, indicating loss/lack of sedation/paralysis; upon evaluation consider/evaluate continued sedation/paralysis of the patient.
 5. Adjust ventilation to meet normal values when appropriate
 6. ETCO2 should be continuously monitored throughout transport
 7. Vomit or other bodily fluids may clog the sensor and require you to change to another one.

CAPNOGRAPHY - ELECTRONIC & Colorimetric / ETCO₂

8. Imperfect positioning of the nasal cannula / capnography device may cause distorted readings. Unique nasal anatomy obstructed nares and mouth breathers can also cause this problem. Oxygen from the mask may lower the reading by 10%.
9. Document PCR values and waveform after intubation and after each time you move the patient.



Procedure

Colorimetric

- Attach End-Tidal CO₂ Detector between the supra glottic airway or endotracheal tube and ventilation bag.
 1. The color will change according to the concentration of CO₂ in the exhaled air:

Purple	To	Yellow	Proper Placement
Purple	To	Tan	Indeterminate
No Change			Improper Placement

CERTIFICATION REQUIREMENTS

Advanced EMT
Paramedic

NOTES

Carbon Monoxide Oximetry

Indications

- Firefighters in rehab or other patients in or around fire
- Persons found in a potentially high CO environment
- Patients with weakness, flushed skin, and/or “flu” like symptoms
- Any person working around any combustion-generated device
- May be considered for all patients

Procedure

- Outside of environmental sources, individuals with hemolytic anemia, sepsis, and critical illness can have higher than normal CO levels. Patients with CO poisoning may have a wide range of varying signs and symptoms including
 - Seizures
 - Lethargy
 - Tachycardia
 - Confusion
 - Nausea
 - Unconsciousness.
1. Turn the machine on. Some CO monitors can take longer to “warm up”
 2. Apply probe to patient’s finger. Most CO monitors are very light-sensitive and do not work as well on ears lobes and feet. Cover the finger probe with dark material (not a white towel – dark material)) for better results
 3. Some monitors have a PI indicator bar used to indicate accuracy. This should be monitored for fullness and continuity with the heart rate
 4. Allow time for the machine to get an accurate reading. (up to 1 minute)
 5. Note CO levels. Use the chart below for reference.
 6. The accuracy of a SpCO monitor can vary by 3%
 7. Pregnant women should always be transported to a hospital. Fetal CO is much higher than in mothers.

0-5%	Considered normal in non-smokers. When >3% with symptoms, consider high-flow O2 and evaluate the environment for possible CO sources. Consider transport if symptoms persist.
5-10%	Considered normal in smokers, abnormal in non-smokers. If symptoms are present, consider high flow O2 and inquire if others are ill. Consider transport if symptoms persist. <u>Alert Fire Department to monitor air quality for high CO levels.</u>
10-15%	Abnormal in any patient. Assess for symptoms, and consider high-flow O2. Evaluate others for illness. These patients should be transported to the local ED. <u>Alert Fire Department and monitor air quality for high CO levels.</u> Consider Placing the patient on CPAP/BIPAP

Carbon Monoxide Oximetry

>15%	Significantly abnormal in any patient. Administer high-flow O ₂ , assess for symptoms, and evaluate others for illness. These patients should be transported to the local ED. <u>Alert Fire Department to evaluate air quality for high CO levels.</u> Place patient on CPAP/BIPAP
>20%	<u>Alert Fire Department to evaluate air quality for high CO levels.</u> Place Pt on CPAP/BIPAP

CERTIFICATION REQUIREMENTS

- EMR
- EMT
- Advanced EMT
- Paramedic

NOTES

- Consider administration of CynoKit.

CHEST DECOMPRESSION

I. I. Purpose:

To establish indications, guidelines, and the standard procedure for performing a needle thoracostomy in the pre-hospital setting by paramedics.

II. II. Policy:

- A. Needle Thoracostomy may be established on a rapidly deteriorating patient with severe respiratory distress who has signs and symptoms of life-threatening tension pneumothorax such as:
 - 1) Progressively worsening dyspnea
 - 2) Hypotension
 - 3) Shock
 - 4) Decreased or diminished breath sounds on the affected side (required)
 - 5) Distended neck veins
 - 6) Tracheal deviation away from the affected side
 - 7) Traumatic cardiac arrest with tension pneumothorax suspected
- B. Contraindicated when unable to locate landmarks, or tension pneumothorax is not suspected
- C. Risks and complications to pleural decompression include but are not limited to:
 - 1. Lacerated lung tissue
 - 2. Pneumothorax
 - 3. Subcutaneous emphysema
 - 4. Intercostal vein and/or artery hemorrhage
- D. Document the following in the patient care record:
 - 1. Signs and symptoms indicating a need for the procedure
 - 2. Location, number of attempts, size needle used
 - 3. Complications
 - 4. Response to treatment
- E. For Adult patients (> 14 years old or larger than pediatric measurement tape) this procedure is a standing order
- F. For Pediatric patients < 14 contact Medical Control for instruction

III. III. Procedure:

- A. Equipment needed:
 - 1. Alcohol pads
 - 2. 10–14-gauge angiocath or commercially available product
 - 3. 10- or 20-ml syringe
 - 4. One-way valve or cover
 - 5. Tape
- B. Explain the procedure to the patient and place in an upright position if tolerated


CERTIFICATION REQUIREMENTS:

- PARAMEDIC

CHEST DECOMPRESSION

C. Prepare the area with alcohol wipes, either:

1. In the second intercostal space, midclavicular line, **OR**
2. Between the fourth and fifth intercostal space, anterior axillary line
 - 1) Pregnant patients should have the procedure performed between the third and fourth intercostal spaces in the anterior axillary line



- D. Insert needle perpendicular to the chest wall between in the intercostal space, just above the rib
1. The neurovascular bundle runs along the inferior rib, and incorrect placement could result in life-threatening bleeding
- E. Advance the needle until the pleural space is entered, as evidenced by one or more of the following:
1. A “popping” sound or “giving way” sensation
 2. A sudden rush of air
 3. Ability to aspirate free air into the syringe
- F. Remove the needle and fully insert the cannula in place
- G. Place a one-way valve, and secure it to the chest with tape

CERTIFICATION REQUIREMENTS:

- PARAMEDIC

CHEST DECOMPRESSION

- H. Evaluate the effectiveness of the procedure by:
 - 1. Immediate and obvious improvement in respiratory status
 - 2. Improved vital signs
 - 3. Improved bilateral lung sounds
- I. If no improvements are noted, or there is no evidence of entering the pleural space, consider using the other site on the same side of the patient
- J. Move 1 cm toward the midaxillary line and perform decompression as above.
- K. Secure the cannula with dressing and tape allowing the cannula to remain in place
- L. May repeat the procedure in the second approved location if no improvement is seen, or symptoms recur
 - 1. Tension pneumothorax may recur if the air is not able to escape the decompression site with the one-way valve
- M. Contact the destination facility and advise that you have performed a Pleural Decompression

CERTIFICATION REQUIREMENTS:

- PARAMEDIC

Regional Protocol **Guideline Section – To be used only as indicated in Protocol Section**
CONTINUOUS POSITIVE AIRWAY PRESSURE (CPAP)/ BIPAP

INDICATIONS

- Patients with moderate to severe respiratory distress secondary to pulmonary edema (wet lung sounds), COPD, asthma, or submersion incidents.
- Indications of moderate to severe respiratory distress can include:
 - Retractions
 - Accessory muscle use
 - Tachypnea (respiratory rate >25/min)
 - Pulse oximetry reading <90%
 - Tripod position
 - 1 to 2 word sentences

CONTRAINDICATIONS

- Respiratory or cardiac arrest
- Systolic blood pressure <90mmHg
- Severely depressed level of consciousness
- Inability to maintain airway patency
- Major trauma, especially head injury with increased ICP or significant chest trauma
- Facial trauma affecting mask seal
- Signs and symptoms of pneumothorax
- Vomiting
- Gastric distention or active GI bleed
- Inability to tolerate mask on face
- CPAP is contraindicated during inferior wall STEMI with right ventricular involvement or with unknown right ventricular involvement.

PROCEDURE

1. Advise the patient of the need for and efficacy of CPAP therapy
2. Continuously monitor the patient's vitals including ECG, pulse oximetry, and waveform capnography
3. Place patient in a seated position with legs dependent or position of comfort
4. Set the CPAP pressure at 8 cmH₂O or BIPAP – Insp Pressure (IPAP) at 12 and PEEP at 8
5. Use appropriate sized and fitted mask (note patient may tolerate better if they hold the mask as opposed to strapping to their face)
6. Allow the patient to adjust to the procedure
7. Titrate pressure up to patient response or comfort
8. Do not delay other EMS treatment guidelines (Nitro, Morphine, Bronchodilators...)
9. Continuously monitor patient for improvement or decline

NOTES

- Monitor O₂ usage as some units will consume oxygen rapidly
- COPD and Asthma patient will typically require lower pressures around 5 cmH₂O
- Pulmonary edema may require pressures of 10 cmH₂O
- If patient fails to show improvement, endotracheal intubation should be considered.
- Monitor EtCO₂ closely for patients who are known CO₂ retainers
- Document all use of CPAP with thorough assessments including a full set of vitals before and after treatment.

CERTIFICATION REQUIREMENTS:

- ADVANCED EMT
- PARAMEDIC

CRICOTHYROTOMY--NEEDLE


Indication

- Pediatric patients, 0 to 9 yrs old, with complete airway obstruction where all other efforts at airway management have failed.

Procedure

1. Continue attempts at ventilation while assembling equipment.
2. The cricothyroid membrane is best identified by palpating the laryngeal prominence at the anterior, superior aspect of the larynx. Approximately one of the patient's fingerbreadths inferior to the laryngeal prominence is a small depression bounded on its inferior aspect by a rigid, horizontal structure, the cricoid cartilage. This small depression is the cricothyroid membrane and the rigid structure below is the cricoid cartilage. Prep the site with Betadine.
3. Attach a syringe to an over-the-needle catheter.
4. Insert the needle down through the midline of the cricothyroid membrane at a 45-60 angle toward the patient's carina. Take caution not to pass through the back of the trachea.
5. During insertion, apply negative pressure to the syringe. The entrance of air into the syringe indicates that the needle is in the trachea.
6. Remove the needle and syringe, leaving the catheter in the trachea.
7. Stabilize the catheter, connect a 3.0 ETT adapter to the hub of the catheter, and ventilate with a bag-valve device and 100 % oxygen.
8. Remove the bag valve to allow for exhalation.
9. Observe for chest rise, increased pulse ox, and other clinical signs of oxygenation.

CRICOTHYROTOMY--NEEDLE



CERTIFICATION REQUIREMENTS

- Paramedic

NOTES

-

CRICOTHYROTOMY--SURGICAL

Indication

- Complete airway obstruction not responding to all other attempts to ventilate
- Destructive facial injuries
- This procedure shall be used only after all other attempts of establishing an airway and ventilating a patient have failed. Limited to patients over 10 years old.

Procedure

- if using a commercial device – follow their instructions on use
- Assemble Equipment


○

# 20 Scalpel	Tracheal Hook
Bougie	Tracheal Tube (5.5 – 6.5) with 10 cc Syringe
Bag-valve-mask	Cloth Tape for securing the tube
Capnography	4x4 gauze to control bleeding

1. **Puncture** (do not incise) the lower third of the cricothyroid membrane
 - a. Cleanse & Locate the cricothyroid membrane
 - i. Move the finger down from the top of the thyroid cartilage to the membrane.
 - ii. Move finger up from jugular notch to membrane.
 - b. Stabilize the thyroid cartilage with fingers. Cleanse with betadine.
 - c. Puncture the lower 1/3 of the cricothyroid membrane with a # 20 scalpel until the widest part of the blade is inserted beneath the skin.
2. **Insert the tracheal hook**
 - a. Insert horizontally along the flat of the scalpel and then turn caudally making sure you have hooked all tissue planes. Remove scalpel.
 - b. Retract at a 45° angle up and toward the feet.
3. **Insert tracheal tube**
 - a. A. Insert bougie straight down then immediately turn down toward the chest. Feel your way... the bougie should move freely. Continue insertion of bougie until "hold up" confirms tracheal placement.
 - b. Insert the tube until the cuff is in the trachea. (Cord mark should barely disappear)
 - c. Inflate the cuff
4. **Ventilate**
 - a. Check placement with capnography, stethoscope, pulse oximetry, bougie, and or esophageal detection device.

CRICOTHYROTOMY--SURGICAL

- b. Secure the tube with 1" cloth tape.
- c. Control bleeding with dressings and direct pressure.



CERTIFICATION REQUIREMENTS

- Paramedic

NOTES

- A thinner scalpel blade will not make an adequate opening without an incision.
- Beware false passage into tissue planes around the trachea - the tube should pass easily once the tip enters the trachea.
 - If this occurs, back the tube out and correct by reinserting the bougie.
- Esophageal detector device easily demonstrates false passage.
- Use a scalpel, then hook, then bougie with your dominant hand.
- Practice this in your head and blindfolded (feel your way)... You may not be able to see because of bleeding.

CRIME SCENE

Policy:

The primary responsibility of EMS is patient care; however, EMS should take all possible precautions to preserve evidence while at a crime scene.

Purpose:

To establish guidelines by which EMS personnel may provide patient care in a potential or known crime scene.

Procedure:

1. Make certain that law enforcement has been notified or is on scene (preferred).
2. The primary EMS responsibility is to provide medical help to a patient or patients. The secondary responsibility is to preserve evidence.
3. The entire scene (including roadway, driveway, parking lot, outside areas) may contain evidence, which may be contaminated or destroyed by EMS.
4. Limit the number of EMS and fire responders entering the crime scene. All personnel should enter and exit by one route, taking care not to touch or move anything not directly related to the care of the patient.
5. Weapons should not be touched or moved by EMS or fire personnel. If a weapon presents a real threat or hindrance to patient care, have law enforcement secure it if possible.
6. The clothing and personnel effects of the patient are evidence. If clothing must be removed from the patient to provide care, EMS or Fire should use care to cut around holes or tears in the clothing and not cut through them.
7. EMS and fire personnel are not detectives. Searches of the premises should be left up to law enforcement.
8. There should be no cleanup of the scene prior to an “ok” from law enforcement. Used dressings, packaging, and other EMS trash should be left in place until after other evidence has been processed by law enforcement.
9. The PCR (patient care report) should reflect the name(s) of all EMS personnel who have physical contact with the scene, including students and riders.
10. The PCR should contain only factual information obtained by EMS about the patient and the patient’s relationship to the scene. The PCR should describe the injuries to a patient and not the apparent cause of those injuries.
11. The PCR will become part of the legal record of the incident.

CERTIFICATION REQUIREMENTS:

- EMR
- EMT
- ADVANCED EMT
- PARAMEDIC

DECONTAMINATION - EMERGENCY

INDICATIONS

- If hazardous materials are suspected, immediately activate the Hazmat Team
- Emergency decontamination shall be performed whenever a patient has been contaminated with a chemical that may present a risk to the patient, caregiver or hospital staff.
- Activate the Hazmat Team for large-scale or multiple-patient contamination, or hazardous environment situations.

PROCEDURE- FOLLOW HAZMAT TEAM RECOMMENDATIONS – SOME RECOMMENDATIONS MAY INCLUDE:

1. Remove the patient from the Hazard Area (Hot Zone). Do not enter Hot Zone unless authorized to do so.
2. If patient is capable have the patient follow the procedure without assistance.
3. Take actions to preserve the patient's dignity.
4. Remove contaminated clothing. This may be accomplished simultaneously with rinsing especially if the patient is critical or chemical burns are occurring.
5. Triple bag contaminated clothing (valuables may be bagged separately)
6. Rinse the patient with copious amounts of water at low pressure. If patient is stable and/or staffing and equipment allows take actions to protect the environment (plastic sheeting, impoundment, etc.)
7. Wash the patient with mild soap, if available, and rinse.
8. Re-dress patient in hospital gown and/or cover with sheet and blanket(s)
9. Procedure need not take too much time—balance amount of decontamination with the need to reduce the risk to all involved.
10. Notify ER that you have a hazmat patient.

POISON CONTROL: 1-800-222-1222
CHEMTREC 1-800-424-9300

CERTIFICATION REQUIREMENTS

- EMR
- EMT
- ADVANCED EMT
- PARAMEDIC

ELECTRICAL THERAPY

INDICATIONS

- Defibrillation—ventricular fibrillation or pulseless ventricular tachycardia
- Cardioversion—unstable tachycardia
- Transcutaneous Pacing—symptomatic bradycardia

PROCEDURE (FOLLOW CURRENT AHA STANDARDS AND GUIDELINES)

Defibrillation:

1. Initial treatment for ventricular fibrillation and pulseless ventricular tachycardia is immediate defibrillation.
2. Every minute that passes reduces the chances of successful cardioversion.
3. Apply defibrillation pads to patient's chest.
4. Perform quick look and assess for shockable rhythm.
5. Charge to highest joule setting for monophasic defibrillators.
6.

Physio Control Recommendations	ZOLL Recommendations	Phillips Recommendations
200 J, 300J, 360J	120J, 150J, 200J	150J, 150J, 150J
7. Deliver shock and begin CPR immediately for 5 cycles.
8. If the patient re-arrests after converting from a shock, use the energy level of the last shock used before increasing.

Cardioversion:

1. Initial treatment for unstable tachycardia and the subsequent treatment of tachycardia not responding to antiarrhythmic medications.
2. Do not delay the delivery of cardioversion for IV attempts, medication administration, or failure of the EKG monitor to "synch".
3. Apply three or four lead EKG monitor cables.
4. Assess rhythm and determine if patient is unstable.
5. Apply pads to patient's chest.
6. Depress the "synch" button. If there is difficulty synchronizing increase QRS size.
7.

Physio Control Recommendations	ZOLL Recommendations	Phillips Recommendations
100 J, 200J, 300J, 360J	100J, 120J, 150J, 200J	100J, 120J, 150J
8. Deliver shock and assess for rhythm change.
9. If shock will not deliver, Turn off synchronizer and shock patient.

Transcutaneous Pacing:

1. If patient is symptomatic, do not delay TCP while awaiting IV access.
2. Apply three or four lead EKG monitor cables.
3. Apply pads to patient's chest. Anterior-Posterior or Anterior-Lateral position may be used.
4. Set rate. Increase output until capture is achieved. Access pulse on right side of the body for mechanical capture.
5. Continue pacing at an output level slightly (10%) higher than threshold of initial capture.
6. ER notification to the is required if patient is being paced so they can prepare.

CERTIFICATION REQUIREMENTS

- EMR (defibrillation only with AED)
- EMT (defibrillation only with AED)
- ADVANCED EMT (defibrillation only)
- PARAMEDIC

GLASGOW COMA SCORE

ADULT

MOTOR RESPONSE		EYE OPENING		VERBAL RESPONSE	
Obeys commands	6	Spontaneous	4	Oriented	5
Localizes	5	To Voice	3	Confused	4
Withdrawal	4	To Pain	2	Inappropriate	3
Flexion	3	None	1	Incomprehensible	2
Extension	2			None	1
None	1				

PEDIATRIC

PEDIATRIC GLASGOW COMA SCALE (PGCS)				
	> 1 Year		< 1 Year	Score
EYE OPENING	Spontaneously		Spontaneously	4
	To verbal command		To shout	3
	To pain		To pain	2
	No response		No response	1
MOTOR RESPONSE	Obeys		Spontaneous	6
	Localizes pain		Localizes pain	5
	Flexion-withdrawal		Flexion-withdrawal	4
	Flexion-abnormal (decorticate rigidity)		Flexion-abnormal (decorticate rigidity)	3
	Extension (decerebrate rigidity)		Extension (decerebrate rigidity)	2
	No response		No response	1
	> 5 Years	2-5 Years	0-23 months	
VERBAL RESPONSE	Oriented	Appropriate words/phrases	Smiles/coos appropriately	5
	Disoriented/confused	Inappropriate words	Cries and is consolable	4
	Inappropriate words	Persistent cries and screams	Persistent inappropriate crying and/or screaming	3
	Incomprehensible sounds	Grunts	Grunts, agitated, and restless	2
	No response	No response	No response	1
TOTAL PEDIATRIC GLASGOW COMA SCORE (3-15):				

CERTIFICATION REQUIREMENTS

- EMR
- EMT
- ADVANCED EMT
- PARAMEDIC

INTRAOSSEOUS ACCESS HUMERAL HEAD**INDICATIONS**

- Adult or Pediatric (if appropriate device is available) patient with life threatening illness or injury and urgent need for IV but veins are not readily available. Humeral Head IO is preferred route for decreased pain and increased flow rates. IV is preferable over IO is possible.

CONTRAINDICATIONS

- Fracture (target bone)
- Previous orthopedic procedure of target bone (IO within 24 hours, prosthetic limb or joint)
- Infection at insertion site
- Inability to locate landmarks or excessive tissue

PROCEDURE

- Place the patient in a supine position
- Expose shoulder and adduct humerus (place the patient's arm against the patient's body) resting the elbow on the stretcher or ground
- Palpate and identify the mid-shaft humerus and continue palpating toward the proximal aspect or humeral head. As you near the shoulder, you will note a protrusion. This is the base of the greater tubercle insertion site.
- With the opposite hand, you may consider "pinching" the anterior and inferior aspects of the humeral head while confirming the identification of the greater tubercle. This will ensure that you have identified the midline of the humerus itself.
- Immobilize with sling and swath or commercial sling

CONSIDERATIONS

- Flow rates:
Due to the anatomy of the intraosseous space, flow rates will be slower than those achieved with IV catheters. Use a pressure bag or infusion pump to ensure continuous infusion.
- Pain:
Insertion of the IO needle in conscious patients causes mild to moderate discomfort but is usually no more painful than a large bore IV. IO infusion can cause severe discomfort for conscious patients. Administer 40mg (or 2mLs) 2% IV Lidocaine Prior to IO flush

**CERTIFICATION REQUIREMENTS**

- Paramedic
- Advanced EMT

INTRAOSSEROUS ACCESS TIBIAL**INDICATION**

Adult or Pediatric (if appropriate device is available) patient with life threatening illness or injury and urgent need for IV but veins are not readily available. Humeral Head IO is preferred route for decreased pain and increased flow rates. . IV is preferable over IO is possible.

CONTRAINDICATIONS (consider alternate tibia)

Fracture of the tibia or femur on side of procedure
 Previous orthopedic procedures (IO within 48 hours, knee replacement)
 Pre-existing medical condition or infection near insertion site
 Inability to locate landmarks (significant edema. or excessive tissue at insertion site)

PROCEDURE

Expose the lower leg
 Locate insertion site one finger breath medial of the tibial tuberosity
 Prep the site as per peripheral IV site
 Prepare the IO driver and needle set
 Using aseptic technique, stabilize the leg and insert IO needle
 Remove IO driver from needle set while stabilizing catheter hub.
 Remove stylet from needle set and dispose in sharps container and connect the extension tubing.
 Confirm placement – do not insert needle if it is too short and the 5mm line is not visible after placing through tissue.
 Proper placement?: consider these: IO Needle stands firm, can flush without infiltration..

Pain control and Flushing:

Flush the IO space with 10 ml of fluid (If the patient is conscious, SLOWLY administer 40mg (2mLs) 2% Lidocaine IO and wait up to 2 minutes prior to initial bolus). If pain persists after the bolus of fluids 10- 20 mg of Lidocaine may be administered.
 Connect IV tubing, monitor, and document as per IV access procedure.

CONSIDERATIONS

Flow rates:


Due to the anatomy of the intraosseous space, flow rates will be slower than those achieved with IV catheters. Use a pressure bag or infusion pump to ensure continuous infusion.

Pain:

Insertion of the IO needle in conscious patients causes mild to moderate discomfort but is usually no more painful than a large bore IV. IO infusion can cause severe discomfort for conscious patients. Administer 40mg (or 2mLs) 2% IV Lidocaine Prior to IO flush

CERTIFICATION REQUIREMENTS

- Advanced EMT
- PARAMEDIC



INTUBATION--NASOTRACHEAL (BLIND)**INDICATION**

A spontaneously breathing patient in need of intubation—inadequate respiratory effort, evidence of hypoxia or carbon dioxide retention, or need for airway protection. Patient must be 12 years of age or older.

CONTRAINDICATIONS

- Apneic patients, midfacial fractures, suspected basilar skull fractures, bleeding disorders, taking Coumadin, likely to receive Heparin or thrombolytics, severe nasal trauma, pharyngeal hemorrhage, acute epiglottitis, suspected laryngeal fracture, and suspected increased intracranial pressure.

PROCEDURE

1. Hyperoxygenate patient while preparing equipment.
2. Select the nostril that appears larger and the ET tube at least 1 mm size smaller than that which would be used for orotracheal approach. Pre-flex the ET tube, and lubricate with water soluble gel.
3. Apply topical vasoconstrictor and/or topical anesthetic. If time permits insert a lubricated nasal pharyngeal airway to help dilate the nostril.
4. Place the patient in the “sniffing position” with elevation of the head on a pillow and the jaw forward.
5. Attach the BAAM whistle (if available) on the ET tube. The patient’s breathing will activate the BAAM and whistling will be observed with inhalation and exhalation.
6. Insert the ET tube with tip directed along the floor of the nostril and then in a plane across the midline toward the opposite shoulder so that the tip of the tube will be near the midline at the level of the larynx.
7. Gently advance the ET tube on early inspiration.
8. If using the BAAM whistle the intensity of the whistling will increase as the ET tube is properly placed. Deviation out of the airflow tract will result in immediate decrease and loss of whistle sound. Withdraw the ET tube a short distance and redirect it laterally by twisting the tube, anteriorly by extending the head, or posteriorly by elevating the jaw and/or slight flexion of the neck until the whistle sound is again maximal.
9. Inflate ET tube cuff with appropriate amount of air. Note measurement.
10. All ET tube placements shall be confirmed using the following steps:

Primary Confirmation

- ETCO₂
- Continued increase in whistle through BAAM as ET tube passes through the vocal cords.
- 5 point auscultations anterior L/R, mid-axillary, over epigastrium.
- Chest rise and fall with each ventilation.

Secondary Confirmation

- With a perfusing patient – ETCO₂
- Monitor oxygen saturation and CO₂ levels

11. If still in doubt, remove the ET tube and ventilate the patient with bag-valve mask and follow the oxygen titration protocol.
12. Upon confirmation of correct ET tube placement, secure with appropriate device and note the tube depth measurement.
13. Reassess placement frequently- each time patient is moved, change in patient condition, transfer of care, (consider C Collar), etc.
14. Continuous ETCO₂ via capnography should be applied.

CERTIFICATION REQUIREMENTS

- PARAMEDIC

INTUBATION--OROTRACHEAL

INDICATIONS

- Cardiac arrest with ongoing chest compressions.
- Inability of patient in respiratory compromise to breathe adequately.
- Inability of the patient to protect their airway—coma, areflexia, or cardiac arrest.
- Inability of the rescuer to ventilate the unresponsive patient with bag-valve device.

Endotracheal intubation is the gold standard of airway management. However, failure to intubate does not mean failure to ventilate.

PROCEDURE

1. Hyperoxygenate while preparing intubation equipment. Remember suction.
2. For patients with suspected spinal injuries, maintain neutral position of the cervical spine during intubation. In all other patients use the position that best accommodates visualization of the vocal cords.
3. Insert the laryngoscope blade into the oropharynx to visualize the vocal cords. Avoid pressure on the patient's lips and teeth.
4. Apply downward pressure on the larynx or use the BURP (Backward, Upward, Right, Pressure) to assist in visualization of the cords.
5. While visualizing the cords, insert the proper size ET tube with stylette through the vocal cords, advance one-half to one inch farther.
6. While holding the ET tube in place, inflate the cuff with just enough air to inflate the cuff of the ET Tube..
7. All ET tube placement shall be confirmed using the following steps:

Primary Confirmation	Secondary Confirmation
• <u>Capnography with all advanced airway patients</u> • Direct visualization of ET tube passing through the vocal cords. • 5 point auscultations anterior L/R, mid-axillary, over epigastrium.	• Chest rise and fall with each ventilation. • With a perfusing patient -Monitor oxygen saturation.

8. If at any time placement of the ET tube is in doubt, insert the laryngoscope into the oropharynx and note if the ET tube passes through the vocal cords.
9. If still in doubt, remove the ET tube and ventilate the patient with bag-valve mask and follow the oxygen protocol..
10. Upon confirmation of correct ET tube placement, secure with appropriate device and note the tube depth measurement.
11. Continuous ETCO₂ via capnography should be applied.
12. Reassess placement frequently—each time patient is moved, change in patient condition, transfer of care, etc.

CERTIFICATION REQUIREMENTS

- PARAMEDIC

IV ACCESS: EXISTING CENTRAL VENOUS LINES

INDICATION

In cases of severe illness or injury requiring immediate fluid or drug administration when peripheral IV access has been unobtainable or is unlikely

PRECAUTIONS

It is important to remember that many patients with central venous lines in place are immunosuppressed or severely debilitated. Thus, they are very susceptible to routine pathogens. Special care should be taken to avoid contamination.

- Access of a port used for dialysis is discouraged. If you are unclear or have doubts you should consider other means of vascular access. Use patient history to help make this decision.

Introduction of air can be extremely hazardous. DO NOT remove injection cap from catheter or allow IV fluids to run dry.

PROCEDURE

BROVIAC / HICKMAN / GROSHONG AND OTHER DOUBLE/TRIPLE LUMEN CATHETERS:

1. Select appropriate port for access. If more than one port is seen then access the venous side. Although blue generally indicates the venous side, many different colors are available. You cannot use color only as a guide. Aspirate blood and examine. Do not use a line with gauze wrapped around the port or with a medication sticker.
2. Lay patient supine.
3. Thoroughly cleanse injectable port cap prior to use. (Red cap is likely arterial and should not be accessed)
4. Aspirate 10cc of blood from catheter (this prevents an inadvertent anticoagulant bolus from occurring). If you are unable to draw blood, reposition patient and try again. If problem persists do not use line.
5. Confirm venous access by examination of aspirated blood and then discard blood. If in doubt use other port or consider use of other vascular access methods.
6. Attach IV line to injection port. Begin IV fluid flow and adjust according to patient presentation. Make sure clamps are closed except when aspirating, flushing or infusing.
7. Inject all necessary medications and fluids. Always aspirate before flushing or infusing. This prevents embolization of any clots that may have formed since last administration.

NOTE: If at any time you are unable to aspirate blood or infuse fluids, do not use line as

Continued on next page

IV ACCESS Continued:
EXISTING CENTRAL VENOUS LINES
 (page 2/2)

PICC LINE ACCESS

1. Select desired port, when two sizes are available select the larger of the two.
2. Thoroughly cleanse injectable port prior to use.
3. Aspirate 6-10cc of blood and discard. Attempt to inject 5cc of NS into port. If resistance is met, withdraw needle and attempt same procedure on different port. Do this until you find a catheter that does not present with resistance to administration of NS. If resistance continues, do not use PICC line.
4. If you are unable to draw blood or suspect arterial placement do not use PICC line.
5. Attach IV tubing, open line, and ensure patency. You should aspirate 5cc of blood before each flush or medication administration. Begin IV fluid flow and adjust according to patient presentation.

PORT ACCESS

1. **Use only if access to a Huber needle is available.**
2. Position patient supine.
3. Locate the sight by visualization and palpation, these ports are generally found in the upper right chest and present as a dome shaped protrusion.
4. Thoroughly cleanse site prior to use. The preferred method is to use Iodine and vigorous rubbing in a circular pattern starting at the center and working your way out. Allow time to dry completely.
5. Using a non-coring Huber Needle attached to a syringe, insert into the site at a 90-degree angle until resistance is met. This means you have reached the back plate of the hub.
6. Aspirate 10cc of blood and discard. Then draw additional blood for lab work and inject 5-10cc of NS. If resistance is met or blood cannot be aspirated change the patient's position and try again. If problem persists withdraw needle and consider alternative vascular access.
7. Remove syringe, attach IV tubing, open line, and ensure patency.
8. Secure the site with large occlusive dressing.

**PICC LINE
Central Catheter):
SUBCUTANEOUS**



**(Peripherally
INTERNAL
INFUSION**



**Inserted
PORTS:**



CERTIFICATION REQUIREMENTS:

- Paramedic

IV ACCESS—EXTERNAL JUGULAR

INDICATION

Inability to obtain an IV or IO on a critically ill or injured patient ≥ 12 years of age who requires intravenous access for fluid or medication administration, where no obvious peripheral site or Intraosseous is accessible.

PROCEDURE

1. Place the patient in a supine, head-down position. This helps distend the vein and prevents air embolism.
2. Turn the patient's head toward the opposite side if no risk of cervical injury exists.
3. Prep the site as per peripheral IV site.
4. Align the catheter with the vein and aim downward toward the same side shoulder.
5. "Tourniquet" the vein lightly with one finger above the clavicle and cannulate the vein in the usual method.
6. Remember that air can easily enter the blood stream from a large Vein such as this so quickly attach ext. set to prevent air from entering the blood stream.
7. Attach the IV and secure the catheter avoiding circumferential dressing or taping.

CERTIFICATION REQUIREMENTS

- PARAMEDIC
- Successfully complete an annual skill evaluation inclusive of the indications, contraindications, technique, and possible complications of the procedure.

KING AIRWAY**INDICATION**

- An alternative to endotracheal intubation for airway management in patients greater than 35 inches tall to secure a patent airway and deliver ventilations.

CONTRAINDICATIONS

- Responsive patients with an intact gag reflex.
- Patients with known esophageal disease.
- Any patients that have ingested caustic substances.
- Patients who are less than 35 inches tall.

PROCEDURE (Reference King EMS kit insert)

Use BSI including gloves, mask, and eye protection. Assemble the equipment while continuing ventilations.

1. Choose the correct tube size based on the patient's height.

Pt. height	Size	Color
35"-45"	2	Green
41"-51"	2.5	Orange
4'-5'	3	Yellow
5'-6'	4	Red
> 6'	5	Purple

2. Check inflatable cuffs for leaks.
3. Apply water soluble lubrication to the tip.
4. Prepare and turn on suction.
5. Apply chin lift and introduce the King airway in to the corner of the mouth.
6. Advance tip under the base of the tongue while rotating the tube back to midline.
7. Without excessive force, advance the tube until the base of the connector is aligned with the patient's teeth or gums.
8. Inflate cuff based on tube size Using a pressure gauge: 60 cm H₂O. Typical inflation volume is as follows:

Size	KLTD/ml	KLTSd/ml
2	25-35	n/a
2.5	25-35	n/a
3	45-60	40-55
4	60-80	50-70
5	70-90	60-80


9. Attach the BVM. While gently bagging slowly withdraw the tube until ventilation is easy to administer a large tidal volume with minimal airway pressure.
10. Adjust cuff inflation, if necessary, to obtain an airway seal at peak ventilation pressure.
11. Assess for proper tube placement.
 - a. Assess breath sounds; Assure chest rise and fall; Attach patient to continuous end tidal CO₂ monitoring; Continue to reassess that tube is properly placed and that patient ventilation is easy and free flowing with chest rise and adequate breath sounds
 - b. If at any time the provider is unsure of proper placement – deflate cuff, remove and use BVM for ventilation.

NOTES

1. Preparation: Use only water soluble lubricant. Do not apply lubricant near ventilatory openings.
2. Induction: Patient should be "deep enough", do NOT insert the KLTD/KLTSd if the patient is swallowing, retching, moving or gagging.
3. Insertion: Hold the KLTD/KLTSd with the dominant hand at the proximal end (connector) such that insertion will be accomplished in a single, continuous motion. Use the lateral approach with chin lift. Insert the KLTD/KLTSd until the base of the connector is aligned with teeth or gums.
 - TCheck that the blue (pharyngeal) cuff is not visible in the oropharynx.
4. Final Positioning: Withdraw the KLTD/KLTSd until ventilation is optimized. Readjust cuff inflation.
5. Taping: Disconnect the circuit and aggressively tape the KLTD/KLTSd in the midline to the maxilla.
 - For the KLTSd, avoid taping over the opening to the gastric access lumen.

CERTIFICATION**CERTIFICATION REQUIREMENTS:**

- ADVANCED EMT
- PARAMEDIC



MEDICATION ADMINISTRATION

Six “Rights” of Drug Administration:

1. Right Patient
2. Right Drug
3. Right Time
4. Right Dose
5. Right Route
6. Right Documentation

Paramedics should carefully read the drug package and/or label prior to administering any drug to help assure the correct preparation is being administered

Sublingual Administration:

Sublingual medications are placed or sprayed under the tongue and allowed to dissolve. Absorption occurs via the rich supply of superficial vessels under the tongue. Examples include Nitroglycerin and Narcan

Inhalation Administration:

Because the respiratory tract offers an enormous absorption surface with a rich blood supply, drugs administered via this route can have both local and systemic effects. Examples include drugs administered via nebulizer and metered-dose inhaler.

Endotracheal Administration:

Instillation of a drug into the trachea via the endotracheal tube. Patient is then ventilated with a bag valve mask to disperse the drug across the alveoli where it is absorbed into the circulation. Drugs that can be given endotracheal include Lidocaine, epinephrine, atropine and Narcan (LEAN is the mnemonic to help remember this). Endotracheal use of diazepam (Valium) is controversial because it is not water soluble and can be damaging to lung tissue. Diluting drugs given endotracheal with 5—10 ml of normal saline can help absorption. Establish IV access ASAP as repeated dosing down ETT can add excess fluid volume to the lungs. In general, endotracheal doses are increased by 2 to 2.5 times the recommended IV dose.

Procedure:

1. Dilute drug in 5—10 ml of normal saline (Many drugs are diluted as packaged.)
2. Remove the needle from the syringe if possible before instilling medication down the tube. The American Heart Association recommends passing a suction catheter beyond the tip of the endotracheal tube and administering the medication through the catheter. Some preloaded syringes do not have removable needles and, in the interest of time, are used as is. With such syringes, care should be taken to prevent damaging the endotracheal tube with the needle.
3. If CPR is being done, briefly interrupt chest compressions while the drug is instilled into the endotracheal tube.
4. Follow instillation of drug with two to three ventilations via bag-valve mask to disperse drug.
5. Assess the patient's response.

Subcutaneous Injection:

In SC or SQ injection medication is injected into the loose connective tissue between the dermis and muscle layer. This route allows for slow absorption of drugs and is used when a sustained effect is desired.

(CONTINUED ON NEXT PAGE)

MEDICATION ADMINISTRATION – CONTINUED

SQ Injection Procedure:

1. Review 6 Rights
2. Explain procedure to patient
3. Take BSI precautions
4. Select and cleanse site with alcohol and allow it to dry or wipe it dry with sterile gauze before proceeding
5. Pinch the skin up slightly between the thumb and other fingers.
6. Insert the needle using a quick, dart-like motion, using the appropriate angle:
 - When using a 5/8-inch needle, a 45-degree angle should be used with most adults
 - In very obese patients, increase the angle to 60 degrees
 - In very thin patients, reduce the angle to 30 degrees
7. Aspirate to check for blood (if blood is drawn, withdraw needle and discard medication and then prepare another dose)
8. Gently inject medication
9. Discard needle in sharps container
10. Massage injection site to reduce discomfort and disperse medication
11. Consider applying band aid if time permits


Intramuscular Injection:

This route has several advantages over the subcutaneous route: larger amounts of fluid can be injected (up to 5 ml in adults), absorption is faster, and drugs that are irritating to SC tissues are better tolerated when given IM. For volumes greater than 3—5 ml, more than one injection site should be used. Because of the depth of IM injections, special care must be taken to avoid damaging nerves. Common sites used for IM injections include the arm (deltoid), the thigh (rectus femoris or vastus lateralis), and the hip (dorsogluteal or ventrogluteal).

Note: While the deltoid (upper arm) is an easily accessible site and well perfused, it has several disadvantages. It can only accommodate small amounts of fluid (i.e., 1 ml in women and children and up to 2 ml in males with well-developed deltoids). The vastus lateralis and rectus femoris (thigh) are the preferred sites for IM injections in infants.

IM injection procedure:

1. Review 5 rights
2. Explain procedure to patient
3. Take BSI precautions
4. Select appropriate size needle syringe and
5. Select appropriate injection site. Consider:
 - Ability of patient to cooperate
 - Amount of drug to be given
 - Type of drug to be given (very irritating should be given in large gluteal muscles, Promethazine, Hydroxyzine, Diazepam.
6. Cleanse and prep site.
7. Using Z-track technique, Insert needle at 90 degree angle to skin using a quick dart-like motion.
8. Aspirate before injecting to check for blood return. (if blood is drawn, withdraw needle and discard medication and then prepare another dose)
9. Discard needle and syringe appropriately
10. Do not massage needle site if Z-track technique utilized.
11. Consider applying Band-Aid if time permits



drugs
i.e.,

Intravenous Administration:

Administering a drug intravenously places the medication directly into the patient's bloodstream. Therefore, the onset of action is more rapid than with other parenteral routes. The general rule to follow when selecting a catheter is to choose the smallest bore that will accomplish the purpose of the IV. However, large-bore catheters (18-14) should be selected for adult patients with life threatening emergencies in which rapid fluid replacement is required. It is also important to note that damage to veins and other complications of IV therapy are often result of utilizing large catheters.

If at all possible, IV catheters should be placed in the hands or forearm with the antecubital space reserved for patients in cardiac arrest or patients with life threatening emergencies requiring rapid fluid administration. The external jugular vein along with veins in the feet or ankle can also be used in emergent patients with limited accessible upper extremity IV sites.

Troubleshooting IV Lines:

In the event that an IV will not infuse, the following steps should be taken:

1. Ensure that the tourniquet has been removed.
2. Check the line for kinks or obstruction caused by the patient or nearby equipment
3. Check to see that the roller clamp and line clamps are open.
4. Raise the height of the IV infusion bag.
5. Gently manipulate the position of the IV line and the patient's extremity.

Do not forcefully irrigate an apparently occluded line.

Procedure for administering IV medication:

1. Review 6 rights.
2. Take BSI precautions.
3. Typically select and cleanse the most distal medication port (closest to IV site).
4. Check patency of the primary line by aspirating gently and checking for blood return.
5. Occlude the tubing above the injection port by using the roller clamp or crimping the tube with the other hand
6. Administer the medication at the specified flow rate.
7. Release the occlusion on the tubing.
8. Readjust the flow rate
9. Document the date, time, and amount of drug administered.
10. Assess the patient's response to the drug.

Procedure for administering IV piggyback (IVPB) medication:

1. Review 6 rights
2. Take BSI precautions
3. Prepare the medication
4. If not using premix, cleanse injection port on bag used for IVPB and inject medication into bag.
5. Invert the bag several times to mix the solution.
6. Attach a medication label to the bag indicating the name and amount of drug injected along with the date and time.
7. Connect an appropriate drip set to IVPB bag and flush tubing.
8. Connect to port on main line (distal main line roller clamp) with either a needle or needle-less adapter.
9. If not using infusion pump, hang IVPB bag higher than main bag and adjust flow rate.
10. Main line can be clamped off at this time or infused along with IVPB.

(CONTINUED ON NEXT PAGE)

Intraosseous Administration:

The IO route is intended for short-term use until other venous access can be obtained. It is recommended that an IO line be placed after 90 seconds or two unsuccessful attempts to start a peripheral line. The distribution of fluid and drugs given the IO route is similar to that of IV administration. Fluids or medications are injected into the bone marrow cavity and pass into the venous sinusoids to the central venous channels and then to the systemic circulation via the emissary and nutrient veins.

The insertion sites for IO infusion commonly used are the humeral head, proximal tibia, distal tibia, and distal femur. Placement can be found under the guideline for IO placement.

Rectal Administration:

The only drug commonly administered via the rectal route in the prehospital setting is valium for status seizure activity in the pediatric patient when IV access is not possible. It is also sometimes used for adult patients who are seizing. Ativan can also be administered rectally.

Procedure for rectal administration of valium (Diazepam):

1. Review 6 rights.
2. Take BSI precautions.
3. Draw up appropriate dose in TB or 1cc syringe.
4. Remove needle.
5. Lubricate tip of syringe.
6. Insert in rectum approximately 3 cm.
7. Inject solution.
8. Facilitate drug retention by elevating and squeezing buttocks together with manual pressure.

Using existing central venous lines and implantable ports for fluid and drug administration:

Central lines may be used for fluid and drug administration in emergency situations. It is important to remember that many patients with central venous lines in place are immunosuppressed or severely debilitated. Thus, they are very susceptible to routine pathogens. Special care should be taken by the paramedic to avoid contamination.

Procedure for using peripheral or central lines for drug or fluid administration:

1. Review 5 rights.
2. Take BSI precautions.
3. Draw up 3 ml of normal saline.
4. Wipe connection port with Betadine and allow it to dry.
5. Connect 5-10ml syringe, release clamp and withdraw 5 ml of blood (do not use this for specimen as it is typically heparinized)
6. Secure clamp
7. Attach syringe with 3ml of normal saline to port, release clamp and flush with saline. Take precautions to insure that you do not flush air from the syringe into the line.
8. Remove syringe and secure clamp
9. You may now connect IV tubing to port (be sure tubing is flushed)
10. Release clamp and adjust flow from drip set.
11. If injecting medication directly into port, be sure to follow with heparin flush and then re-secure the clamp.

(CONTINUED ON NEXT PAGE)

Procedure for using Implantable Ports (Port-a-Cath):

Implantable ports are venous access devices that are surgically implanted under the skin with the distal end of the catheter inserted into a large central vein. The injection end of the catheter is implanted subcutaneously, often on the chest wall, and has a self-sealing septum over a small chamber or reservoir. (Most require Huber needles)

1. Consider 5 rights
2. Take BSI precautions
3. Swab site with Betadine or alcohol
4. Locate the device and stabilize it with one hand.
5. Puncture the skin and septum with a Huber needle attached to a 3 ml syringe containing normal saline. (Huber needles are special stainless-steel needles : they may be straight or angled 90 degrees) Do not use regular needles or IV catheters with Port-a-Caths)
6. Aspirate blood to determine patency and then inject the saline to flush the system.
7. Connect air-free IV tubing to reservoir and begin infusion.
8. Tape connection site to prevent displacement.
9. After use, flush the device with a heparinized solution.

Intranasal Drug Administration

NAD (Nasal Administration Device) or MAD (Mucosal Administration Device).

Some medications can be administered intranasal. If injection or IV is not possible or undesired, this route is an option for the following drugs: Fentanyl, Glucagon, Versed, Narcan and Ketamine. Other drugs may be approved for intranasal administration and will be listed in the drug section.

To administer medication intranasal:

1. Place medication in syringe, add 0.1ml more than the dose needed for dead space.
2. Place the NAD device to syringe to atomize the medication in the nasal mucosa.
3. Have patient sitting up if possible (45 degrees is best)
4. Place the NAD into the nostril aim toward the superior part of the ear) and pinch the other nostril
5. Have patient inhale while spraying
6. Spray ½ of the medication in each nostril. Should not exceed 1 cc per nostril.

PUSH DOSE PRESSERS (PDP)***Procedure:***

1. Gather 10ml ns flush, 1:10,000 cardiac Epi, Twin pack or 18ga. Needle
2. Push out 1ml of NS from 10ml Flush = (9ml of NS)
3. Attach Twin pack (metal point) or 18ga Needle to 9ml Flush
4. Remove top from 1:10,000 Cardiac Epi (Exposing RubberPlunger)
5. Place metal tip or 18ga needle (Attached to 9ml NS Flush) into rubber plunger of 1:10,000 Epi.
6. Draw 1ml from 1:10,000 Epi Into 9ml NS Flush. Making 10ml of 1:100,000 Epi. (10mcg/1ml Epi)
7. Label 1:100,000 Epi with Tape and Marker

Dosing:

- Give 1ml of 1:100,000 Epi IV Slow IVP = (10mcg/ml) Every Minute until Target HR and or Blood Pressure is achieved.
- Monitor very carefully and repeat as needed.

CERTIFICATION REQUIREMENTS

- PARAMEDIC

MIST REPORT

Emergency Department physicians and nursing staff expect concise, precise and pertinent information. The following is the standard format they are most accustomed to. This report, if given in person, should last no more than 30 seconds or so:

M – Mechanism of Injury / Chief Complaint

I – Injuries

S – Signs and Symptoms

T - Treatments

Arkansas Trauma Team Activation Time out – allows 30 seconds for a brief and concise verbal report on trauma patients to the receiving physician and staff. Using MIST will assist in keeping this information brief while still allowing for pertinent information to be given.

Incoming radio report may also follow this format for brevity and consistency.

CERTIFICATION REQUIREMENTS:

- EMR
- EMT
- ADVANCED EMT
- PARAMEDIC

ORTHOSTATIC VITAL SIGNS**INDICATION**

Patient with suspected blood or fluid loss, dehydration or syncope, as a diagnostic aid.

PROCEDURE

1. Assess the need for orthostatic vital signs.
2. Obtain patient's pulse and blood pressure while supine.
3. Have patient stand for one minute.
4. Obtain patient's pulse and blood pressure while standing.
5. If pulse has increased by 20 BPM and systolic BP decreases by 20 mmHG, the orthostatic vital signs are considered positive.
6. If patient is unable to stand, orthostatic vital signs may be taken with patient sitting with feet dangling.
7. Document the vital signs for supine and standing positions.
8. Determine the appropriate treatment based on protocol.

CERTIFICATION REQUIREMENTS:

- EMR
- EMT
- ADVANCED EMT
- PARAMEDIC



Arkansas Department of Health EMS Field Patient Care Report – Short Form
 All Pertinent Sections Should Be Completed for all Patients at Time of Care Transfer to ED Staff
 Please print legibly. Complete this form and leave with the RN receiving the patient.
THIS FORM DOES NOT REPLACE THE OFFICIAL EMS PATIENT CARE REPORT



Agency Name _____ Phone _____ Receiving Hospital _____	
Patient Name _____ DOB ____/____/____ Age _____ Gender _____	
Date ____/____/____ Time ____:____ Level of Consciousness ☐ Alert ☐ Verbal ☐ Pain ☐ Unresponsive	

General / Treatments	Vitals	Time	Blood Pressure	Pulse	BPM	O ₂ Sat	EKG	Airway	
			/					Oxygen	NC NRB BVM
			/					OPA/NPA	☐ Yes ☐ No
	Meds	Time	Medication/Fluid	Dose	Rate	Total Administered	Tube	ETT King Other	
							Size		
							Depth		
							PAI	☐ Yes ☐ No	
	PMHx:			Meds:			Allergies:		
	Primary Impression:								
Findings/Notes (If more room needed, check box at bottom-right and write additional):									

Stroke	Prehospital Stroke Screen Performed: ☐ Yes ☐ No		Stroke Alert called to hospital by EMS: ☐ Yes ☐ No	
	Glucose 60–400 ☐ Yes ☐ No		Last Known Well Time: _____ Date: _____	
	Balance - Sudden loss of balance? ☐ Yes ☐ No		Name providing well time: _____	
	Eyes - Sudden change in vision or trouble seeing? ☐ Yes ☐ No		Phone number: _____ - _____	
	Face - Facial drooping? ☐ Yes ☐ No		Times (approximation - in whole minutes)	
	Arms - Does one arm drift downward?? ☐ Yes ☐ No		Dispatch to patient contact (goal<8):	Minutes
	Speech - Is their speech slurred or strange? ☐ Yes ☐ No		Arrival to first vital set (goal<5):	Minutes
	Time - Did you document last known well time? ☐ Yes ☐ No		Arrival to glucose check (goal<5):	Minutes
	Potential Stroke Patient? ☐ Yes ☐ No		Total time on scene (goal <15):	Minutes
	Stroke Band # - starts with "S" ☒ S		Total transport time:	Minutes

STEMI	12, 15, or 18 Lead Obtained: ☐ Yes ☐ No		STEMI Alert called to hospital by EMS: ☐ Yes ☐ No	
	STEMI Indicated: ☐ Yes ☐ No		Elevation in leads: _____	
	Initial Rhythm: _____		Depression in leads: _____	
	Was ECG transmitted to the receiving facility: ☐ Yes ☐ No			

Trauma	Trauma Band #: _____ (starts with "A" or "B")		ATCC Contacted: ☐ Yes ☐ No		Trauma Alert by EMS: ☐ Yes ☐ No			
	GCS		Eye Opening	Spontaneous To Speech To Pain None 4 3 2 1	Best Verbal Response	Oriented Disoriented Monosyllabic Incomprehensible None 5 4 3 2 1		
	Initial:						Best Motor Response	Obey's Commands Localizes Pain Withdraws from Pain Abnormal Flexion Extension to Pain None 5 4 3 2 1
	Post:							
	MOI: _____							
	Injuries: _____							

EMS Field HCP: Name & Title: _____ Transfer of Care Time: _____

Receiving HCP: Name & Title: _____

☐ If checked, see back side

PHYSICIAN ON SCENE

Primary Physician

A paramedic is permitted to take orders by telephone or other means of communication from the patient's primary/attending physician under any of the following conditions:

- The paramedic knows the physician by voice.
- The physician identifies him or herself and repeats orders to at least two members of the on-scene EMS crew.
- Written, signed orders are presented.

The medical control physician should be notified and this notification documented on the patient care report.

Non-Primary Physician

This pertains only to those situations in which a non-primary physician (i.e. not the patient's physician) is physically present on scene of an emergency. In the event that the physician on scene wishes to direct the care of the patient(s) and therefore, accepts responsibility for the patient(s), the physician on scene must be informed of and agree to the following conditions prior to assuming the care of the patient:

- The physician must show proper identification and a current Arkansas physician's license.
- The physician must agree to sign a written statement attesting to physician's assumption of responsibility for patient care.
- The physician must remain with the patient(s) on scene and during transport to the receiving hospital. Patient care may be transferred at the receiving hospital, with report by the physician, to the medical staff.
- The physician on scene must be informed that the medical control at the receiving hospital will be contacted and medical control will make the final decision regarding assumption of patient care by the physician on scene.

If the above conditions are agreed to, the physician on scene may assume the responsibility for patient care.

Paramedic's Responsibility

1. Remain tactful, calm and courteous.
2. Follow the procedure conditions.
3. Offer assistance to the physician on scene. The paramedic may perform any procedures that are within the scope of practice of that individual as defined by these protocols.
4. Maintain control of medications and equipment.
5. Inform the physician on scene of equipment available.
6. Maintain active communication with medical control.
7. Complete the necessary patient care form and obtain appropriate signatures.

Physician's Responsibility Form

Physicians, please read carefully.

The emergency personnel are trained extensively and function under protocols developed to address situations that occur in the pre-hospital emergency and non-emergency setting.

If you wish to take charge of the injury/illness scene, you must:

1. Show your current Arkansas medical license to the Paramedic or EMT on scene.
2. Agree to take full responsibility for care and treatment of the patient(s) involved in this incident.
3. Accompany the patient(s) in the ambulance to the most appropriate receiving hospital.

Physician's signature

License #, Type

Date

Witness

Date

CERTIFICATION REQUIREMENTS:

- ADVANCED EMT
- PARAMEDIC

PULSE OXIMETRY

INDICATIONS

- Patients with suspected hypoxemia.
- Patients being administered oxygen by EMS crew.
- A trending tool to monitor O₂ saturation as one indication of perfusion to be used in conjunction with other clinical findings.

PROCEDURE

Pulse oximetry measures the percentage of hemoglobin saturated with oxygen and is denoted as SaO₂. Several factors may have an impact on the reading: PCO₂, pH, temperature, CO, and whether hemoglobin is normal or altered. Pulse oximetry changes may be delayed and not a direct reflection of patient's oxygenation. Therefore, clinical findings should determine care of patient. Pulse oximetry should be used as one of those findings along with others to make treatment decisions.

1. Turn the machine on and allow for self-tests.
2. Apply probe to the patient's finger, ear lobe, forehead, or foot.
3. Allow machine to register saturation level. This may take up to 45 seconds.
4. Record saturation percent (SaO₂), pulse rate and time.
5. Verify pulse rate on machine with actual pulse of patient.
6. Monitor critical patients continuously until arrival at the destination.
7. Document percent of oxygen saturation every time vital signs are recorded and in response to efforts to correct hypoxemia.
8. In general, normal saturation is 97—99%. Below 94%, suspect a respiratory compromise.
9. Use the pulse oximetry reading as an added tool for patient evaluation, another clinical finding. Remember to treat the patient, not the machine.
10. Factors which may reduce or otherwise alter reliability of pulse oximetry readings:
 - Poor peripheral circulation—blood volume, hypotension, hypothermia.
 - Low blood hemoglobin concentration.
 - Excessive pulse ox sensor movement.
 - Fingernail polish—should be removed with nail polish remover.
 - Carbon monoxide bound to hemoglobin—250 times greater than oxygen to hemoglobin.
 - Irregular or rapid heart rhythms—atrial fibrillation, SVT, etc.
 - Jaundice.

CERTIFICATION REQUIREMENTS

- EMR
- EMT
- ADVANCED EMT
- PARAMEDIC

RESTRAINTS

INDICATION


Patients with actual or potential threat to self or others.

PROCEDURE

1. Evaluate the need for restraints. Restraints should be considered only as a last resort after verbal techniques have failed.
2. If threat to self or others is due to behavioral problems (including drugs or alcohol) or criminal behavior (including resisting arrest), request law enforcement assistance.
3. Consult Medical Control.
4. The least amount of restraint necessary to accomplish the desired purpose should be used.
5. The restraints should not be limiting to the patients peripheral or central circulation or respiratory status.
6. Soft restraints such as cravats or roller bandage can be used for extremity restraints. Sheets may be used to limit upper body or lower extremity movement. This does not restrict the use of equipment specifically designed for patient restraint.
7. Restraints should be frequently monitored during transport. Neurovascular status of restrained parts should be assessed.
8. Documentation should include the reason for the use of restraints, the type of restraints used, and the time restraints were placed.

CERTIFICATION REQUIREMENTS

- EMT
- ADVANCED EMT
- PARAMEDIC

RULE OF NINES**CERTIFICATION REQUIREMENTS:**

- EMR
- EMT
- ADVANCED EMT
- PARAMEDIC

SAFETY EQUIPMENT REMOVAL

INDICATION

Mechanism of injury or signs and symptoms that suggest potential spinal injury.

PROCEDURE

1. There are many different athletic events where the potential for spinal injury is high. Some athletes wear protective equipment which may vary from sport to sport, level to level, or school to school. Individuals may also wear helmets or padding for non-sport related activity such as riding a motorcycle.
 2. In sports situations, the paramedic must coordinate activities with the team's athletic trainer or physician when possible.
 3. Removal of helmets and padding is recommended for assessment as well as for maintaining the spine in a neutral in-line position.
 4. Removal of helmet and shoulder pads:
 - o Helmet Removal:
 - a. Manually immobilize the helmet.
 - b. Remove any face mask or face shield as able.
 - c. A second rescuer should provide anterior and posterior, or lateral, stabilization and support of the patient's head and neck during removal of the helmet.
 - d. Deflate any air bladders in the helmet.
 - e. Remove any cheek/jaw pad if necessary.
 - f. Gently slide the helmet off rotating the helmet slightly anterior so as not to lift the occiput. Do not pull apart from side to side unless absolutely necessary for removal.
 - g. Be sure to maintain neutral alignment of the cervical spine once the helmet is removed as well as during removal of the shoulder pads.
 - o Shoulder pad removal:
 - a. Expose the anterior portion of pads and cut any center strings/straps.
 - b. Cut any straps under the arms. (These straps can be disconnected if doing so does not cause excessive movement.)
 - c. Maintain stabilization and support of the head and neck with hands underneath the shoulder pads from side of the patient's torso.
 - d. With the appropriate number of rescuers, lift the patient's thorax, maintaining neutral alignment just enough to gently slide the shoulder pads off.
- OR**
- e. Gently slide the shoulder pads off over the patient's head if the surface the patient is on allows for smooth sliding.
5. Several sports require many different types of protective equipment. Approach each suspected spinal injury with the goal of maintaining neutral alignment.
 6. Complete a thorough neurological assessment prior to, and after, spinal restriction.

CERTIFICATION REQUIREMENTS:

EMT

ADVANCED EMT

PARAMEDIC

STROKE SCREEN BEFAST

 STROKE SCALE			
B	BALANCE	Trouble walking, dizzy, loss of coordination	
E	EYES	Trouble seeing, blurred vision, loss of vision	
F	FACE	Droop, numbness, unequal	
A	ARMS	Paralysis, numbness, unequal strength, weakness	
S	SPEECH	Slurred, loss, unable to find right words	
T	TIME	Document time last seen normal AND time symptoms noted	

CERTIFICATION REQUIREMENTS:

- EMR
- EMT
- ADVANCED EMT
- PARAMEDIC

TOXICOLOGICAL SYNDROMES

COMMON SIGNS	CAUSATIVE AGENT	SPECIFIC TREATMENT RECOMMENDATIONS
Cholinergic (“Wet” patient presentation) Confusion, CNS depression, weakness, SLUDGE (salivation, lacrimation, urination, defecation, emesis), bradycardia, wheezing, bronchoconstriction, miosis, coma, convulsion, diaphoresis, seizure	Organophosphate and Carbamate insecticides, nerve agents, some mushrooms	Atropine, pralidoxine (2-PAM Chloride), diazepam, activated charcoal
Anticholinergic (“Dry” Patient Presentation) Delirium, tachycardia, dry flushed skin, dilated pupils, seizures and dysrhythmias (in severe cases)	Antihistamines, antiparkinson medications, atropine, antipsychotic agents, antidepressants, skeletal muscle relaxants, many plants (e.g., jimson weed, and Amanita muscaria)	Benzodiazepine, activated charcoal, rarely physostigmine (Antilirium)
Hallucinogen Visual illusions, delusions, bizarre behavior, flashbacks, respiratory and CNS depression	LSD, PCP, mescaline, some mushrooms, marijuana, jimson weed, nutmeg, mace, some amphetamines	Minimal sensory stimulation and calming measures, benzodiazepine if necessary
Opioids Euphoria, hypotension, respiratory depression, arrest, nausea, pinpoint pupils, seizures, coma	Heroin, morphine, codeine, meperidine (Demerol), propoxyphene (Darvon), fentanyl (Duragesic), OxyContin	Naloxone (Narcan), nalmeferene (Revex)
Sympathomimetic Delusions, paranoia, tachycardia or bradycardia, hypertension, diaphoresis, seizures, hypotension and dysrhythmias in severe cases	Cocaine, amphetamine, methamphetamine, over-the-counter decongestants	Minimal sensory stimulation and calming measures, benzodiazepine if necessary

TRAUMA ALERT

The following reflects the Pre-hospital Triage and Decision Scheme of the ADH Rules and Regulations for Trauma Systems, March 2000. All trauma patients shall be evaluated against the criteria to determine the need for rapid transport. If the trauma patient meets any of the items listed below consider the patient a **"trauma alert"** and notify dispatch and ATCC as soon as possible. The dispatch center shall notify the receiving facility immediately and document the trauma alert time. On-scene times for patients meeting the trauma alert criteria shall be 10 minutes or less, unless there are extrication delays. Transport of the **"trauma alert"** patient to the receiving facility shall be in the emergency mode, unless otherwise determined by Medical Control.

VITAL SIGNS & LEVEL OF CONSCIOUSNESS

- Shock Systolic Blood Pressure of **90** mmHg or less with other signs & symptoms of shock
- Respiratory Respiratory Rate of **10** or less; or **29** or higher.
Distress Stridor or retractions.
- Altered Mentation Glasgow Coma Scale of **13** or less
 Pediatric Coma Scale of **9** or less
 Trauma Score of **11** or less
 Pediatric Trauma Score of **9** or less

ASSESS ANATOMY OF INJURY

- Penetrating injury to the head/open or depressed skull fracture
- Penetrating injury of the neck, torso, or groin
- Amputation above the wrist or ankle
- Spinal cord injury with limb paralysis or alteration of SMC's
- Flail chest
- Pelvic fracture
- Two or more obvious long bone fractures above the elbows or knees
- Major burns: 15%BSA or greater and/or with respiratory involvement
- High voltage electrical burns

For trauma patients meeting any one of the above criteria, initiate Trauma Alert and Rapid Transport

For Trauma patients not meeting any one of the above criteria, consider the following to determine the need for TRAUMA ALERT and rapid transport. Consult Medical Control and ATCC for assistance if necessary.

MECHANISM OF INJURY

- Speed 40 mph or greater
- Vehicle rollover
- Death of same vehicle occupant
- Pedestrian vs. vehicle 5mph or greater
- Vehicle deformity 20" or greater
- Ejection from moving vehicle
- Motorcycle, ATV or bicycle 20mph or greater
- Falls 20ft or greater (consider pediatric rules if applicable)

CO-MORBID FACTORS

The following factors may compound the severity of injury and shall increase the index of suspicion:

- Extremes in age: 12 or less/55 or more
- Hostile environment (e.g. extremes of heat or cold)
- Medical illness (e.g. COPD, CHF, renal failure)
- Presence of intoxicants/substance abuse
- Pregnancy

CERTIFICATION REQUIREMENTS:

- PARAMEDIC

TRAUMA SCORE ADULT

Respiratory Rate		Respiratory Expansion		Systolic Blood Pressure	
10–24/ minute	4	Normal	1	90 mmHg or greater	4
25–35/ minute	3	Retractive	0	70–89 mmHg	3
36/ minute or greater	2			50–69 mmHg	2
1–9/ minute	1			0–49 mmHg	1
None	0			None	0

Capillary Refill			Add points for Glasgow Coma Score	
Normal	2	Nail bed, forehead, or lip color refill (less than) < 2 seconds	14–15	5
Delayed	1	> 2 seconds	11–13	4
None	0	No capillary refill	8–10	3
			5–7	2
			3–4	1

Trauma Score: ____ + Points for GCS: ____ = TOTAL Trauma Score (revised): _____

CERTIFICATION REQUIREMENTS:

- EMR
- EMT
- ADVANCED EMT
- PARAMEDIC

TRIAGE

INDICATION

Multiple-patient scenarios, to categorize patients based on the severity of their injuries, prioritize their need for treatment and transportation and stabilize life-threatening injuries before additional resources arrive on-scene.

PROCEDURE

This procedure is based on START triage system.

1. Determine the location, number and condition of patients.
2. Determine, in close coordination with Extrication sector, if triage will be performed in place or at the entrance to the treatment area.
3. Determine resources.
4. Assign triage teams.
5. Direct minor patients (walking wounded) to a gathering place and tag them later. *
6. Identify and treat as necessary, remaining patients.
 - Evaluate patient using START
 - Attach triage tag or ribbon to patient
7. When triage is complete, provide COMMAND with a “Triage Report.”
8. Once “Immediate” have been treated/transported, Reassess “Delayed” by Mechanism of Injury and upgrade as necessary. May be done continuously if resources allow *

*At smaller incidents (up to 10 patients) “MINOR” patients should not be relocated and reassessment should be continuous

The S.T.A.R.T. Algorithm

<u>ACTION</u>	<u>Tagged as</u>
Move the walking wounded	MINOR
No respirations (after head tilt or insertion of an OPA)	DEAD/DYING
Respiration over 30	IMMEDIATE
Pulse—No radial pulse	IMMEDIATE
Mental Status—Unable to follow simple commands	IMMEDIATE
All others	DELAYED

CERTIFICATION REQUIREMENTS:

- EMT
- ADVANCED EMT
- PARAMEDIC